

Wuhan Institute of Virology analysis of bronchial lavage specimens from ICU patients at Wuhan Jinyintan Hospital in December 2019 contain both SARS-CoV-2 and adenovirus vaccine sequences, consistent with a vaccine challenge trial

By

Steven Carl Quay, MD, PhD

www.DrQuay.com

Steven@DrQuay.com

Dedication. This work is dedicated to the men, women, and children who were infected with SARS-CoV-2 over the last year. It is my hope that this work becomes part of the body of evidence to help inform the public about gain-of-function pathogen research and that a renewed debate can be had about the benefits and risks of this research in the context of world health.

COVID-19 CORONAVIRUS / CASES

[UPDATES](#) - [Graphs](#) - [Countries](#) - [Death Rate](#) - [Incubation](#) - [Age](#) - [Symptoms](#)

Last updated: January 26, 2021, 23:49 GMT

Coronavirus Cases

There are currently **100,789,048 confirmed cases** and **2,164,336 deaths** from the coronavirus COVID-19 outbreak as of January 26, 2021, 23:49 GMT.

Source: <https://www.worldometers.info/coronavirus/>

Acknowledgements. Despite having collaborated over many decades on numerous scientific projects, research during 2020 into COVID-19, SARS-CoV-2, and therapeutic approaches has been a unique experience. With lockdowns and international travel bans, all collaborative work has been virtual. With an apparent bias surrounding investigation into the origin of SARS-CoV-2, ad hoc groups of Citizen-Scientists, often anonymous, have worked together via email, videoconference, micro-blogging, and social messaging networks to advance our understanding of this horrific pandemic.

I want to thank a Twitter group called #DRASTIC for many useful discussions that found their way into this document. Dr. Martin Lee, Ph.D., Adjunct Professor of Statistics at UCLA provided statistical support throughout this work. H. Lawrence Remmel provided input on the adenovirus vaccine as a dual target vaccine. I want to thank D.A. for originally suggesting performing a Bayesian analysis on the work I had done on SARS-CoV-2 and for his facilitation of the review of this work by a diverse group of scientists and policy makers.

In all cases, however, this is my own work product.

A Bayesian analysis concludes beyond a reasonable doubt that SARS-CoV-2 is not a natural zoonosis but instead is laboratory derived

Wuhan Institute of Virology analysis of lavage specimens from ICU patients at Wuhan Jinyintan Hospital in December 2019 contain both SARS-CoV-2 and adenovirus vaccine sequences consistent with a vaccine challenge trial

Executive Summary. The one-year anniversary of the COVID-19 pandemic records 2.1 million deaths, over 100 million confirmed cases,¹ and trillions of dollars of economic damage. Although there is universal agreement that a coronavirus identified as Severe Acute Respiratory Syndrome Coronavirus 2 or SARS-CoV-2 (abbreviated CoV-2 henceforth) causes the disease COVID-19, there is no understanding or consensus on the origin of the disease.

The Chinese government, WHO, media, and many academic virologists have stated with strong conviction that the coronavirus came from nature, either directly from bats or indirectly from bats through another species. Transmission of a virus from animals to humans is called a zoonosis.

A small but growing number of scientists have considered another hypothesis: that an ancestral bat coronavirus was collected in the wild, genetically manipulated in a laboratory to make it more infectious, training it to infect human cells, and ultimately released, probably by accident, in Wuhan, China. For most of 2020 this hypothesis was considered a crackpot idea, but in the last few weeks, more media attention has been given to the possibility that the Wuhan Institute of Virology, located near the Wuhan city center and with a population of over 11 million inhabitants, may have been the source of the field specimen collection effort, laboratory genetic manipulation, and subsequent leak. On January 15, 2021, the U.S. Department of State issued a statement requesting the WHO investigation of the origin of COVID-19 include specific assertions related to a laboratory origin of the pandemic.²

Given the strong sentiment in the scientific community in favor of a zoonosis and the massive effort undertaken by China to find the natural animal source, one can assume that any evidence in favor of a natural origin, no matter how trivial, would become widely disseminated and known. This provides a potential evidence bias within the scientific community in favor of a natural origin which isn't quantifiable but should be kept in mind.

This becomes especially important background when evidence that could support a laboratory origin has been directly provided by leading Chinese scientists themselves, like Dr. Zhengli Shi, head of coronavirus research at the Wuhan Institute of Virology and Gao Fu (George Fu Gao), Director of Chinese CDC; by the Chinese government, as well as by powerful and vocal, pro-natural origin scientists, like Dr. Peter Daszak, of the NYC-based NGO, EcoHealth Alliance.

¹ <https://www.worldometers.info/coronavirus/>


² <https://www.state.gov/ensuring-a-transparent-thorough-investigation-of-covid-19s-origin/>

This report uses Bayesian inference, a common statistical tool in which Bayes' theorem, a well-known statistical equation, is used to update the likelihood for a particular hypothesis as more evidence or information becomes available. It is widely used in the sciences and medicine and has begun to be used in the law.

The starting probability for origin of SARS-CoV-2 was set with the zoonotic or natural hypothesis at 98.8% likelihood with the laboratory origin hypothesis set at 1.2%. The initial state was biased as much as possible towards a zoonotic origin, with the starting point selected as the upper bounds of the 95% confidence interval for the mean and standard deviation of three independent estimates, including one by Daszak and colleagues. Each piece of new evidence for or against each hypothesis was then used to adjust the probabilities. If evidence favored a natural origin the math adjusts upward the probability of a natural origin, and so on.

The most significant evidence provided herein is the finding from RNA-Seq performed by the Wuhan Institute of Virology (WIV) of lavage patient samples collected on December 30, 2019.³ These ICU patients were the subject of the seminal paper, entitled, “A pneumonia outbreak associated with a new coronavirus of probable bat origin,” from Dr. Zhengli Shi and colleagues that first characterized SARS-CoV-2.⁴ This author has confirmed that the RNA-Seq of all five patients contained SARS-CoV-2 sequences.

Surprisingly the specimens also contained the adenovirus “pShuttle” vector, developed by Chinese scientists in 2005 for SARS-CoV-1.⁵ Two immunogens were identified, the Spike Protein gene of SARS-CoV-2 and the synthetic construct H7N9 HA gene.⁶ Hundreds of perfectly homologous (150/150) raw reads suggest this is not an artefact. Reads that cross the vector-immunogen junction are identified. An example of the read contigs for CoV-2 is shown in this figure:



³ The detailed evidence for the adenovirus vaccine sequences is given at the end of this document.

⁴ <https://www.nature.com/articles/s41586-020-2012-7>

⁵ <https://www.ncbi.nlm.nih.gov/nuccore/AY862402.1>

⁶ <https://www.ncbi.nlm.nih.gov/nuccore/KY199425.1/>

While adenovirus is a common infection the wildtype viruses have low homology to the vaccine vector sequence, by design, to avoid rejection of the vaccine due to prior exposure to wildtype adenoviruses.

Two patients from the same hospital who had bronchial lavage on the same day but had their specimens sent to the Hubei CDC did not have adenovirus vaccine sequences.

Three explanations come to mind from this evidence:

1. These represent sample preparation artifacts at the WIV, such as sample spillover on the sequencer.
2. These patients were admitted with an unknown infection, were not responding to the treatment protocols for a infection of unknown origin, and they were vaccinated with an experimental vaccine in a desperate but compassionate therapeutic “Hail Mary.”
3. A clinical trial of a combination influenza/SARS-CoV-2 vaccine was being conducted and an accidental release into Wuhan occurred.

Only WIV scientists and Chinese authorities can answer these questions. Until the evidence of the adenovirus sequences has been confirmed by other scientists, this author will not include this evidence in the Bayesian analysis.

Obviously if a vaccine containing the Spike Protein of SARS-CoV-2 was being administered to patients in Wuhan in December 2019 the question of laboratory origin is a settled matter.

The remaining analysis is being conducted without the adenovirus vaccine evidence unless and until it is corroborated. The outcome of this report is the conclusion that the probability of a laboratory origin for CoV-2 is 99.8% with a corresponding probability of a zoonotic origin of 0.2%. This exceeds most academic law school discussions of how to quantify ‘beyond a reasonable doubt,’ the threshold for finding guilt in a criminal case. The report contains the detailed analysis and quantitative basis for the statistics and conclusion. It should be noted that because of the commutative property of the collected adjustments to the probabilities, the order in which they are used in the overall calculation is immaterial and the same end likelihoods will be reached regardless of the order of input.

The following Text-Table summarizes the evidence examined and the changes in probabilities:

Evidence	Zoonotic Origin	Laboratory Origin
Initial State	98.8%	1.2%
International committees to determine CoV-2 origin may not be impartial	98.8%	1.2%
Three key zoonotic papers: pros and cons	98.8%	1.2%
SARS-like infections among employees of the Wuhan Institute of Virology in the fall of 2019 reported by US Government	98.8%	1.2%
Location of first cases near Wuhan Institute of Virology	95.1%	4.9%
Lack of evidence of seroconversion in Wuhan and Shanghai	80.9%	19.1%
Lack of posterior diversity	30.8%	69.2%
Opportunity: The Wuhan Institute of Virology has publicly disclosed that by 2017 it had developed the techniques to collect novel coronaviruses, systematically modify the receptor binding domain to improve binding or alter zoonotic tropism and transmission, insert a furin site to permit human cell infection, make chimera and synthetic viruses, perform experiments in humanized mice, and optimize the ORF8 gene to increase human cell death.	30.8%	69.2%
Lack of furin cleavage sites in any other sarbecovirus	4.7%	95.3%
Rare usage of -CGG- single codons & no CGG-CGG pairs	0.5%	99.5%
Routine use of CGG in laboratory codon optimization, including Daszak & Shi	0.2%	99.8%
Spike Protein receptor binding region (200 amino acids) optimized for humans	0.2%	99.8%
Whole genome analysis shows pre-adaption of CoV-2	0.2%	99.8%
The finding of CoV-2 in Barcelona wastewater in early 2019 was an artifact	0.2%	99.8%
Shi and the WHO comment early on that CoV-2 seemed to begin with a single patient	0.2%	99.8%
Mammalian biodiversity between Yunnan and Hubei is significantly different, limiting a potential common intermediate host	0.2%	99.8%
The ancestor of CoV-2 can only obtain a furin site from other subgenera viruses but recombination is limited/non-existent between subgenera	0.2%	99.8%
Canvas of 410 animals shows humans and primates are the best, bats are the worst, for ACE2-Spike Protein interaction	0.2%	99.8%
A government requested review of samples collected from a mineshaft may have caused the COVID-19 pandemic	0.2%	99.8%
The Hunan Seafood Market and farmed animals in Hubei province are not the source of CoV-2	0.2%	99.8%
Line 2 of the Wuhan Metro System is the likely conduit of the pandemic and is the closest subway line to the WIV	0.2%	99.8%
Feral and domestic cats are not the intermediate host	0.2%	99.8%
Extraordinary pre-adaption for the use of human tRNA is observed	0.2%	99.8%
Evidence of lax operations and disregard of laboratory safety protocols and regulations in China	0.2%	99.8%
Previous SARS-CoV-1 laboratory accidents	0.2%	99.8%
Shi and Daszak use Wuhan residents as negative control for zoonotic coronavirus exposure	0.2%	99.8%
RaTG13 could be CoV-2 precursor using the synthetic biology 'No See 'Em' technique	0.2%	99.8%
Location, location, location: Based on the distance between known SARS-CoV-1 laboratory-acquired infections and the hospital of admission of the infected personnel, the WIV is within the expected hospital catchment for a CoV-2 LAI	0.2%	99.8%

The summary which follows will simply be a review and discussion of the evidence in the context of the two hypotheses.

Zoonosis Hypothesis

A viral zoonosis has at least three elements, a host, a virus, and the human population. With some viruses there are often two hosts. One is a 'reservoir host' where the virus can live for years or even decades in a relatively stable relationship. The reservoir host is never decimated by the virus, and the virus is never burned out by the reservoir host, disappearing completely. For coronaviruses the reservoir host is always one or more bat species. If there is a reservoir host that some viruses that cannot jump directly into the human population, there is a need for an second host, an intermediate host. In this case the virus spends time jumping into the intermediate host, 'practicing' adaption through random mutation and Darwinian selection for fitness to reproduce, infect, and transmit in the intermediate host. This process is then repeated between the intermediate host and the human population. Alternatively, the virus can jump directly between the bat reservoir and humans, without the need for an intermediate host.

For two prior human coronavirus epidemics, an intermediate or proximate host was identified. For SARS-CoV-1 in 2003-4 it was the civet cat while for Middle Eastern Respiratory Syndrome (MERS) in 2012-4 it was the camel. In both of these human epidemics, the intermediate host was identified within four to ten months of the first clinically identified human infection. With CoV-2 we are at 12 months since the pandemic began and still waiting for evidence of, despite a much larger effort inside China to find an intermediate host. For both of these previous pandemics, a bat species reservoir host was also identified, but not in the case of SARS-CoV-2.⁷

Based on the genome sequence of CoV-2, Drs. Shi and Daszak have proposed that the reservoir host for CoV-2 is the intermediate horseshoe bat (*Rhinolophus affinis*), which is found in Yunnan Province. Yunnan Province is in southern, rural China and about 1900 km from the north central province of Hubei, where the 11 million people of Wuhan live. In the US this would be equivalent in distance, climate change, and human population density difference to going from the Everglades in Florida to Manhattan, in New York City. The intermediate horseshoe bat isn't found at all in Hubei province, making a direct bat-to-human transmission improbable.⁸ Experiments in three independent laboratories also demonstrate that CoV-2 has changed genetically so much that it can no longer infect any bat species cell culture tested. So, while the leading US coronavirus expert, Dr. Ralph Baric of The University of North Carolina suggested in early 2020 that CoV-2 may have jumped into the human population directly from bats without an intermediate host, this hypothesis seems to no longer be viable.

For the zoonosis hypothesis to be advanced, it is now necessary to find an intermediate host. In January 2020 a theory was proposed that CoV-2 arose in the Huanan Seafood Market, a traditional Chinese “wet market” where live animals are butchered and sold for food. The market theory was based on the observation that about 40% of early patients worked or shopped there. This was reminiscent of the wet market sources for civet cats infected with SARS-CoV-1 or the camel markets for the MERS coronavirus. The Chinese authorities closed the market on December 31, 2019 after performing extensive environmental sampling and sanitation.

But by May 2020 Dr. Gao Fu, Director of the Chinese CDC, announced that the market was not the source of CoV-2, as all of the animal specimens tested negative for CoV-2. And while SARS-CoV-1 was found in 100% of local farmed civets when tested, CoV-2 was different. In July 2020 Dr. Shi reported that extensive testing of farmed animals throughout Hubei Province failed to find CoV-2 in any animals.

For about six months, the pangolin, a scaly anteater, was suspected to be the intermediate host but finally Dr. Daszak reported that CoV-2 was not found in pangolins in the wild or from the (illegal) market trade.⁹ Domestic and feral cats also were ruled out as a possible source. A

⁷ I am distinguishing here the difference between SARS-CoV-2 being a descendent of a bat coronavirus (with 3.8% or 1100 nucleotide (nt) differences between them) and the finding of the immediate precursor of SARS-CoV-2 in a bat colony population somewhere in the wild, which usually is <100 nt differences.

⁸ “We have done bat virus surveillance in Hubei Province for many years but have not found that bats in Wuhan or even the wider Hubei Province carry any coronaviruses that are closely related to SARS-CoV-2. I don't think the spillover from bats to humans occurred in Wuhan or in Hubei Province,” said Dr. Shi. [Science, July 2020](https://doi.org/10.1126/science.abc.2020.01.03.328582)

⁹ <https://link.springer.com/article/10.1007/s10393-020-01503-x>

comprehensive computer-based screen of 410 different animals reported the remarkable finding that the best ACE2 receptor matches to CoV-2 were human and other primates (or primate cells in the laboratory), including the favorite laboratory coronavirus host, the VERO monkey cell culture, and that all bat species were the worst host. At the time of this writing, there is not even a working hypothesis for the species of an intermediate host.

A typical zoonosis has a number of characteristic properties that can allow identification of a zoonotic infection, even in the absence of identifying an intermediate host. None of these properties are found for CoV-2.

All zoonotic infections have in common the principle that when a virus in nature uses evolution to move from, for example, a bat host to a camel host and then to a human host, it is a hit and miss, slow process. After all, evolution is the result of random genetic changes, mutations, and then enrichment of the ones that are helpful by amplification during reproduction. With both SARS-CoV-1 and MERS, the coronavirus spent months and years jumping from the intermediate host into humans, not having all of the necessary mutations needed to be aggressive, grow, and then spread, but spending enough time in humans to cause an infection and leaving behind a corresponding immune response.

The hallmark evidence of this ‘practice’ in abortive host jumping is in stored, archived human blood specimens taken from before the epidemic, where one can find evidence of pre-epidemic, usually sub-clinical, community spread from the antibodies to the eventual epidemic virus. For SARS-CoV-1 and MERS, about 0.6% of people in the region where the epidemic began showed signs of an infection in archived blood. With CoV-2, this seroconversion, as it is called, has never been observed, including in 540 specimens collected from ‘fever clinics’ in Wuhan between October 2019 and January 2020, reported by the WHO. Because this is such a potent signal of a zoonosis, and because I believe that China has over 100,000 stored specimens from Wuhan taken in the fall of 2019, the lack of reports of seroconversion, the silence from China on this evidence, speaks volumes.

Another hallmark of a slow, natural zoonosis can be found in the virus. In SARS-CoV-1 and MERS, the coronavirus spent years in the intermediate host, passing back and forth among populations of hosts, the civets or camels, that were living in close proximity. During this time, they would accumulate a background of genetic mistakes, i.e., mutations- usually about one mistake every two weeks. When the final chip falls, and a mutation(s) happens allowing the jump into humans, the virus with that new mutation(s) also jumps around within the intermediate host population. The consequence of this latter behavior for a true zoonosis is that the genome sequences found in humans don’t all descend from a single jump into a single human but show jumps from viruses that are only cousins of each other, not direct lineal descendants.

In a true zoonosis, the family tree of virus genome sequences doesn’t pass back through the first patient but instead tracks all the way back to an ancestor months or years earlier. This is called posterior diversity, and it is an easy genetic test to perform. With CoV-2, every one of the more than 294,000 virus genomes sequenced can be traced back to the first genomic cluster and in the first patient in that cluster, a 39-year-old man who was seen at the People’s Liberation Army

(PLA) Hospital about one mile from the Wuhan Institute of Virology. The CoV-2 pandemic has the phylogenetic signature of one pure virus sequence infecting one human, with human-to-human spread thereafter; there is just the one and only jump into the human population ever seen. This lack of posterior diversity has been alluded to by Dr. Shi, by the WHO, and by other prominent virologists; they just never take that critical piece of the evidence to the next the proper inference.

The virus in a true zoonosis also contains the signature record of the gradual changes and adaptations it made in the protein key, the Spike Protein, it uses to unlock human cells and cause infection. With SARS-CoV-1 the Spike Protein had fewer than one-third of all the changes it would later develop by the time it became an epidemic. With CoV-2 the Spike Protein was almost perfectly adapted to the human lock, using 99.5% of the best amino acids possible.

Since with CoV-2 we have no evidence from stored blood that it was quietly practicing on humans in the community of Wuhan, it is surprising that when it finds its first patient, it has perfected to 99.5% the spike protein amino acid sequence, its ability to attack and infect humans. If this adaption couldn't have happened in the community, the only place it could have happened is in a laboratory, by what is called serial passage, a common laboratory process that repeatedly gives the virus a chance to practice on humanized mice or VERO monkey cells.¹⁰ A related study showing human adaption right from the start of the pandemic looked at which of the dozens of protein manufacturing tools that CoV-2 uses (called tRNAs). It showed the same uncanny adaptation to the human tools with no evidence that the tools from other potential intermediate hosts would be suitable.

This evidence presented makes a strong case that CoV-2 did not come from nature. But is there affirmative evidence that it could have come from a laboratory? The answer is yes.

Laboratory Origin Hypothesis

The spike protein that gives the coronavirus its name, corona or crown, is the key to match with the lock found in host cells. But before it can inject its genetic material in the host cell, the spike protein needs to be cut, to loosen it in preparation for infection. The host cell has the scissors or enzymes that do the cutting. The singular, unique feature of CoV-2 is that it requires a host enzyme called furin to activate it at a spot called the S1/S2 junction. No other coronavirus in the same subgenera has a furin cleavage site, as it is called. The other coronaviruses are cleaved at a site downstream from the S1/S2 site, called the S' site.

This is of course a major problem for the zoonosis theory, but it gets worse.

Since 1992 the virology community has known that the one sure way to make a virus deadlier is to give it a furin cleavage site at the S1/S2 junction in the laboratory. At least eleven gain-of-function experiments, adding a furin site to make a virus more infective, are published in the open literature, including Dr. Zhengli Shi, head of coronavirus research at the WIV. This has

¹⁰ It is noteworthy that the furin cleavage site is actually unstable in passage in VERO cells and is often deleted within a few passages. A laboratory origin theory needs to account for this observation. On the other hand, mutations in the furin site among the human CoV-2 genomes are exceedingly rare.

caused a flurry of Chinese papers since the pandemic began trying to show a natural furin site in a related virus (this one example was later shown to be an error in interpretation) or to show that furin sites from distant cousins of CoV-2 might be the source through a process called recombination, where two different viruses infect the same host and then make a mistake in copying their genetic material, and swap sequences.

These convoluted, hypothetical methods each fail, however. It turns out that it is Daszak himself who has shown that the subgenera of coronaviruses that have furin sites are found in different bat hosts, which live in different regions of China, than the sarbecovirus subgenera of which CoV-2 is a member. And even with these barriers, they apparently are too far apart to recombine. “For the three focal subgenera, *Sarbecoviruses*, *Merbecoviruses* and *Embecoviruses*...none of the three focal subgenera recombines with one another.”¹¹ As noted previously² Dr. Shi also does not believe the bats of Hubei province are capable of being a host for CoV-2-related coronaviruses.

But it gets worse still for the zoonosis theory. The gene sequence for the amino acids in the furin site in CoV-2 uses a very rare set of two codons, three letter words so six letters in a row, that are rarely used individually and have never been seen together in tandem in any coronaviruses in nature. But these same ‘rare in nature’ codons turn out to be the very ones that are always used by scientists in the laboratory when researchers want to add the amino acid arginine, the ones that are found in the furin site. When scientists add a dimer of arginine codons to a coronavirus, they invariably use the word, CGG-CGG, but coronaviruses in nature rarely (<1%) use this codon pair. For example, in the 580,000 codons of 58 Sarbecoviruses the only CGG pair is CoV-2; none of the other 57 sarbecoviruses have such a pair.¹²

So, there is no natural example of a furin protein site in nature that could be introduced into CoV-2 by recombination, there is no natural example of the particular gene sequence for the furin protein site contained in CoV-2 being used to code for anything in nature, but this particular coding is exactly what Dr. Shi, Baric, and others have used previously in published experiments to insert or optimize arginine codons.

It is telling that when Dr. Shi introduced the world to CoV-2 for the first time in January 2020 she showed hundreds of gene sequences of this novel virus but stopped just short of showing the furin site, the one she is purported to have introduced, seemingly not wanting to call attention to her handywork. She apparently failed to realize that an accomplished but innocent virologist, finding the first furin site ever seen in this class of viruses apparently coming from nature, would have featured the presence of the furin site prominently, and also would have used its presence and her experience with furin sites in other viruses to predict what it would foretell for the world due to its aggressive nature.

She could have perhaps saved many lives just by telling the world that she saw a furin site in the virus sequence. It would be left to a French and Canadian team to later identify the furin site in a

¹¹ CoV-2 is in the subgenera Sarbecoviruses.

<https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1009272>

¹² <https://virological.org/t/alignment-of-58-sarbecovirus-genomes-for-conservation-analysis-of-sars-cov-2/430>

paper.¹³ They would write: “This furin-like cleavage site...may provide a **gain-of-function** to the 2019-nCoV for efficient spreading in the human population compared to other lineage B betacoronaviruses.” [Emphasis added.]

Dr. Shi has denied the virus came from her lab, but she has created such a record of multiple examples of obfuscation, half-truths, contrived specimens, genetic sequences taken from thin air but published in premier journals and US NIH databases, etc. that her veracity is deeply damaged. Perhaps her words and actions on December 30, 2019 show the truth. Her very first response when told there was an unknown outbreak in Wuhan and to return back quickly from a meeting she was attending in Shanghai was to say, “Could this have come from our lab?”¹⁴

“I wondered if [the municipal health authority] got it wrong,” she says. “I had never expected this kind of thing to happen in Wuhan, in central China.” Her studies had shown that the southern, subtropical provinces of Guangdong, Guangxi and Yunnan have the greatest risk of coronaviruses jumping to humans from animals—particularly bats, a known reservoir. After all, the US equivalent of the distance, climate change, and human population density change between Yunnan and Wuhan is comparing the Everglades National Park in Florida and New York City.

Her other action on December 30 was to alter WIV computer databases of novel coronaviruses used by the world’s virologists for research to make it more difficult to search for which coronaviruses she had in her building. In short, the day she was asked to address the pandemic in Wuhan, she chose to spend time to make unavailable to her fellow scientists of the world her decades of coronavirus work.

The notion that CoV-2 was a laboratory creation, designed for maximum virulence, that escaped the laboratory accidentally has additional rings of evidence. From President Xi announcing in February new laws about laboratory security, to abundant evidence that the WIV was closed in October with few personnel inside, to the top military medical research doctor, General Chen Wei, being placed in charge of the WIV, to many more clues, it is clear an event occurred in Wuhan sometime in late 2019 that is most consistent with a laboratory escape.

The Asian region has a two-decade record of a little less than one laboratory-acquired infection per year. After the first SARS-CoV-1 epidemic was ended, SARS-CoV-1 jumped four more times into the human population, all from laboratories, with two in China. The last smallpox death in the entire world was a secretary who worked two floors above a research lab in England and contracted it through the ventilation system. The head of that laboratory committed suicide over his anguish for causing her death.

Over and over again, there is a long history and record of laboratory acquired infections that provides the background for considering what happened here.

¹³ <https://www.sciencedirect.com/science/article/pii/S0166354220300528?via%3Dihub>

¹⁴ https://www.scientificamerican.com/index.cfm/_api/render/file/?method=inline&fileID=E1FDF8DE-9E22-4CE5-AD8B2E4682F52A86

Lab-made Bio-Weapon Hypothesis

But was SARS-CoV-2 more than just a gain-of-function experiment that escaped a laboratory? Could it have been one part of a two-part novel virus-vaccine bioweapons program?

General Chen Wei has been involved in vaccine research since joining the People's Liberation Army after college. In a 2017 internal speech at the AMMS (Academy of Military Medical Sciences) she said: "只要有矛. 才能研究盾." which translates roughly as, "you need to have an arrow to study a shield." I believe a Rubicon has been crossed by the world with this pandemic and framing the proper understanding of how we got here, and the proper response will be the critical next steps.

Evidence of adenovirus vaccine sequences in early patients would suggest both that SARS-CoV-2 was created in a laboratory and that there was sufficient priority set on this project to create a specific vaccine for the chimera coronavirus.

When Oppenheimer saw the application of Einstein's physics in the embodiment of the atomic bomb, he is said to have quoted a line from the Hindu scripture, the Bhagavad Gita, which reads: 'Now I am become Death, the destroyer of worlds.' The contribution of physics' research to human killing would total less than 300,000 people in two ten-square mile zones in Japan, and the horrors of those events led the world to regulate the raw materials of such bombs and to sanction sovereign nations who attempted to violate the rules.

This had followed the contribution of chemistry to human killing in the form of chemical warfare during World War I, in which 100,000 were killed, and led the nations of the world to an historic agreement to never use chemical warfare again. It is now only 'rogue' operators who violate the norms civilized nations have agreed to.

It seems to be biology's turn to show its dark arts. If it is generally understood that biology/biotechnology has been harnessed to create a pandemic that has killed more people than physics and chemistry research combined, and to be a weapon where no place on earth is safe from its effects (SARS-CoV-2 has been detected in the deepest Amazon jungles and at research stations in Antarctica), there needs to be developed a new set of regulations, rules, etc. to both honor the 1.8 million innocent people who died from COVID-19 and to protect the world so this never happens again. It is also urgent to gather further data to support or refute if this was a Chinese bioweapons program, as the consequences of that would be significant.

Pre-publication peer review. The manuscript was provided by email to the following medical and scientific peers to afford an opportunity to review, comment, and critique the manuscript before publication. Those highlighted in yellow are members of the WHO-convened Global Study of the Origins of SARS-CoV-2¹⁵, The Lancet COVID-19 Commission¹⁶, or both.

¹⁵ <https://www.who.int/health-topics/coronavirus/origins-of-the-virus>

¹⁶ <https://covid19commission.org/origins-of-the-pandemic>

First Name	Last Name
John	Amuasi
Kristian	Andersen
Danielle	Anderson
Ralph	Baric
Francis	Collins
Carlos	das Neves
Peter	Daszak
Vladimir	Dedkov
Dominic	Dwyer
Anthony	Fauci
Hume	Field
Tedros Adhanom	Ghebreyesus
Eddie	Holmes
Gerald	Keusch
Marion	Koopmans
Dato' Sai Kit (Ken)	Lam
Fabian	Lendertz
W. Ian	Lipkin
Ken	Maeda
Hung	Nguyen
Stanley	Perlman
David	Quammen
Andrew	Rambaut
Angelie	Rasmussen
Linda	Saif
Zhengli	Shi
Supaporn	Wacharapluesadde


A Bayesian analysis concludes beyond a reasonable doubt that SARS-CoV-2 is not a natural zoonosis but instead is laboratory derived

Introduction. A two-hypothesis, Bayesian analysis was conducted to determine the origin of the SARS-CoV-2 pandemic. The conclusion was that it was created in a laboratory with synthetic biology tools from a bat beta coronavirus, subgenera sarbecovirus backbone (98.9% probability) and not from a natural, zoonotic transmission (1.1%).

There is no direct evidence of whether the release was accidental, or deliberate but circumstantial evidence makes it is highly likely it was accidental.

At the one-year anniversary of the first cases of COVID-19, the coronavirus pandemic caused by the SARS-CoV-2 virus, the origin of the virus remains unknown. While leading institutions and experts have been consistently adamant that it is a zoonotic disease which jumped from a bat reservoir host to humans directly or through an intermediate host the alternative possibility that it escaped from a laboratory conducting research remains a viable option.

In fact, in 2015 Peter Daszak, a leading zoonotic proponent of CoV-2 origin, wrote in, “Spillover and pandemic properties of zoonotic viruses with high host plasticity,”¹⁷ that transmission from laboratories was a major source of zoonotic disease. The Figure below from the Daszak paper shows this important relationship (green arrow):




Daszak et al. also writes: “**Zoonotic virus spillover** from wildlife was most frequent in and around human dwellings and in agricultural fields, as well as **at interfaces with occupational exposure to animals** (hunters, **laboratory workers**, veterinarians, **researchers**, wildlife management, zoo and sanctuary staff). **Primate hosts were most frequently cited as the source of viruses transmitted by direct contact** during hunting (exact $P = 0.051$) and **in laboratories**

¹⁷ <https://www.nature.com/articles/srep14830>


(exact $P = 0.009$).” [Emphasis added]. Primate “hosts” can presumably include monkey cell culture, such as the ubiquitous VERO cell used in all virology laboratories, including the WIV.

In 2015 Dr. Daszak spoke of the spillover danger of certain types of laboratory research:



Assessing Coronavirus threats

Peter Daszak
EcoHealth Alliance, New York, USA
www.ecohealthalliance.org



Local conservation.
Global health.

Follow up Genetic and Experimental studies (post-PREDICT) to Further Assess Spillover Potential

With each step, increased risk possible

- Virus isolation
- Sequence whole genome
- With temporally sampled viruses, measure mutation rates and phylodynamics
- Sequence receptor binding domain, if known
- Structural comparison with human receptors (e.g. 3D models, *In silico*)
- Cell line infection experiments (*in vitro*)
- Humanized mice and other animal experiments

He writes: “with each step, increased risk possible” with “Humanized mice and other animal experiments” the highest risk work.

In a prescient Twitter post in November 2019, he highlights the work he is doing using recombinant viruses with humanized mice and making viruses that “**don’t respond to MABs, vaccines...**” in response to criticism his work is of limited value:



Clearly, before the beginning of the pandemic, Daszak, now a member of both the WHO and Lancet teams being sent to China to explore the origin of CoV-2, could entertain the eal possibility of a laboratory created virus escaping into the human population/community.

The purpose of this analysis is to use a Bayesian Inference Network approach to the collected circumstantial evidence that is available to provide likelihoods of the alternative hypotheses as to the origin of SARS-CoV-2. The analysis also will include certain prior probabilistic conclusions to help set the initial state before the proprietary evidence is used.

Origin hypotheses: Initial States to establish the posterior probabilities.

Two published Bayesian analyses and two independent studies of zoonotic spillover from nature and laboratory-acquired infections in Asia will be used to establish the posterior probabilities for this analysis.

Zoonotic spillover frequency versus laboratory acquired infection frequency based on two published papers, one by Daszak et al.

In 2015 Daszak et al. published a paper entitled, “Spillover and pandemic properties of zoonotic viruses with high host plasticity,”¹ in which they identified 162 zoonotic viruses with naturally occurring animal-to-human transmission from 1990-2010. This is a frequency of $162/20 = 8.1$ events per year.

They also note: “The majority (94%) of zoonotic viruses described to date ($n = 162$) are RNA viruses, which is 28 times higher (95% CI 13.9–62.5, exact $P < 0.001$) than the proportion of RNA viruses among all vertebrate viruses recognized, indicating that RNA viruses are far more likely to be zoonotic than DNA viruses.” CoV-2 is an RNA virus.

Finally, they note that: “In general, wild animals were suggested as the source of zoonotic transmission for 91% (86/95) of zoonotic viruses compared to 34% (32/95) of viruses transmitted from domestic animals and 25% (24/95) with transmission described from both wild and domestic animals.”

One of the caveats of the Daszak data is that it categorizes a laboratory-acquired infection (LAI) from an animal collected from the wild as a zoonotic spillover. There is no data in the paper to assess this issue and leaving it uncorrected is a conservative approach since it only inflates the natural zoonotic frequency.

In 2018 a paper by Siengsan-Lamont entitled, “A Review of Laboratory-Acquired Infections in the Asia-Pacific: Understanding Risk and the Need for Improved Biosafety for Veterinary and Zoonotic Diseases,” was published.¹⁸ They reported 27 LAIs between 1982 and 2016, a frequency of $27/(2016 - 1982) = 0.8$ events per year.

Using these historical frequencies of zoonotic spillover versus LAI to predict a future event can be calculated in the following manner:

Evidence	Zoonotic Origin	Laboratory Origin
Frequency per year from Daszak paper	8.1	NA
Frequency per year from Siengsan-Lamont paper	NA	0.8
Total events per year	$8.1 + 0.8 = 8.9$	$8.1 + 0.8 = 8.9$
Likelihood of future event based on historical frequency	$8.1/8.9 \times 100 = 0.91$	$0.8/8.9 \times 100 = 0.9$

Daszak’s initial state analysis. This evidence sets the likelihood that CoV-2 was a zoonotic origin event at 91% and a laboratory origin event at 9%.

¹⁸ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6073996/>

Independent prior analyses: Rootclaim.

The next data that will be used is a recent analysis published on the Rootclaim website.¹⁹ Three hypotheses below were analyzed through a series of evidence statements and the probabilities that each was the origin of SARS-CoV-2 determined:

Hypothesis	Calculated Probability
Lab escape: The virus was the subject of genetic research, including gain-of-function, and was released by accident	81%
Zoonotic: The virus evolved in nature and was transmitted to humans from a non-human vertebrate animal	16%
Bioweapon: The virus was genetically engineered as a bioweapon and was deliberately released	3%

As can be seen, the highest likelihood probability is an accidental lab escape, the lowest a bioweapon. The details of the evidence used to arrive at this conclusion is contained in Appendix 1. A summary of the changes in probability at each level of evidence analysis is shown in this table:

Evidence	Laboratory	Zoonosis	Bioweapon
Starting point	1.2%	82%	16%
Contagion and mortality	1.4%	97%	1.9%
Outbreak location: Wuhan	42%	56%	2.8%
Virus sources near Wuhan	16%	83%	1.0%
Chimera	37%	60%	2.5%
Furin cleavage	72%	23%	4.8%
WIV lab procedures	80%	17%	3.5%
WIV disassociation	89%	9%	2.0%
Chinese response	90%	8%	1.7%
No reported infections at WIV	86%	11%	2.4%
No whistleblowers	81%	16%	2.8%

As can be seen, the starting point assumed an 82% probability of a zoonotic origin. This starting point is a reasonable value and will be used here. Since some of the evidence in the above analysis will be used here, only the starting point will be used and not the probability changes from there.

For purposes of this analysis only the Rootclaim initial state will be used since much of their evidence is also covered in the analysis here.

¹⁹ <https://www.rootclaim.com/analysis/what-is-the-source-of-covid-19-sars-cov-2>

In a paper by Daszak and colleagues it states: “In general, wild animals were suggested as the source of zoonotic transmission for 91% (86/95) of zoonotic viruses compared to 34% (32/95) of viruses transmitted from domestic animals and 25% (24/95) with transmission described from both wild and domestic animals.”¹

On the other hand, domestic animals seem to have been ruled out for SARS-CoV-2. In an interview for *Science* in July 2020, Dr. Zhengli Shi, head of coronavirus research at the Wuhan Institute of Virology, stated: “Under the deployment of the Hubei Provincial Government, our team and researchers from Huazhong Agricultural University collected samples of farmed animals and livestock from farms around Wuhan and in other places in Hubei Province. We did not detect any SARS-CoV-2 nucleic acids in these samples.”²⁰

Reanalysis of Rootclaim initial state to remove Bioweapons option.

The US government uses the following definitions:

“Gain-of-function (GOF) studies, or research that improves the ability of a pathogen to cause disease, help define the fundamental nature of human-pathogen interactions, thereby enabling assessment of the pandemic potential of emerging infectious agents, informing public health and preparedness efforts, and furthering medical countermeasure development.

Gain-of-function studies may entail biosafety and biosecurity risks; therefore, the risks and benefits of gain-of function research must be evaluated, both in the context of recent U.S. biosafety incidents and to keep pace with new technological developments, in order to determine which types of studies should go forward and under what conditions.”²¹

“Dual use research of concern (DURC) is life sciences research that, based on current understanding, can be reasonably anticipated to provide knowledge, information, products, or technologies that could be directly misapplied to pose a significant threat with broad potential consequences to public health and safety, agricultural crops, and other plants, animals, the environment, materiel, or national security.”²²

For this analysis, the assumption is made that GOF and DURC are largely the same processes and techniques in the laboratory and thus can only be distinguished by direct, documentary evidence of the intent of the research from administrators in the facilities conducting the work.

In the absence of any such documentary evidence that bioweapon research was being conducted or that SARS-CoV-2 is a bioweapon and to take the least inflammatory posture, the initial state for the above prior analysis will be recalculated by eliminating the hypothesis, and its accompanying probability, that SARS-CoV-2 was created as a bioweapon. The revised initial state calculation is shown in this table:²³

²⁰ <https://www.sciencemag.org/sites/default/files/Shi%20Zhengli%20Q%26A.pdf>

²¹ <https://www.phe.gov/s3/dualuse/Pages/GainOfFunction.aspx>

²² <https://www.phe.gov/s3/dualuse/Pages/default.aspx>

²³ For clarity, the 3% bioweapon probability was simply dropped and the remaining likelihoods, 81% and 16%, were normalized.

Evidence	Zoonotic Origin	Laboratory Origin	Bioweapons Origin
Rootclaim initial state	0.86	0.012	0.16
Remove bioweapons	NA	NA	0
Normalize remaining hypotheses	$0.86/(0.86 + 0.012) = 0.986$	$0.012/(0.86 + 0.012) = 0.014$	NA

Rootclaim Initial state analysis, adjusted. This evidence sets the likelihood that CoV-2 was a zoonotic origin event at 98.6% and a laboratory origin event at 1.4%.

Additional Prior Evidence by Demaneuf and De Maistre. A second prior Bayesian analysis was performed by professionally educated risk assessment personnel and Chinese-language speaking professionals²⁴ and is included herein in its entirety. For the sake of brevity, the zoonotic origin evidence was based primarily on population size, distribution, and geographic distribution of bat populations relative to Wuhan. With respect to a lab accident, they separately analyze probabilities of a virus escape during collection, transport, and direct lab accidents and then separately the probability of a community outbreak following a lab escape. They also use primary Mandarin-language sources for Chinese estimates of the same events, showing corroboration of the probabilities. Their conclusion is that the probability of a lab escape ranges from 6% to 55% with a zoonotic origin a zoonotic origin probability being 45% to 94%.

Second Bayesian analysis. Using the most conservative probabilities, this evidence sets the likelihood that CoV-2 was a zoonotic origin event at 94% and a laboratory origin event at 6%.

Selection of initial state for Bayesian analysis.

The Text-Table below summarizes the three approaches to an initial state as to the origin of CoV-2. While the Demaneuf and De Maistre analyses set a range for the zoonotic origin of 45% to 94%, I have used the top of the range of their probability of a zoonotic origin to be conservative.

Prior Analysis	Zoonotic Origin	Laboratory Origin
Daszak et al. paper	91%	9%
Rootclaim Bayesian analysis	98.6%	1.4%
Demaneuf and De Maistre Bayesian analysis	94%	6%

Using a simple online calculator²⁵ the mean of these three value sets is 94.5%, the standard deviation is $\pm 3.8\%$, and the 95% confidence interval is $\pm 4.3\%$. Using these data, the upper bound of the 95% confidence interval is 98.8% and, to be most conservative, this will be used as the starting probability of a zoonotic origin.

Initial state for this analysis. The likelihood that SARS-CoV-2 began as a zoonotic event is 98.8% and the likelihood it began as a laboratory event is 1.2%.

²⁴ <https://zenodo.org/record/4067919#.X-qlm9gzbOj> . For reference purposes, this paper comes with a spreadsheet listing 112 individual BSL-3 labs in China across 62 lab-complexes.

²⁵ <https://www.calculator.net/standard-deviation-calculator.html?numberinputs=91%2C+94%2C+98.6&ctype=s&x=48&y=19>

1. **General approach of this analysis**²⁶

This analysis is intended to examine two competing and mutually exclusive theories of the origin of the coronavirus, SARS-CoV-2 (CoV-2), and the pandemic it has caused, COVID-19.

At the time of this writing there have been 83 million confirmed cases and 1.8 million deaths.²⁷ Some sources place the economic damage at \$21 trillion USD.

Bayes Theorem

This brief description of the Bayes Theorem was taken from the work of Jon Seymour:²⁸

“The eponymously named [Bayes Theorem](#) was discovered by the Reverend Thomas Bayes in the 1700’s and saved for posterity by an archivist of his papers who discovered the work posthumously. In common language, it provides a rational technique for revising a prior belief in light of new evidence. The equation for Bayes Theorem is given below:

$$P(H|E) = \frac{P(E|H).P(H)}{P(E)}$$

where:

- H is the statement of the hypothesis of interest
- P(H) is the prior probability that the hypothesis is true, independent of the evidence.
- E is the evidence being used to revise the belief in hypothesis
- P(E) is the marginal likelihood of the evidence, independent of the hypothesis
- P(E|H) is the likelihood the evidence, given that the hypothesis is true
- P(H|E) is the posterior probability of the hypothesis, given the evidence.

P(E) is sometimes difficult to estimate, but the following identity must hold:

$$P(E) = P(E|H).P(H) + P(E|\hat{H}).P(\hat{H})$$

Here P(E| \hat{H}) is the probability of the evidence, assuming the hypothesis is false and P(\hat{H}) is the probability the hypothesis is false which is the same as 1-P(H). Estimating the two conditional probabilities P(E|H) and P(E| \hat{H}) is generally easier than estimating the unconditional probability, P(E).”

²⁶ The statistical approach and many of the individual statistical analyses were performed by Dr. Martin Lee, PhD, Adjunct Professor of Biostatistics, UCLA. <https://ph.ucla.edu/faculty/lee> The likelihood adjustments to the Bayesian analysis, which you can see are routine math, were conducted by the author.

²⁷ <https://www.worldometers.info/coronavirus/coronavirus-cases/>

²⁸ <https://jonseymour.medium.com/a-bayesian-analysis-of-one-aspect-of-the-sars-cov-2-origin-story-where-the-first-recorded-1fbdcbea0a2b>

Theory One. The zoonotic theory is that a vertebrate animal was infected with CoV-2 or an ancestor (Index Host) and that a human was infected with contact to that Index Host in some manner. Human-to-human spread then followed.

Theory Two. The laboratory origin theory is that CoV-2 or an ancestor was being used in laboratory experiments and that it ‘escaped’ from the lab via an infected person, lab animal, experimental waste, etc.

I have found no evidence of a deliberate release and early firsthand accounts of local officials and scientists suggest surprise and consternation. If this was a deliberate release, such evidence would be extremely local, limited in distribution, and highly compartmentalized. It is beyond the scope of this analysis.

Weight of the evidence. For purposes of the calculation of posterior probabilities in the Bayesian analysis, evidence which has a statistical basis will be used directly to adjust the probabilities.

Statistically significant evidence. Since some of the probability calculations have astronomical values which would make a single such evidence statement, if inputted directly, swamp any further calculation and make their later contribution mute, a decision was made to simply treat quantitative probabilities as significant at the $p = 0.05$ level, no matter how much ‘more significant’ the calculation suggested.

So, for example, a probability of certain codon usage coming from nature may be one in 440 or $p = 0.002$, the contribution of this evidence to the input to the posterior probability adjustment would be set at a p -value of 0.05. In such cases the adjustment would be to change the ‘winning’ hypothesis by multiplying by 19, since a $p = 0.05$ is the same as a 19 out of 20 likelihood event. This is a conservative treatment of what would be highly significant data.

Other quantitative evidence. If a piece of evidence can be quantified but it does not reach a significance of $p = 0.05$ it will be used directly in the likelihood adjustment.

Non-quantitative evidence. For evidence that cannot be quantified, the decision was made to treat these as quantitative outcomes with a 51% to 49% likelihood value with respect to the ‘winning’ hypothesis. This has the effect of increasing the probability of that hypothesis for that step in the Bayesian analysis by 1.04. This 51%/49% concept is related to the legal standard of the ‘preponderance of the evidence’ used in civil litigation.

Independence. An important qualitative assessment that must be made is whether or not two pieces of evidence are independent of each other. If they are independent, they can each be used in determining a new likelihood calculation. If they are dependent on each other then they must be combined and only a single new likelihood analysis can be made. Where ever possible, evidence statements that could be considered as dependent are called out and this rule is followed on their contribution to the analysis.

Subjective Discount Factor. The impact of each piece of evidence was adjusted further by a subjective discount factor. This is a qualitative assessment of the overall veracity of a particular

piece of evidence when all factors, samples, methods, data sources, etc. are taken into context. It varies from 60% to 100% and is used as a fraction to reduce the impact of a single piece of evidence even further.

Hearsay. Just as in a court of law, evidence, usually attributed to a given person or persons, that is not directly available but instead relies on statements of others is usually not allowed in a court trial and will accordingly not be used here to adjust the Bayesian analysis. It may be recorded and preserved as a placeholder and reminder for further research. If new, direct evidence can be found then the bar of using it is lifted and it can be used for adjustment.

Significant figures. Because of the overall nature of the analyses here, all math calculations related to likelihoods are performed and carried forward at the ‘one significant figure’ level, with standard rounding rules applied. This has the effect, near the end of the cumulative evidence, of failing to change the relative probabilities as the small adjustments are reversed in the rounding process.

Evidence. International committees to investigate the origin of SARS-CoV-2 may not be impartial.

At the time of the writing of this manuscript there are two committees charged with examining the evidence and determining the origin of the SARS-CoV-2 virus. One committee is commissioned by the World Health Organization (WHO) and the other is an ad hoc committee established by the British medical journal, *The Lancet*.

The composition of the two committees is shown in the Text-Table below:

	Lancet Commission of CoV-2	WHO Commission on CoV-2 origin
	Dr. Peter Daszak, Chair	Dr. Peter Daszak, Ph.D (EcoHealth Alliance, USA)
	Dr. John Amuasi	Prof. John Watson (Public Health England, United Kingdom)
	Dr. Danielle Anderson	Prof. Dr. Marion Koopmans, DVM PhD (Erasmus MC, Netherlands)
	Dr. Isabella Eckerle	Prof. Dr. Dominic Dwyer, MD (Westmead Hospital, Australia)
Also co-author	Dr. Hume Field	Vladimir Dedkov, Ph.D (Institute Pasteur, Russia)
	Dr. Gerald Keusch	Dr. Hung Nguyen, PhD (International Livestock Research Institute (ILRI), Vietnam)
	Dr. Dato' Sai Kit (Ken) Lam	PD. Dr. med vet. Fabian Lendertz (Robert Koch-Institute, Germany)
	Dr. Carlos das Neves	Prof. Dr. Thea Fisher, MD, DMSc(PhD) (Nordsjællands Hospital, Denmark)
	Dr. Malik Peiris	Dr. Farag El Moubasher, Ph.D (Ministry of Public Health, Qatar)
	Dr. Stanley Perlman	Prof. Dr. Ken Maeda, PhD, DVM (National Institute of Infectious Diseases, Japan)
	Dr. Linda J. Saif	WHO Commission of CoV-2 origin
	Dr. Supaporn Wacharapluesadee	
	Lancet Commission on CoV-2	
	Signed Lancet letter	
	Co-author with Daszak	

There are a number of potential conflicts of interest:

Fully half of The Lancet's team had already suggested that any lab-leak hypothesis was a “conspiracy theory” in a January 2020 paper that has been shown elsewhere within to have been orchestrated behind the scenes to appear spontaneous.



The above paper published in August 2020 has as co-authors Drs. Hume, Daszak, and Shi. Having two of these scientists be asked to investigate a third co-author is a clear conflict of interest.

A newspaper piece about Peter Daszak entitled, “The doctor who denied COVID-19 was leaked from a lab had this major bias,”²⁹ questions his ability to be unbiased due to a deep, long history of work with Dr. Zhengli Shi of the WIV.

A lengthy piece in Wired was subtitled, “The two major investigations into the origins of the pandemic are compromised by potential conflicts of interest.”³⁰

Since the purpose of this manuscript is to evaluate the scientific evidence concerning the origin of SARS-CoV-2 no further effort will be put into these matters. If and when a report is prepared from either committee there will be time to analysis the work in the reports and compare it to prior publications and statements from the committee members to look for bias.

Likelihood from initial state is unchanged following this evidence analysis:

Zoonotic origin (98.8%) and laboratory origin (1.2%)

²⁹ <https://nypost.com/2021/01/16/doctor-who-denied-covid-was-leaked-from-a-lab-had-this-major-bias/>

³⁰ https://www.wired.com/story/if-covid-19-did-start-with-a-lab-leak-would-we-ever-know/?utm_source=twitter&utm_medium=social&utm_campaign=onsite-share&utm_brand=wired&utm_social-type=earned

Evidence. Three high visibility papers grounded the zoonotic origin hypothesis in the public conversation from February to May 2020: a pros and cons analysis.

Introduction. The two key data points from December 2019 concerning the origin of the SARS-CoV-2 coronavirus infection, the cause of COVID-19, are the observation that a large number of the earliest patients worked or had visited the Hunan Seafood Market in Wuhan, China and that the hospitals where the first patients were admitted were a short distance from the Wuhan Institute of Virology (WIV), the only high security, BSL-4 laboratory in all of China, and arguably the leading research institute in the world studying coronaviruses of the type causing COVID-19.

The first data point is reminiscent of the origin of SARS-CoV-1, a zoonosis with interspecies transmission from bats to civet cats and then to humans, identified in wet markets in southern China. The second data point is reminiscent of the four SARS-CoV-1 human spillovers that occurred after the 2003 epidemic ended and were each a laboratory-acquired infection (LAI) by a scientist working in a government research laboratory, much like the WIV, and then local human-to-human spread and nearby hospital admission.

To be clear in this paper, the term zoonosis will only be used to describe a interspecies transmission outside of a laboratory. This point seems important to clarify since Dr. Zhengli Shi, head of coronavirus research at the WIV, has previously reported: “An outbreak of hemorrhagic fever with renal syndrome occurred among students in a college (College A) in Kunming, Yunnan province, China in 2003. Subsequent investigations revealed the presence of hantavirus antibodies and antigens in laboratory rats at College A and two other institutions. Hantavirus antibodies were detected in 15 additional individuals other than the index case in these three locations. Epidemiologic data indicated that the human infections were a result of **zoonotic transmission** of the virus from laboratory rats.”³¹ [emphasis added.] The author has found no other support for the use of the term zoonotic transmission with respect to an LAI and its dual use could be confusing, and so will be avoided.

While the two initial data points would suggest that a balanced approach should be taken with respect to investigations of the origin of SARS-CoV-2, three high visibility publications that argued the laboratory origin idea was a “conspiracy theory” and strongly argued that it was of zoonotic origin foreclosed legitimate debate for much of 2019. The purpose of this evidence analysis is to examine these papers and weigh the strength of the evidence.

Paper 1: The February 3, 2020 paper by WIV scientist Dr. Shi et al. entitled: “A pneumonia outbreak associated with a new coronavirus of probable bat origin.”

This seminal paper set the stage for the zoonotic origin of SARS-CoV-2 and has been accessed over one million times. According to *Nature*, this article is in the 99th percentile (ranked 24th) of the 326,159 tracked articles of a similar age in all journals and the 99th percentile (ranked 2nd) of the 783 tracked articles of a similar age in *Nature*.

³¹ <https://pubmed.ncbi.nlm.nih.gov/20380897/>

However, a careful analysis of it shows serious issues which suggest it is unreliable. The following analysis is in the form of an independent manuscript:

The seminal paper from the Wuhan Institute of Virology claiming SARS-CoV-2 probably originated in bats appears to contain a contrived specimen, an incomplete and inaccurate genomic assembly, and the signature of laboratory-derived synthetic biology

The coronavirus RaTG13 was purportedly identified in a bat “fecal” specimen that is probably not feces, has significant unresolved method-dependent genome sequence errors and an incomplete assembly with significant gaps, and has an anomalous base substitution pattern that has never been seen in nature but is routinely used in codon-optimized synthetic genome constructions performed in the laboratory

Abstract. The species of origin for the SARS-CoV-2 coronavirus that has caused the COVID-19 pandemic remains unknown after over six months of intense research by investigators around the world. The current consensus theory among the scientific community is that it originated in bats and transferred to humans either directly or through an intermediate species; no credible intermediate species exists at this time. The suggested origin early on from a Wuhan “wet market” has been determined to be a red herring and the pangolin is no longer considered a likely intermediate by the virology community.

The basis for the hypothesis that SARS-CoV-2 probably evolved from bats initially came from a February 2020 paper³² from Dr. Zheng-Li Shi’s laboratory at the Wuhan Institute of Virology (WIV). In that paper the Wuhan laboratory made two claims: 1), “a bat fecal sample collected from Tongguan town, Mojiang county in Yunnan province in 2013” contained a coronavirus, originally designated “Rhinolophus bat coronavirus BtCoV/4991³³” in 2016 but renamed in their paper, RaTG13; and 2), the genomes of RaTG13 and SARS-CoV-2 had an overall identity of 96.2%, making it the closest match to SARS-CoV-2 of any coronavirus identified at that time. RaTG13 remains the closest match to SARS-CoV-2 at the current time.

In this paper I document that:

- 1) The RaTG13 specimen was not a bat fecal specimen, based on a comparison of the relative bacterial and eukaryotic genetic material in the purported fecal specimen to nine authentic bat fecal specimens collected in the same field visits as RaTG13 was collected by the Wuhan laboratory, run on the same Illumina instrument (id ST-J00123), and published in a second paper in February 2020.¹⁵ While the authentic bat fecal

³² Zhou, P., Yang, X., Wang, X. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270–273 (2020). <https://doi.org/10.1038/s41586-020-2012-7>.

³³ A [Coronavirus BtCoV/4991 Genbank entry](#) by Dr. Shi records: organism="Rhinolophus bat coronavirus BtCoV/4991." In July 2020 she wrote: “Ra4991 is the ID for a bat sample while RaTG13 is the ID for the coronavirus detected in the sample. We changed the name as we wanted it to reflect the time and location for the sample collection. 13 means it was collected in 2013, and TG is the abbreviation of Tongguan town, the location where the sample was collected.”

samples were, as expected, largely bacterial (specifically, 65% bacteria and 12% eukaryotic genetic sequences), the purported RaTG13 specimen had a reversed composition, with mostly eukaryotic genes and almost no bacterial genetic material (0.7% bacteria and 68% eukaryotic). The RaTG13 specimen was also only 0.01% virus genes compared to an average of 1.4% for authentic bat fecal specimens. A Krona analysis identified 3% primate sequences consistent with VERO cell contamination, the standard monkey cell culture used for coronavirus research, including at the Wuhan laboratory. Based on using the mean and standard deviation of the nine authentic bat fecal specimens from the Wuhan laboratory, the probability that RaTG13 came from a true fecal sample but had the composition reported by the Wuhan laboratory is one in thirteen million;

- 2) According to multiple references, RaTG13 was identified via Sanger dideoxy sequencing before 2016, partially sequenced by amplicon sequencing in 2017 and 2018, and then complete sequencing and assembly by RNA-Seq in 2020, although some reports from WIV suggest the timing of the RNA-Seq experiments may have been performed earlier than 2020. In any case, a Blast analysis of sequences from the amplicon and RNA-Seq experiments indicates an approximate 5% nucleotide difference, 50-fold higher than the technical error rate for RNA-Seq of about 0.1%. At least two gaps of over 60 base-pairs, with no coverage in the RNA-Seq data, were easily identified. The incomplete assembly and anomalous, method-dependent sequence divergence for RaTG13 is troublesome;
- 3) The pattern of synonymous to non-synonymous (S/NS) sequence differences between RaTG13 and SARS-CoV-2 in a 2201 nucleotide region flanking the S1/S2 junction of the Spike Protein records 112 synonymous mutation differences with only three non-synonymous changes. Based on the S/NS mutational frequencies elsewhere in these two genomes and generally in other coronaviruses the probability that this mutation pattern arose naturally is approximately one in ten million. A similar pattern of unnatural S/SN substitutions was seen in a 10,818 nt region of the pp1ab gene. This pp1ab gene pattern has a probability of occurring naturally of less than one in 100 billion. A total of four regions of the RaTG13 genome, coding for 7,938 nt and about one-quarter of the entire genome, contain over 200 synonymous mutations without a single non-synonymous mutation. This has a probability of one in 10^{-17} . A possible explanation, the absolute criticality of the specific amino acid sequence in the regions which might make a non-synonymous change non-infective, is ruled out by the rapid appearance of an abundance of non-synonymous mutations in these very regions when examining the over 80,000 human SARS-CoV-2 specimens sequenced to date. An alternative hypothesis, that this arose by codon substitution is examined. It is demonstrated, by example from a published codon-optimized SARS-Cov-2 Spike Protein experiment, that the anomalous S/SN pattern is precisely the pattern which is produced, by design, when synthetic biology is used and represents a signature of laboratory construction.

Based on the findings concerning the RaTG13 data, including anomalies and inconsistent statements about RaTG13, its origin, renaming, and sequencing timing; the finding that the specimen it is purported to have come from is not bat feces and has a signature of cell culture contamination; the unexplained method-dependent 5% sequence difference for RaTG13; and the S/SN mutation pattern reported, which to my knowledge has never been seen in nature, it can be concluded that RaTG13 is not a pristine biological entity but shows evidence of genetic manipulation in the laboratory.

Until a satisfactory explanation of the findings in this paper have been offered by the Wuhan laboratory, all hypotheses of the proximal origin of the entry of SARS-CoV-2 into the human population should now include the likelihood that the seminal paper contains contrived data. For example, the hypothesis that SARS-CoV-2 was the subject of laboratory research and at some point escaped the laboratory should be included in the narrative of the origin of SARS-CoV-2 research.

Introduction. Since the first reported patient on December 1, 2019 with a SARS-CoV-2 infection, the virus has caused a pandemic that has led to twenty-five million cases worldwide and over 840,000 deaths as of August 30, 2020. To make progress on treating this disease and preventing the next viral outbreak, knowing the origin of the virus and how it entered the human population is critical.

On February 3, 2020 a paper was published from the Wuhan Institute of Virology that identified a bat coronavirus, RaTG13, as having a 96.2% identity to SARS-CoV-2, quickly providing support for a zoonotic origin, either from bats directly or from bats to humans through an unknown intermediary species. If true, this would replicate the model of SARS-CoV 2003 in which the transmission was from bats to civets to humans and for MERS in which the transmission was from bats to camels to humans. At the time of this paper and through August 30, 2020, no other virus has been identified with a closer sequence homology to SARS-CoV-2 than RaTG13. The publication containing the RaTG13 sequence has been cited over 1600 times in the six months since publication. None of these studies contain research on the isolated virus itself since the virus has never been isolated or cultured. It was apparently found in only one sample from 2013 and that sample has been exhausted.³⁴

An examination of the raw data associated with RaTG13 immediately identified serious anomalies, bringing into question the existence of RaTG13 as a biological entity of completely nature origin.

³⁴ [Dr. Shi Science interview July 2020](#)

Materials and Methods.

GenBank accession URL table for sequences used in this paper.

The GenBank accession URLs for the specimens, raw reads, and sequences that are used in this paper are contained in the following Table, which can be used to reach the raw data.

Descriptor	URL Hyperlink
SARS-CoV-2 reference sequence in GenBank	SARS-CoV-2 complete genome
Bat coronavirus RaTG13, complete genome, Genbank	RaTG13 complete genome
RaTG13 purported bat fecal specimen	SRR11085797
Rhinolophus bat coronavirus BtCoV/4991 RNA-dependent RNA polymerase (RdRp) gene, partial cds	BtCoV/4991 RdRp gene
SRX8357956: amplicon _sequences of RaTG13	Specimen descriptor
RNA-Seq data for RaTG13	RNA-Seq data for RaTG13
Reference fecal bat specimens from WIV	SRR11085736
Reference fecal bat specimens from WIV	SRR11085734
Reference fecal bat specimens from WIV	SRR11085737
Reference fecal bat specimens from WIV	SRR11085733
Reference fecal bat specimens from WIV	SRR11085735
Reference fecal bat specimens from WIV	SRR11085738
Reference fecal bat specimens from WIV	SRR11085739
Reference fecal bat specimens from WIV	SRR11085740
Reference fecal bat specimens from WIV	SRR11085741

Below is a screen shot of the GenBank entry for the purported specimen from which RaTG13 was identified and upon which RNA-Seq was performed. While the title claims it is a “Rhinolophus affinis fecal swab” specimen it also records in the design of work entry that “(t)otal RNA was extracted from bronchoalveolar lavage fluid.” These descriptions are clearly inconsistent.

SRX7724752: RNA-Seq of Rhinolophus affinis:Fecal swab 1 ILLUMINA (Illumina HiSeq 3000) run: 11.6M spots, 3.3G bases, 1.7Gb downloads				
Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit following the manufacturers instructions. An RNA library was then constructed using the TruSeq Stranded mRNA Library Preparation Kit (Illumina, USA). Paired-end (150 bp) sequencing of the RNA library was performed on the HiSeq 3000 platform (Illumina).				
Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences				
Study: Bat coronavirus RaTG13 Genome sequencing PRJNA606165 • SRP249482 • All experiments • All runs show Abstract				
Sample: SAMN14082201 • SRS6146537 • All experiments • All runs Organism: unidentified coronavirus				
Library: Name: RaTG13 Instrument: Illumina HiSeq 3000 Strategy: RNA-Seq Source: METAGENOMIC Selection: RANDOM Layout: PAIRED				
Runs: 1 run, 11.6M spots, 3.3G bases, 1.7Gb				
Run	# of Spots	# of Bases	Size	Published
SRR11085797	11,604,666	3.3G	1.7Gb	2020-02-13

Apparent missing amplicon reads for RaTG13 in GenBank.

There are 33 amplicon reads in GenBank for RaTG13 from experiments recorded as having been performed in 2017 and 2018. A file naming pattern was noticed among the data sets which suggests there may be amplicon runs that were not deposited in GenBank. These files, if related to RaTG13, may contain useful sequence data and an effort should be made to retrieve them and, if appropriate, upload them to GenBank. A Table with the apparently missing data (yellow) is shown here.

Date	Amplicon file name endings						
3-Jun-17	A07	A08					
17-Jun-17	A05	A06					
20-Jun-17					F03	G03	H03
27-Sep-18	A06	B06	C06		E05	F05	G05/G06
29-Sep-18				D05	E05		G04
30-Sep-18	A02	B11					
8-Oct-18			C11				G10
11-Oct-18	A12	B12					
14-Oct-18	A02	B02	C02	D02			

Relationship of *Rhinolophus* bat coronavirus BtCoV/4991 and Bat coronavirus RaTG13.

The Wuhan laboratory has reported on the bat coronaviruses, BtCoV/4991 and RaTG13, in two peer-reviewed publications, one in 2016 and one in February 2020.³⁵ They have submitted three entries to GenBank for these two viruses, in 2016, February 2020, and May 2020.³⁶ The GenBank entries confirm sequencing experiments using Sanger dideoxy sequencing in 2016, PCR-generated amplicon sequencing performed on an AB 310 Genetic Analyzer in 2017 and 2018, and RNA-seq performed on an Illumina HiSeq 3000 (instrument id ST-J00123) in 2020. A single GISAID entry records that the RNA-seq data was obtained from an original specimen without passage.³⁷ This is an important detail since evidence of primate sequences, consistent with VERO cell contamination, is found in this specimen, as reported below, which would suggest laboratory passage.

None of these disclosures report that BtCoV/4991 and RaTG13 are the same coronavirus, simply renamed. This information was only disclosed in a written Question and Answer publication from *Science* magazine by Dr. Shi on July 31, 2020.^{4, 38} Given this disclosure months after the original publication concerning RaTG13 in *Nature* it is possible that the omission of the original publication and sequence data concerning BtCoV/4991 violated the “Reporting

³⁵ [2016 Virologica Sinica paper](#) and [February 2020 Nature paper](#)

³⁶ [RaTG13 complete genome Feb 2020](#), [Raw sequence reads for RaTG13 published Feb 2020](#), [Amplicon reads for RaTG13 from 2017 and 2018 published in May 2020](#).

³⁷ The GISAID entry is EPI_ISL_402131.

³⁸ Dr. Shi wrote: “Ra4991 is the ID for a bat sample while RaTG13 is the ID for the coronavirus detected in the sample. We changed the name as we wanted it to reflect the time and location for the sample collection. 13 means it was collected in 2013, and TG is the abbreviation of Tongguan town, the location where the sample was collected.”

standards and availability of data, materials, code and protocols” required for *Nature* publications.³⁹

The February 2020 papers uses the RNA-Seq data for RaTG13 genome determination but fails to disclose the previous data obtained by Sanger dideoxy sequencing in 2016 and by amplicon sequencing in 2017 and 2018. Since these unrecorded data establish method-dependent sequencing differences of up to 4% the failure to disclose this data or to reconcile these differences is troubling.

In addition, the raw assembly accession data for RaTG13 are not described or linked to the Genbank entry, MN669532, and also no assembly method is specified in the raw data SRX7724752.12 and the Illumina run. And the amplicon sequencing data has sequence gaps of approximately 20% of the genome. Therefore, no primary assembly data has been made available by the WIV for the RaTG13 genome. This is contrary to the *Nature* Reporting Standards⁹ as they state: “When publishing reference genomes, the assembly must be made available in addition to the sequence reads.”

Relationship of RaTG13 and SARS-CoV-2.

There have been two descriptions of the process by which the RaTG13 genome was identified as closely homologous to SARS-CoV-2. These seem to be inconsistent with each other.

In the February 2020 *Nature* paper⁵ it states:

“We then found that a short region of RNA-dependent RNA polymerase (RdRp) from a bat coronavirus (BatCoV RaTG13)—which was previously detected in *Rhinolophus affinis* from Yunnan province—showed high sequence identity to 2019-nCoV. We carried out full-length sequencing on this RNA sample (GISAID accession number EPI_ISL_402131). Simplot analysis showed that 2019-nCoV was highly similar throughout the genome to RaTG13, with an overall genome sequence identity of 96.2%.”

In a July 2020 interview the process was described:

“We detected the virus by pan-coronavirus RT-PCR in a bat fecal sample collected from Tongguan town, Mojiang county in Yunnan province in 2013, and obtained its partial RdRp sequence. Because the low similarity of this virus to SARS-CoV, we did not pay special attention to this sequence. In 2018, as the NGS sequencing technology and capability in our lab was improved, we did further sequencing of the virus using our remaining samples, and obtained the full-length genome sequence of RaTG13 except the 15 nucleotides at the 5’ end. As the sample was used many times for the purpose of viral nucleic acid extraction, there was no more sample after we finished genome sequencing, and we did not do virus isolation and other studies on it. Among all the bat samples we collected, the RaTG13 virus was detected in only one single sample. In 2020, we compared the sequence of SARS-CoV-2 and our unpublished bat

³⁹ [Nature research reporting standards for availability of data](#)

coronavirus sequences and found it shared a 96.2% identity with RaTG13. RaTG13 has never been isolated or cultured.”

If the full-length genome of RaTG13 was available by 2018 it is unclear why a database search within the WIV for coronaviruses that resembled SARS-CoV-2 would lead to identifying the 370-nt segment representing the RdRp gene (as stated in the February paper) but not the full length RaTG13 genome (which was stated to have been sequenced by 2018). In addition, an assembly of all available amplicon data for RaTG13 from 2017 and 2018 contains gaps of approximately 20% of the genome. If the sample was completely consumed during the 2017-8 sequencing it is unclear how RNA-Seq was conducted in 2020 to permit the full-length genome to be determined.

Analytical methods. Taxonomy of specimens was determined in the NCBI Sequence Read Archive and KRONA.⁴⁰ Blast was used for sequence alignment and comparisons.⁴¹

To evaluate the data from the bat species relative to the RaTG13 fecal sample analysis, the latter was treated as a fixed result with the comparison to the taxonomy results of the nine bat feces specimens. It also was noted that the data were clearly right skewed (and descriptively both mean/median and standard deviation/interquartile range were used). Therefore, a non-parametric procedure, the Wilcoxon signed-rank test was used with the p-value calculated by an exact procedure because of the small sample size. Considering the synonymous to non-synonymous mutation frequency and how to evaluate that for the various protein coding regions of the virus, it was noted that for all of the genes pooled, the ratio of the synonymous to non-synonymous regions was approximately 0.83. To analyze the corresponding distribution for each gene, we assumed that each mutation was an independent observation from a Bernoulli random variable and, therefore the number of synonymous mutations in the gene would have a binomial distribution (with probability 0.83). A probability was then computed for the actual number of synonymous mutations on this basis (the probability was determined on a one-sided basis, i.e. excess mutations, and was calculated as a strict inequality).

Results.

Original characterization of RaBtCoV/4991 (RaTG13) and related bat fecal specimen.

In 2016 Dr. Shi and colleagues published a paper entitled, “Coexistence of multiple coronaviruses in several bat colonies in an abandoned mineshaft⁴²” in which a number of novel bat coronaviruses were isolated from bat fecal specimens collected during 2012 and 2013. The viruses were named, according to the paper, in the following fashion:

⁴⁰ [NCBI Sequence Archive](#)

⁴¹ [Blast alignment](#)

⁴² Xing-Yi Ge, et. al., Coexistence of multiple coronaviruses in several bat colonies in an abandoned mineshaft, *Virologica Sinica*, 2016, 31 (1): 31–40. DOI: 10.1007/s12250-016-3713-9

“The positive samples detected in this study were named using the abbreviated bat species name plus the bat sample number abbreviation. For example, a virus detected from *Rhinolophus sinicus* in sample number 4017 was named RsBtCoV/4017. If the bat was co-infected by two different coronaviruses, numbers were appended to the sample names, such as RsBtCoV/4017-1 and RsBtCoV/4017-2.”

In the July 2020 interview Dr. Shi wrote:

“Ra4991 is the ID for a bat sample while RaTG13 is the ID for the coronavirus detected in the sample. We changed the name as we wanted it to reflect the time and location for the sample collection. 13 means it was collected in 2013, and TG is the abbreviation of Tongguan town, the location where the sample was collected.”

The 2016 and 2020 statements about the naming of virus RsBtCoV/4991 appear inconsistent with each other.

Of the 152 coronaviruses identified, 150 were classified as alphacoronaviruses while only two were classified as betacoronaviruses, HiBtCoV/3740-2 and RaBtCoV/4991. The naming convention from the paper means this latter coronavirus was identified in a fecal specimen from a *Rhinolophus affinis* bat and was sample number 4991.

The latter virus was described in the paper as follows:

“Virus RaBtCoV/4991 was detected in a *R. affinis* sample and was related to SL-CoV. The conserved 440-bp RdRp fragment of RaBtCoV/4991 had 89% nt identity and 95% aa identity with SL-CoV Rs672. In the phylogenetic tree, RaBtCoV/4991 showed more divergence from human SARS-CoV than other bat SL-CoVs and could be considered as a new strain of this virus lineage.”

The Genbank accession number for RaBtCoV/4991 is [MN KP876546.1](#) and in Genbank it is identified as having been collected in July 2013 as a “feces/swabs” specimen.

The RATG13 genome sequence was assembled from low coverage RNA-Seq data.


A Blast analysis of the RaTG13 genome against [SRR11085797](#) retrieved about 1700 reads which covers only about 252,000 nt of the total reads of 3.3 Gb. Since the genome size of RaTG13 is known to be about 30,000 nt this represents an 8-fold coverage, typically insufficient for a definitive assembly. For example, some have suggested a 30-fold coverage is necessary to create high quality assemblies.⁴³

⁴³ Sims, D. *et al.* Sequencing depth and coverage: key considerations in genomic analyses. *Nature Reviews – Genetics*. (2014) 15: 121-132. doi:10.1038/nrg3642.

At an eight-fold coverage and based on the typical practice of having four or more reads to call a SNP,⁴⁴ the 8-fold coverage of RaTG13 would have 4.2% bases or about 1260 calls of less than 4 reads and about 10 bases would be missed completely, with no calls at all.

A Blast of the RaTG13 published genome onto the RNA-Seq data documents at least two 60 base-pair gaps with no coverage, precluding a complete assembly.

Given the low coverage in the RNA-Seq data, an exploratory, non-exhaustive Blast search was conducted against the published RaTG13 sequence. Two gaps of over 60 nt, shown below, were easily found:




It is conceivable there are additional gaps but the above two are sufficient to document that the complete RaTG13 genome sequence could not have been assembled solely from the RNA-Seq data, as stated.²

Taxonomy analysis of the RaTG13 specimen is inconsistent with being from bat feces and shows evidence of laboratory cell culture contamination.

According to the Wuhan laboratory, the RaTG13 coronavirus was a fecal swab specimen collected from a *Rhinolophus affinis* bat in 2013. Unexpectedly, (Text-Figure below) the taxonomy analysis is primarily eukaryotic (green arrow; 67.91%) with only traces of bacteria (blue arrow; 0.65%). The viral genomes also make only a trace contribution (red arrow; 0.01%):

⁴⁴[Illumina Technical Bulletin Call Coverage](#)



[Taxonomy analysis for RaTG13 data SRR11085797](#)

To compare this specimen composition to bat fecal specimens collected by Dr. Shi and her WIV colleagues and analyzed in other studies, a paper from Dr. Shi’s laboratory, also published in February 2020, was identified. In this paper, entitled, “Discovery of Bat Coronaviruses through Surveillance and Probe Capture-Based Next-Generation Sequencing,”⁴⁵ a total of nine specimens “collected during previous bat CoV surveillance projects, (were) extracted from bat rectal swabs.” According to the Methods section in this paper, the “previous bat CoV surveillance projects” include the field work in 2013 when the RaTG13 was said to have been collected. The comparison below is thus the same specimens collected on the same field surveillance projects by the same investigators from the Wuhan laboratory and sequenced on the same Illumina instrument. These nine specimens will be referred to as “reference fecal specimens” henceforth.

The following Text-Table compares the taxonomical analysis of the RaTG13 and reference fecal specimens. The reference fecal specimens have an average eukaryotic genome content of about 12% while RaTG13’s eukaryotic content was 68%. On the other hand, the most abundant genes in the reference fecal specimens were bacterial, with an average of 65%; RaTG13 had less than 1% bacterial genes. And finally, the reference fecal specimens had 1.57% virus genes compared to the 0.01% virus genes of RaTG13.


⁴⁵ [Discovery of bat coronaviruses through surveillance and probe capture-based next-generation sequencing](#)

Specimen ID	Specimen Type	Unidentified Reads	Eukaryota	Bacteria	Viruses	Sum
SRR11085736	<i>Rhinolophus affinis</i>	0.86	4.36	91.07	0.03	96.32
SRR11085734	<i>Miniopterus schreibersii</i>	3.81	16.03	76.15	0.11	96.1
SRR11085737	<i>Scotophilus kuhlii</i>	17.98	8.59	67.81	2.19	96.6
SRR11085733	<i>Hipposideros larvatus</i>	13.27	27.99	42.96	4.1	88.32
SRR11085735	<i>Hipposideros pomona</i>	34.33	7.96	54.78	0.71	97.78
SRR11085738	<i>Pipistrellus abramus</i>	20.33	21.44	47.3	6.45	95.52
SRR11085739	<i>Tylonycteris pachypus</i>	61.75	14.34	20.06	0.06	96.21
SRR11085740	<i>Miniopterus pusillus</i>	0.78	1.46	99.22	0.05	101.51
SRR11085741	<i>Rousettus aegyptiacus</i>	6.44	2.59	88.36	0.45	97.84
Mean +/- SD	Nine bat feces specimens	17.73+/-19.79	11.64+/-9.02	65.30+/-26.10	1.57+/-2.28	96.24+/-3.45
Median +/- IQR	Nine bat feces specimens	13.27+/-24.995	8.59+/-15.26	67.81+/-41.58	0.45+/-3.09	96.32+/-2.00
SRR11085797	RaTG13 fecal specimen	29.38	67.91	0.65	0.01	97.95
	P-value (exact Wilcoxon signed-rank test)	0.16	0.0039	0.0048	0.0039	0.098

As shown in the Text-Table above the RaTG13 specimen is significantly different from the reference fecal specimens in composition. The probabilities for each category, eukaryote, bacteria, and virus, are individually highly statistically significant. They are also independent of each other and therefore the overall probability that RaTG13 has the composition of eukaryote, bacteria, and virus genes that was reported by the Wuhan laboratory but is actually from an authentic bat fecal specimen is less than one in 13 million.

The alternative conclusion is that this sample was not a fecal specimen but was contrived. The data cannot, however, distinguish between a non-fecal specimen that came from true field work on the one hand and a specimen created *de novo* in the laboratory on the other hand.

A graphical comparison of the above data is shown below and visually shows the significant differences between the WIV fecal specimens and the RaTG13 specimen, despite the claim they were collected in the same field surveillance trips:



Another comparison can be made between the reference fecal specimens and the RaTG13 specimen by looking at the taxonomy of the nine to twelve “strong signals” identified on the NCBI Sequence Read Archive. The following Text-Table is a summary of these findings.


Specimen	The identity of the Strong Signals in the Specimens		
	Bacteria	Eukaryotes	Viruses
Rhinolophus affinis anal swab (SRR11085736)	92%	One magnaorder of placental mammals, includes bat	None
Miniopterus schreibersii anal swab (SRR11085734)	88%	One bat , the host bat, Miniopterus sp.	None
Scotophilus kuhlii anal swab (SRR11085737)	56%	Two bats , mouse-eared and big brown bats.	Two viruses, kobuvirus (host includes bats) and a Scotophilus kuhlii coronavirus
Hipposideros larvatus anal swab (SRR11085733)	56%	One bat , the host bat, Hipposideros sp. and one rodent.	Hipposideros pomona bat coronavirus
Hipposideros pomona: Anal swab (SRR11085735)	78%	One bat , the host bat, Hipposideros sp.	None
Pipistrellus abramus: Anal swab (SRR11085738)	73%	Two bats , the big brown bat and the mouse-eared bat.	Pipistrellus abramus bat coronavirus
Tylonycteris pachypus: Anal swab (SRR11085739)	67%	Three bats , the microbat, the great roundleaf bat, and a superorder of mammals, which includes bats.	None
Miniopterus pusillus: Anal swab (SRR11085740)	89%	One bat , the Natal long-fingered bat.	None
Rousettus aegyptiacus: Anal swab (SRR11085741)	91%	One magnaorder of placental mammals, includes bats.	None
Average	77%		
RaTG13 Rhinolophus affinis:Fecal swab (SRR11085797)	None	All nine strong signals are eukaryotes. Five bats , the Great Roundleaf bat, resident of China, the Egyptian fruit bat, which is not found in China, a megabat, mouse-eared bat, and bent-winged bat. Two marmots, the Alpine marmot from Europe and the Yellow-bellied marmot of North America. The paraorder of whales. The red fox.	None

As can be seen, while the strong signals in the authentic specimens contain 56% to 92% (average 77%) bacterial signals, the RaTG13 specimen has no bacteria among the nine strong signals. Most specimens do not have virus strong signals but the three that do are host-related coronaviruses (four) or one host-related kobuvirus.

RaTG13 has no viral strong signals. Among the reference specimens with eukaryotic strong signals, they are either bat-related genes (eleven) or higher order taxonomy signals that include bats (three). There is one anomalous rodent-related signal among the reference specimens.

The RaTG13 specimen is again an outlier with all nine strong signals arising from eukaryotic genes. Five of the nine signals are bats, some resident to China and some with non-Chinese host ranges. Surprisingly, unlike three of the reference bat signals which are identified as host-related, the RaTG13 specimen did not contain *Rhinolophus* sp. host-related strong signals. The remaining four strong signals are marmot-related genes (two), whale-related gene (one), and red fox-related gene (one).

Finally, a Krona analysis (below) identifies 3% primate sequences (red arrow) in the RaTG13 sequence data. This is consistent with contamination by the standard laboratory coronavirus cell culture system, the VERO monkey kidney cell line.



Source: [Krona analysis of RaTG13 specimen](#)

It is unclear why these obviously anomalous findings were not detected during the peer-review process prior to publication of this important work. At this point, an explanation is needed from the WIV to refute the conclusion that the specimen identified as the source of RaTG13 is **not** a bat fecal/anal specimen and that the primate genetic material is consistent with a VERO cell contaminated specimen.

Method-related nt base substitutions in RaTG13.

The original Sanger dideoxy RdRp sequence reported in 2016 is homologous to RNA-seq data from 2020 but is non-homologous to amplicon sequencing data from 2017 and 2018.

As expected, a comparison of the 2016 RdRp GenBank sequence for BtCoV/4991 obtained by Sanger dideoxy sequencing with the RNA-seq sequencing of RaTG13 reported in *Nature* shows 100% identity over the 370 nt segment.

Sequence ID: Query_30201 Length: 370 Number of Matches: 1					
Range 1: 1 to 370 Graphics Next Match Previous Match					
Score	Expect	Identities	Gaps	Strand	
684 bits(370)	0.0	370/370(100%)	0/370(0%)	Plus/Plus	
Query 15322		GCCTCACTTGTCTTCTGCTCGCAACATACAACGTGCTGTAGCTTGTACACCGTTTCTAT			15381
Sbjct 1		GCCTCACTTGTCTTCTGCTCGCAACATACAACGTGCTGTAGCTTGTACACCGTTTCTAT			60
Query 15382		AGATTAGCTAATGAGTGTGCTCAAGTATTGAGTGAAATGGTCATGTGTGGCGGTTCACTA			15441
Sbjct 61		AGATTAGCTAATGAGTGTGCTCAAGTATTGAGTGAAATGGTCATGTGTGGCGGTTCACTA			120
Query 15442		TATGTTAAACAGGTGGAACCTCATCAGGAGATGCCACAACCTGCTTATGCTAATAGTGTC			15501
Sbjct 121		TATGTTAAACAGGTGGAACCTCATCAGGAGATGCCACAACCTGCTTATGCTAATAGTGTC			180
Query 15502		TTTAACATTGTCAAGCTGTTACGGCCAATGTTAATGCACCTTTATCTACTGATGGTAAC			15561
Sbjct 181		TTTAACATTGTCAAGCTGTTACGGCCAATGTTAATGCACCTTTATCTACTGATGGTAAC			240
Query 15562		AAAATTGCCGATAAGCACGTCCGCAATTTACAACACAGACTTTATGAGTGTCTCTATAGA			15621
Sbjct 241		AAAATTGCCGATAAGCACGTCCGCAATTTACAACACAGACTTTATGAGTGTCTCTATAGA			300
Query 15622		AATAGAGATGTTGACACAGACTTTGTGAATGAGTTTTACGCATATTTGCGTAAACATTTTC			15681
Sbjct 301		AATAGAGATGTTGACACAGACTTTGTGAATGAGTTTTACGCATATTTGCGTAAACATTTTC			360
Query 15682		TCAATGATGA			15691
Sbjct 361		TCAATGATGA			370


Surprisingly, the two amplicon sequences from 2017 that partially cover the 370 nt RdRp region have four base substitutions or gaps over a total segment of 219 nt (2% divergence).

Sequence ID: Query_64615 Length: 1100 Number of Matches: 1					
Range 1: 3 to 89 Graphics Next Match Previous Match					
Score	Expect	Identities	Gaps	Strand	
147 bits(79)	2e-39	87/90(97%)	3/90(3%)	Plus/Minus	
Query 15322		GCCTCACTTGTCTTCTGCTCGCAACATACAACGTGCTGTAGCTTGTACACCGTTTCTAT			15381
Sbjct 89		GCCTCACTTGTCTTCTGCTCGCAACATACAACGTGCTGTAGCTTGTACACCGTTTCTAT			30
Query 15382		AGATTAGCTAATGAGTGTGCTCAAGTATTG			15411
Sbjct 29		AGATTAGCTAATGAG-G-GCTCAAGT-TTG			3

Sequence ID: Query_31429 Length: 785 Number of Matches: 1					
Range 1: 655 to 783 Graphics Next Match Previous Match					
Score	Expect	Identities	Gaps	Strand	
233 bits(126)	1e-65	128/129(99%)	0/129(0%)	Plus/Minus	
Query 15563		AAATTGCCGATAAGCACGTCCGCAATTTACAACACAGACTTTATGAGTGTCTCTATAGAA			15622
Sbjct 783		AAATTGCTGATAAGCACGTCCGCAATTTACAACACAGACTTTATGAGTGTCTCTATAGAA			724
Query 15623		ATAGAGATGTTGACACAGACTTTGTGAATGAGTTTTACGCATATTTGCGTAAACATTTCT			15682
Sbjct 723		ATAGAGATGTTGACACAGACTTTGTGAATGAGTTTTACGCATATTTGCGTAAACATTTCT			664
Query 15683		CAATGATGA			15691
Sbjct 663		CAATGATGA			655

RaTG13 Spike Protein gene has 5% substitutions when comparing 2020 RNA-Seq and 2017 amplicon sequencing data.

The segment of RaTG13 which shows the greatest sequence divergence between the RNA-seq and amplicon sequencing methods spans from A8886 to A9987 and is shown here below. It contains 80 base substitutions/indels in a 1107 nt sequence (5% substitution and 2% gaps).

 [Download](#)

[Graphics](#)


[SRA](#)


SRX8357956

Sequence ID: [SRA:SRR11806578.14.1](#) Length: **1100** Number of Matches: **1**

Range 1: 14 to 1100

[Graphics](#)

 [Next Match](#)


 [Previous Match](#)

Score	Expect	Identities	Gaps	Strand
1716 bits(929)	0.0	1052/1107(95%)	25/1107(2%)	Strand Plus/Minus

No explanation has been offered in publications from the WIV for the method-dependent sequencing differences identified here, which are twenty- to 50-fold higher than the 0.1% technical error rate sometimes attributed to RNA-Seq data.

The Spike Protein gene sequence substitution divergence between RaTG13 and SARS-CoV-2 contains an improbable synonymous/non-synonymous pattern.

The functional structure of the SARS-CoV-2 Spike Protein is shown here:



The SARS-CoV-2 Spike protein (above) contains an S1 subunit and S2 subunit with the Polybasic Cleavage Site (PBCS) between R685 and S686. This cleavage is performed by a host cell surface protease, furin, and is an important attribute in explaining the virulence of SARS-CoV-2 compared to other human coronaviruses, which do not have a furin cleavage site. The PBCS also contains the unusual PRRA insertion that has not been previously seen in Clade B coronaviruses and for which no natural mechanism for its appearance has been offered.⁴⁶

The S1 subunit is located within the N-terminal 14–685 amino acids of S protein, containing N-terminal domain (NTD), receptor binding domain (RBD), and receptor binding motif (RBM). The S2 subunit contains a fusion peptide (FP), heptad repeat 1 (HR1), heptad repeat 2 (HR2), transmembrane domain (TM) and cytoplasmic domain (CP).

The base substitution pattern of synonymous and non-synonymous substitutions when comparing RaTG13 and the reference sequence of SARS-CoV-2 demonstrated an anomalous pattern for the coding region for aa 541 to 1273, a 733 aa protein segment representing over 60% of the SP gene.

As shown in the Text-Figure below, there are only three substitutions (red arrow) and the PBCS insertion (blue arrow) when comparing this segment of the RaTG13 and SARS-CoV-2 SP. Excluding the PBCS, the amino acid sequences are 99.6% identical.

⁴⁶ [The proximal origin of SARS-CoV-2.](#)

Score	Expect	Method	Identities	Positives	Gaps
1501 bits(3886)	0.0	Compositional matrix adjust.	726/733(99%)	728/733(99%)	4/733(0%)
Query 541	FNFNGLTGTGVLTESNKKFLPFQQFGRDIADTTDAVRDPQTLEILDITPCSFGGVSVITP				600
Sbjct 541	FNFNGLTGTGVLTESNKKFLPFQQFGRDIADTTDAVRDPQTLEILDITPCSFGGVSVITP				600
Query 601	GTNTSNQVAVLYQDVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIGAEHVNNNSY				660
Sbjct 601	GTN SNQVAVLYQDVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIGAEHVNNNSY				660
Query 661	ECDIPIGAGICASYQTQTN SPRARSVASQSIIAYTMSLGAENSVAYSNNNSIAIPTNFTI				720
Sbjct 661	ECDIPIGAGICASYQTQTN RSVASQSIIAYTMSLGAENSVAYSNNNSIAIPTNFTI				716
Query 721	SVTTEILPVSMTKTSVDCTMYICGDSSTECNLLQLQYGSFCTQLNRALTGIAVEQDKNTQE				780
Sbjct 717	SVTTEILPVSMTKTSVDCTMYICGDSSTECNLLQLQYGSFCTQLNRALTGIAVEQDKNTQE				776
Query 781	VFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPSKRSFIEDLLFNKVTADAGFIKQYGDC				840
Sbjct 777	VFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPSKRSFIEDLLFNKVTADAGFIKQYGDC				836
Query 841	LGDI AARDL ICAQKFNGLTVLPPLL TDEMIAQYTSALLAGTITSGWTFGAGAALQIPFAM				900
Sbjct 837	LGDI AARDL ICAQKFNGLTVLPPLL TDEMIAQYTSALLAGTITSGWTFGAGAALQIPFAM				896
Query 901	QMAYRFNGIGVTQNVLYENQKLIANQFN SAIGKIQDSLSTASALGKLQDVVNQNAQALN				960
Sbjct 897	QMAYRFNGIGVTQNVLYENQKLIANQFN SAIGKIQDSLSTASALGKLQDVVNQNAQALN				956
Query 961	TLVKQLSSNFGAISSVLNDILSRDKVEAEVQIDRLITGRLQSLQTYVTQQLIRAAEIRA				1020
Sbjct 957	TLVKQLSSNFGAISSVLNDILSRDKVEAEVQIDRLITGRLQSLQTYVTQQLIRAAEIRA				1016
Query 1021	SANLAATKMSECVLGQSKRVDFCGKGYHLSFPPQSAPHGVVFLHVTYVPAQEKNF TTAPA				1080
Sbjct 1017	SANLAATKMSECVLGQSKRVDFCGKGYHLSFPPQSAPHGVVFLHVTYVPAQEKNF TTAPA				1076
Query 1081	ICHDGKAHFPREGVFSNGTHWVFTQRFYEPQIITTDNTFVSGNCDVVIGIVNNTVYDP				1140
Sbjct 1077	ICHDGKAHFPREGVFSNGTHWVFTQRFYEPQIITTDNTFVSGNCDVVIGIVNNTVYDP				1136
Query 1141	LQPELDSFKEELDKYFKNHTSPDVLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDL				1200
Sbjct 1137	LQPELDSFKEELDKYFKNHTSPDVLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDL				1196
Query 1201	QELGKYEQYIKWPWYIWLGFIAGLIAIMVTIMLCMTSCCSCCLKGCCSCGSCCKFDEDD				1260
Sbjct 1197	QELGKYEQYIKWPWYIWLGFIAGLIAIMVTIMLCMTSCCSCCLKGCCSCGSCCKFDEDD				1256
Query 1261	SEPV LKG VKLHYT 1273				
Sbjct 1257	SEPV LKG VKLHYT 1269				

Given the high amino acid identity of this 733 amino acid sequence (except for the PBCS insertion) and the typical coronavirus synonymous to non-synonymous mutation frequency of between three and five synonymous mutations for each non-synonymous mutation,⁴⁷ it was expected that a comparison of the nucleotide sequence for this region between SARS-CoV-2 and RaTG13 would show an almost identical sequence as well.

In fact, when the SARS-CoV-2 nt sequence 23,183-25,384 was compared to the RaTG13 nt sequence 23,165-25,354, the corresponding genome sequence to the 99.6% identical protein sequence above, the nucleotide identity was only 94.2% identical, with 122 synonymous substitutions and only the three non-synonymous substitutions.

⁴⁷ [Comparative genomic analysis](#)

To put this in context a comparison of thirteen other protein coding regions of SARS-CoV-2 and RaTG13 (Text-Table below) shows that the overall synonymous to non-synonymous mutation frequency is 549 synonymous to 109 non-synonymous or a ratio of about 5.0.

Gene	Region of Genome	Total Nucleotides	Synonymous mutations	Non-Synonymous mutations	S/NS	Probability of more than the number of synonymous mutations given the probability of a synonymous mutation is 0.83 (based on all genes pooled)
pp1ab	1-21,239	21,239	659	102	6.5	0.003
pp1ab ABSS	7448-18266	10,818	283	13	21.8	5.73×10^{-12}
Spike Protein RBD	1-1814	1814	131	27	4.9	0.48
Anomalous Base Substitution Segment	23,183-25,384	2201	112	3	37.3	$< 1.0 \times 10^{-7}$
Entire Spike Protein	1-3810	3808	231	41	5.6	0.18
ORF1a polyprotein	1-13,215	13215	440	86	5.2	0.33
ORF3a protein	1-828	828	25	6	4.2	0.56
E Protein	1-228	228	1	0	Infinite	0.83
M Protein	1-669	669	27	3	9.0	0.1
ORF6 Protein	1-186	186	3	0	Infinite	0.17
ORF7a Protein	1-366	366	13	3	4.3	0.47
ORF7b Protein	1-132	132	0	1	0	0.83
ORF8 Protein	1-366	366	5	6	0.8	0.99
Nucleocapsid Phosphoprotein	1-1260	1260	35	4	8.75	0.083

With the exception of the anomalous base substitution segment (ABSS) in the Spike Protein gene and the pp1ab gene, the remainder of the S/SN substitution ratios are consistent with the literature values for coronaviruses. Only two genes or gene regions have a higher S/SN ratio than the ABSS because they have no non-synonymous mutations: the E protein gene with 228 nucleotides and the ORF6 protein gene with 186 nucleotides. Because of the short length of these two genes, the probabilities of the results for the E and ORF6 genes were not significant, with p-values of 0.86 and 0.17, respectively.

The p-value for the ABSS, on the other hand, was highly significant, with a p-value of < 0.0000001 . This strongly suggests a non-natural cause for this base substitution pattern, barring some unknown biological mechanism for such a result.

A second highly anomalous sequence was found in the pp1ab gene. This is about five-times larger than the Spike Protein region and is even more unlikely to have happened naturally, a chance of about one in 100 billion times.

Are there only synonymous mutations in these regions because non-synonymous mutations lead to non-replicative viruses?


A simple explanation for these results would be an extreme criticality for the specific sequences of these regions with respect to infectivity. If a single amino acid change yielded a non-transmissible viral particle that strong negative purification process could explain the above results.

This hypothesis can be immediately rejected based on two observations.

In an examination of over 80,000 SARS-CoV-2 genome sequences, the most common Spike Protein non-synonymous mutation is within the ABSS (D614G) which was identified within weeks of the outbreak in January 2020 and which has become “the dominant virus...in every geographical region.”⁴⁸ Specifically, as of August 28, 2020, GISAID reports that 65,738 full length SARS-CoV-2 genomes of a total of 83,387, or 79%, and comprising the G, GH, and GR clades, contain the D614G SNV. Under real world biological conditions, the ABSSN region has in fact, not a strong negative purification process in operation but in fact a strong positive selection process ongoing.

Secondly, in an analysis of mutations in 63,421 SARS-CoV-2 genomes the Spike Protein amino acid 605 to 1120 region had a total of 7,149 mutations. Fully 5,936 of these mutations (83%) are the above noted D614G non-synonymous change. Of the remaining 1213 mutations, 452 were non-synonymous while 755 were synonymous, a ratio of 1.7. There were also four indels and two stop codon mutations.

The following Text-Figure contains a map of the SARS-CoV-2 genome with the location of amino acid changes that have been found during the worldwide spread noted, with the frequency related to the height of the mark. The two ABSS in pp1ab and SP are marked with red brackets and clearly demonstrate an abundance of non-synonymous mutations in these regions during the human-to-human spread.



[Nextstrain SARS-CoV-2 amino acid change events](#)

Clearly, these regions can tolerate many non-synonymous mutations, rejecting the theory of a criticality for the amino acid sequence of this region. No other natural biological mechanism to explain these results has been identified.

Codon modification, enhancement, or optimization is an example from synthetic biology in which the S/SN ratio is, by design, an anomaly when looked at through the lens of nature

⁴⁸ Biswas NK, Majumder PP. Analysis of RNA sequences of 3636 SARS-CoV-2 collected from 55 countries reveals selective sweep of one virus type. Indian J Med Res. 2020;151(5):450-458. doi:10.4103/ijmr.IJMR_1125_20.

Synonymous codon substitution is a decades old, well known method of enhancing gene expression when cloning exogenous genes in a laboratory experiment. In a paper on the immunogenicity of the SARS-CoV-2 Spike Protein⁴⁹ the following synthetic biology methods were used:

“We used the following structure coordinates of the coronavirus spike proteins from the PDB to define the boundaries for the design of **RBD expression constructs: SARS-CoV-2 (6VSB)**, SARS-CoV-1 (6CRV), HKU-1 (5I08), OC43 (6NZK), 229E (6U7H) NL63 (6SZS). Accordingly, a **codon-optimized gene encoding for S1-RBD [SARS-CoV-1 (318 – 514 aa, P59594), SARS-CoV-2 (331 – 528 aa, QIS60558.1)**, OC43 (329 – 613 aa, P36334.1), HKU-1 (310 – 611 aa, Q0ZME7.1), 229E (295 – 433 aa, P15423.1) and NL63 (480 – 617 aa, Q6Q1S2.1)] containing human serum albumin secretion signal sequence, three purification tags (6xHistidine tag, Halo tag, and TwinStrep tag) and two TEV protease cleavage sites was **cloned into the mammalian expression vector pαH. S1 RBDs were expressed in Expi293 cells** (ThermoFisher) and purified from the culture supernatant by nickel-nitrilotriacetic acid agarose (Qiagen).”

The Genbank alignment (below) confirms that the authentic SARS-CoV-2 Spike Protein sequence (<https://www.ncbi.nlm.nih.gov/nuccore/1798174254>) and the [Synthetic construct SARS CoV-2 spike protein receptor binding domain gene, complete cds](#) are 100% homologous at the protein level:

unnamed protein product						
Sequence ID: Query_33917 Length: 581 Number of Matches: 1						
Range 1: 335 to 532 Graphics ▼ Next Match ▲ Pre						
Score	Expect	Method	Identities		Positives	Gaps
414 bits(1064)	6e-149	Compositional matrix adjust.	198/198(100%)		198/198(100%)	0/198(0%)
Query 331	NITNLCPFGEVFNATRFASVYAWNRKRISNCVADYSVLVNSASFSTFKCYGVSPTKLN					390
Sbjct 335	NITNLCPFGEVFNATRFASVYAWNRKRISNCVADYSVLVNSASFSTFKCYGVSPTKLN					394
Query 391	CFTNVYADSFVIRGDEVQRQIAPGQTGKIADYNYKL PDDFTGCVIAWNSNNLDSKVG					450
Sbjct 395	CFTNVYADSFVIRGDEVQRQIAPGQTGKIADYNYKL PDDFTGCVIAWNSNNLDSKVG					454
Query 451	YLYRLFRLSKNLKPFERDISTEIQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYQPYRV					510
Sbjct 455	YLYRLFRLSKNLKPFERDISTEIQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYQPYRV					514
Query 511	VVLSFELLHAPATVCGPK 528					
Sbjct 515	VVLSFELLHAPATVCGPK 532					

But a comparison of the authentic nucleotide sequence of SARS-CoV-2 to the codon-optimized synthetic construct shows no match using the “highly similar Megablast” algorithm setting. When the alignment algorithm is run in a more relaxed mode the impact of codon optimization in this case can be seen, a 70% homology:

⁴⁹ <https://immunology.sciencemag.org/content/5/48/eabc8413/tab-pdf>

Download

Graphics

Sequence ID: Query_50133

Length: 1746

Number of Matches: 1

Range 1: 1003 to 1595

Graphics

Next Match

Previous Match

Score	Expect	Identities	Gaps	Strand
275 bits(304)	2e-76	419/595(70%)	4/595(0%)	Plus/Plus
Query 22553	AATATTACAACTTGTGCCCTTTTGGTGAAGTTTTTAACGCCACCAGATTGTCATCTGTT	22612		
Sbjct 1003	AACATCACCACATCTGTGCCCTTCGGCGAGGTGTTCAACGCCACAAGATTCGCCTCTGTG	1062		
Query 22613	TATGCTTGGAAACAGGAAGAGAATCAGCAACTGTGTTGCTGATTATTCTGTCTATATAAT	22672		
Sbjct 1063	TACGCCCTGGAACCGGAAGCGGATCAGCAATTGCGTGGCCGACTACAGCGTGCTGTACAAC	1122		
Query 22673	TCCGCATCATTTTC--CACTTTTAAGTGTTATGGAGTGTCTCTACTAAATTAATGATC	22730		
Sbjct 1123	AGCGC--CAGCTTCAGCACCTTCAAGTGCTACGGCGTGTCCCTACCAAGCTGAACGACC	1180		
Query 22731	TCTGCTTTACTAATGTCTATGCAGATTCAATTTGTAATTAGAGGTGATGAAGTCAGACAAA	22790		
Sbjct 1181	TGTGCTTCACCAACGTGTACGCCGACAGCTTCGTGATCAGAGGCGACGAAGTGCGGCAGA	1240		
Query 22791	TCGCTCCAGGGCAAACCTGGAAAGATTGCTGATTATAATTATAAATTACCAGATGATTTTA	22850		
Sbjct 1241	TTGCCCTGGACAGACAGGCAAGATCGCCGATTACAACCTACAAGCTGCCCGACGACTTCA	1300		
Query 22851	CAGGCTGCGTTATAGCTTGAATCTAACAATCTTGATTCTAAGGTTGGTGGTAATTATA	22910		
Sbjct 1301	CCGGCTGTGTGATTGCTTGGAAACAGCAACAACCTGGACAGCAAAGTCGGCGGCAACTACA	1360		
Query 22911	ATTACCTGTATAGATTGTTTAGGAAGTCTAATCTCAAACCTTTGAGAGAGATATTTCAA	22970		
Sbjct 1361	ACTACCTGTACCGGCTGTTCCGGAAGTCCAACCTGAAGCCTTTCGAGCGGGACATCAGCA	1420		
Query 22971	CTGAAATCTATCAGGCCGGTAGCACACCTTGTAAATGGTGTGGAAGGTTTAAATTGTTACT	23030		
Sbjct 1421	CCGAGATCTATCAGGCCGGCAGCACCCCTTGCAATGGCGTGGAAGGCTTCAACTGCTACT	1480		
Query 23031	TTCTTTTACAATCATATGGTTTCCAACCCACTAATGGTGTGTTACCAACCATACAGAG	23090		
Sbjct 1481	TCCCACTGCAGTCTACGGCTTCAGCTTACAACCGCGTGGGCTACCAGCCTTACAGAG	1540		
Query 23091	TAGTAGTACTTTCTTTTGAACCTTCTACATGCACCAGCAACTGTTTGTGGACCTAA	23145		
Sbjct 1541	TGGTGGTGCTGAGCTTCGAGCTGCTGCATGCTCCTGCCACAGTGTGTGGACCTAA	1595		


This is a situation in which there are 176 synonymous changes without a single non-synonymous change and is the genome signature of laboratory-derived synthetic biology. If these sequences were compared for phylogenetic divergence without the knowledge of their artificial construction, this synthetic laboratory experiment would create the impression that these two sequences had diverged in the wild from a common ancestor decades earlier.

The following Table identifies four regions of the RaTG13 and SARS-CoV-2 genomes in which there were a total of 220 synonymous mutations without a single non-synonymous change.

Protein/Gene	Protein Region	Total Nucleotides	Synonymous mutations	NS Mutations
S Protein	605-1124	1557	91	0
pp1ab	3607-4534	2781	66	0
pp1ab	4626-5111	1455	26	0
pp1ab	5113-5828	2145	37	0
Total		7938	220	0

These regions represent over 26% of the entire genome and appear analogous to the outcome expected from the application of a synonymous codon modified, laboratory-derived synthetic biology project. They also represent about one-sixth of the 4% apparent phylogenetic divergence between RaTG13 and SARS-CoV-2.

October GenBank update. On October 13, 2020 the sequence for RaTG13 was updated. For the first time the first 15 nucleotides at the 5' end were present. However, these were not found in a blast of either the RNA-Seq raw reads or the Amplicons. The following email was sent to Dr. Shi asking for an explanation of the fecal specimen composition and the source for the 5' nt data.



At the time of this writing a response has not been received.

Discussion. The foundation of the working hypothesis that the COVID-19 pandemic arose via a natural zoonotic transfer from a non-human vertebrate host to man has been built on two publications: the February 3, 2020 *Nature* paper by Dr. Zheng-Li Shi and colleagues, in which the bat coronavirus RaTG13 is first identified as the closest sequence identity to SARS-CoV-2 at 96.2% and the March 17, 2020 *Nature Medicine* paper entitled, "The proximal origin of SARS-CoV-2," by Andersen *et al.*, in which the Shi *et al.* paper is cited as evidence for a bat origin for the pandemic. In the approximately six months since they were published, these two papers have been cited over 1600- and 200-times on PubMed, respectively.

However, research is beginning to question whether a bat species can be considered a natural reservoir for SARS-CoV-2. A recent paper performed an *in silico* simulation of the SARS-CoV-2 Spike Protein interaction with the cell surface receptor, ACE2, from 410 unique vertebrate species, including 252 mammals.⁵⁰ Among primates, 18/19 have an ACE2 receptor which is

⁵⁰ Broad host range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates Joana Damas, et al. Proc. of the Nat. Acad. of Sci. Aug 2020, 202010146; DOI: 10.1073/pnas.2010146117

100% homologous to the human protein in the 25 residues identified to be critical to infection, including the *Chlorocebus sabaeus* (the Old World African Green monkey) and the rhesus macaques.

It is noteworthy that the laboratory workhorse of coronavirus research is the VERO cell, isolated from a female African Green monkey in 1962, and containing an ACE2 receptor that is 100% homologous to the human ACE2 in the 25 critical amino acids for infectivity.

This *in silico* work was confirmed in the laboratory with respect to rhesus macaques. Within weeks of the identification of SARS-CoV-2, the Wuhan laboratory had demonstrated that the pandemic virus would infect and produce a pneumonia in rhesus macaques.⁵¹

A surprising finding from the ACE2 *in silico* surveillance work was the very poor predicted affinity of the ACE2 receptors in both bats and pangolins. Of 37 bat species studied, 8 scored low and 29 scored very low. As expected by these predictions, cell lines derived from big brown bat (*Eptesicus fuscus*),⁵² Lander's horseshoe bat (*Rhinolophus landeri*), and Daubenton's bat (*Myotis daubentonii*) could not be infected with SARS-CoV-2.⁵³

It is unfortunate that growth of the RaTG13 specimen could not have been attempted in the *Rhinolophus sinicus* primary or immortalized cells generated and maintained in the Wuhan laboratory: kidney primary cells (RsKi9409), lung primary cells (RsLu4323), lung immortalized cells (RsLuT), brain immortalized cells (RsBrT) and heart immortalized cells (RsHeT).⁵⁴ However it should be noted that a synthetically created RaTG13 was reported not to infect human cells expressing *Rhinolophus sinicus* ACE2, providing evidence that RaTG13 may not be a viable coronavirus in a wild bat population.⁵⁵

The other proposed intermediate host, the pangolin, also had predicted ACE-2 affinity that was either low or very low.

A recent paper that examined the high synonymous mutation difference between RaTG13 and SARS-CoV-2 used an *in silico* methodology to suggest that the difference could be largely attributed to the RNA modification system of hosts.⁵⁶ However, the authors do not "(t)he

⁵¹ Infection with Novel Coronavirus (SARS-CoV-2) Causes Pneumonia in the *Rhesus Macaques*. C. Shan et al., Research Square, DOI: [10.21203/rs.2.25200/v1](https://doi.org/10.21203/rs.2.25200/v1). Shan, C., Yao, Y., Yang, X. et al. Infection with novel coronavirus (SARS-CoV-2) causes pneumonia in *Rhesus macaques*. *Cell Res* **30**, 670–677 (2020). <https://doi.org/10.1038/s41422-020-0364-z>

⁵² J. Harcourt et al., Severe acute respiratory syndrome coronavirus 2 from patient with coronavirus disease, United States. *Emerg. Infect. Dis.* **26**, 1266–1273 (2020).

⁵³ M. Hoffmann et al., SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* **181**, 271–280.e8 (2020).

⁵⁴ Zhou, P., Fan, H., Lan, T. et al. Fatal swine acute diarrhoea syndrome caused by an HKU2-related coronavirus of bat origin. *Nature* **556**, 255–258 (2018). <https://doi.org/10.1038/s41586-018-0010-9>.

⁵⁵ Y. Li et al., Potential host range of multiple SARS-like coronaviruses and an improved ACE2-Fc variant that is potent against both SARS-CoV-2 and SARS-CoV-1. *bioRxiv*:10.1101/2020.04.10.032342 (18 May 2020).

⁵⁶ [The divergence between SARS-CoV-2 and RaTG13 might be overestimated due to the extensive RNA modification](#)

limitation of our study is that we were currently unable to provide experimental evidence for the modification on viral RNAs.” The low S/SN ratio of 1.7 in the expansion of SARS-CoV-2 in the human population would argue against a robust host RNA modification mechanism.

In summary, the findings reported here are:

1. Inconsistencies between published papers and interviews as to the source and sequencing history of the original specimen that was claimed to have been collected in 2013 (RaBtCoV/4991) and the specimen for the bat RaTG13 virus. For example, two explanations of the discovery of the close relationship between RaTG13 and SARS-Cov-2, a highly homologous match between the RdRp genes of the viruses noticed in 2020 followed by full genome sequencing, or identification in 2020 of a homologous match to full genome sequencing previously done in 2018. Current publicly available data for RaTG13 from 2017 and 2018 is a set of 33 amplicon sequencing runs but they cover only about 80% of the entire genome. In the *Science* interview Dr. Shi’s says the specimen for RaTG was consumed during sequencing in 2018, but if this is true, the RNA-Seq referred to in the *Nature* paper could not have been performed in 2020. At this time, the Wuhan laboratory has not met the requirements of *Nature* with respect to the sharing of primary and sequence assembly data from their seminal paper¹ and this data should be provided immediately.
2. The specimen from which RaTG13 was reported to have been isolated and which has been repeatedly reported to have been a bat fecal specimen has a taxonomical composition of eukaryotes, bacteria, and viruses that is completely different from a set of nine bat fecal specimens collected in the same field visits by the same laboratory personnel from the Wuhan Institute of Virology. The probability that an authentic fecal specimen could have the composition reported is one in ten million, an impossibly low occurrence. Examination of the strong signals in the RaTG13 specimen identifies both a variety of bat genetic material, some that are not native to China, as well as unexpected species, such as marmots and a red fox. It also contains a telltale 3% primate sequence consistent with VERO cell contamination. I propose that this specimen is apparently either a mislabeled specimen (although I cannot conjure what the field source or specimen would be) or was artificially created in a laboratory.
3. The method-dependent sequence differences between the amplicon data and the RNA-Seq data are about 5% or about 50-times higher than expected as a technical error rate of 0.1%. This is an experimental quality issue that needs to be addressed; no explanation has been offered for this to date. In addition, no assembly methodology has been provided and at least two gaps, totaling over 60 nt, were easily identified.
4. The findings, reported here of a mutational drift of synonymous mutations only between SARS-CoV-2 and RaTG13 in the Spike Protein S1/S2 region and the pp1ab gene that has never been seen in nature before and which has a probability of having occurred by chance of less than one in ten million and one in one billion makes it more likely that, at least for these portions of the RaTG13 genome, comprising over one-

quarter of the entire genome, another process is underway. With the demonstration that codon-enhancement or optimization can produce this unnatural S/SN pattern, some form of laboratory-based synthetic biology was performed on RaTG13, SARS-CoV-2, or both.

Apparently, the entire specimen from which RaTG13 was purported to have been found has been consumed in previous sequencing experiments and the Principal Investigator has stated that no virus has ever been isolated or cultured from the specimen at any time in the past. Given the irregularities and anomalies identified in this paper it seems prudent to conclude that all data with respect to RaTG13 must be considered suspect. As such, reliance of the foundational papers of the origin of SARS-CoV-2 as having arisen from bats via a zoonotic mechanism must be reexamined and questioned.

Paper 2: The February 19, 2020 Lancet paper entitled: “Statement in support of the scientists, public health professionals, and medical professionals of China combatting COVID-19.”

On February 19, 2020 *The Lancet* published a Correspondence entitled “Statement in support of the scientists, public health professionals, and medical professionals of China combatting COVID-19⁵⁷” with 27 public health scientists from eight countries as authors. The statement seems to attempt to settle the question of the origin of SARS-CoV-2 and short circuit further debate, as the second sentence reads: “We stand together to strongly condemn conspiracy theories suggesting that COVID-19 does not have a natural origin.” It goes on to state: “Conspiracy theories do nothing but create fear, rumors, and prejudice that jeopardize our global collaboration in the fight against this virus.”

The letter provided an open solicitation for support and at this time has been signed by at over 20,300 people, as if to purport that science can be advanced through polling and the democratic process.⁵⁸ While it is a truism that conspiracy theories have no place in the academia, legitimate debate should not be foreclosed.

The statement itself provides a more nuanced discussion of the evidence for a zoonotic origin and contains 14 references, eight of which contain data about the COVID-19 pandemic and six of which are governmental policy statements without new data, background articles from 2003 and 2004 on zoonotic diseases, or a virus naming statement by the Coronavirus Study Group (CSG) of the International Committee on Taxonomy of Viruses, which is responsible for developing the official classification of viruses and taxa naming (taxonomy) of the Coronaviridae family. The eight articles with data were written at the end of January or early February, when there were fewer than 10,000 patients.

⁵⁷ [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)30418-9/fulltext#back-bib1](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)30418-9/fulltext#back-bib1)

⁵⁸ This is reminiscent of the story attributed to Albert Einstein by Stephen Hawking in his *Brief History of Time*. According to Hawking, a book was published in 1930 in pre-war Germany entitled, “One Hundred Authors Against Einstein.” When he was asked about the book Einstein is reported to have retorted, “If I were wrong, then one would have been enough!”

An analysis of the evidence for a zoonotic source given in support of the above Statement is contained in Text-Table here. The analysis shows there was very little actual data available at the time to permit reaching such a definitive conclusion. There was also the absence of data or discussion that could support a laboratory origin.

Reference	Statements concerning origin of SARS-CoV-2	Response to statements
1.Gorbalenya AE Baker SC Baric RS et al. Severe acute respiratory syndrome-related coronavirus: the species and its viruses—a statement of the Coronavirus Study Group. bioRxiv. 2020; (published online Feb 11. DOI: 2020.02.07.937862 (preprint).)	A naming statement about SARS-CoV-2. The emergence of SARS-CoV-2 as a human pathogen in December 2019 may thus be perceived as completely independent from the SARS-CoV outbreak in 2002–2003. With respect to novelty, SARS-CoV-2 differs from the two other zoonotic coronaviruses, SARS-CoV and MERS-CoV, introduced to humans earlier in the twenty-first century.	Does not provide data on a potential zoonotic source.
2.Zhou P Yang X-L Wang X-G et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature. 2020; (published online Feb 3.)	The sequences of 2019-nCoV BetaCoV/Wuhan/WIV04/2019 among patient specimens are almost identical and share 79.6% sequence identity to SARS-CoV. Furthermore, we show that 2019-nCoV is 96% identical at the whole-genome level to a bat coronavirus. Pairwise protein sequence analysis of seven conserved non-structural proteins domains show that this virus belongs to the species of SARSr-CoV. The close phylogenetic relationship to RaTG13 provides evidence that 2019-nCoV may have originated in bats.	The bat genome identity of 96% described here, coupled with the known mutation rate of SARS-CoV-2 of about 26/year, implies a lowest common ancestor about 44 years ago.

<p>3.Lu R Zhao X Li J et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. Lancet. 2020; (published online Jan 30.)</p>	<p>Genome sequences of 2019-nCoV sampled from nine patients who were among the early cases of this severe infection are almost genetically identical, which suggests very recent emergence of this virus in humans and that the outbreak was detected relatively rapidly. 2019-nCoV is most closely related to other betacoronaviruses of bat origin, indicating that these animals are the likely reservoir hosts for this emerging viral pathogen.</p>	<p>Figure 1A shows 8 sequences and the consensus sequence. These 8 sequences show 3 with 0 mutations, 2 with 1 mutation, 3 with 2 mutations, and none with more than 2 mutations. Based on current estimates of 1 mutation per human passage, these are at most two human-to-human transfers apart. Importantly, there is no background diversity as would be seen in two or more reservoir-to-human events. Fig 2 states strain Bat-SL-CoVZC45 is 87.6% sequence identity to the human virus, which means a difference of about 3700 mutations or over 70 years from lowest common ancestor.</p>
<p>4.Zhu N Zhang D Wang W et al. A novel coronavirus from patients with pneumonia in China, 2019. NEJM. 2020; (published online Jan 24.)</p>	<p>"more than 85% identity with a bat SARS-like CoV (bat-SL-CoVZC45, MG772933.1) genome published previously. Since the sequence identity in conserved replicase domains (ORF 1ab) is less than 90% between 2019-nCoV and other members of betacoronavirus, the 2019-nCoV — the likely causative agent of the viral pneumonia in Wuhan — is a novel betacoronavirus belonging to the</p>	<p>A >85% identity with a bat coronavirus means the human and bat virus have over 70 years to LCA.</p>

	sarbecovirus subgenus of Coronaviridae family."	
5.Ren L Wang Y-M Wu Z-Q et al. Identification of a novel coronavirus causing severe pneumonia in humans: a descriptive study. Chin Med J. 2020; (published online Feb 11.)	All five patients have sequence homology of 99.8% to 99.9%. These isolates showed 79.0% nucleotide identity with the sequence of SARS-CoV (GenBank NC_004718) and 51.8% identity with the sequence of MERS-CoV (GenBank NC_019843). The virus is closest to a bat SARS-like CoV (SL-ZC45, GenBank MG772933) with 87.7% identity, but is in a separate clade. Surprisingly, RNA-dependent RNA polymerase (RdRp), which is the most highly conserved sequence among different CoVs, only showed 86.3% to 86.5% nt identities with bat SL-CoV ZC45.	Similar to reference 3 comments. Lack of conserved sequencing of the most highly conserved sequence with bat coronavirus would suggest a non-bat source.
6.Paraskevis D Kostaki EG Magiorkinis G Panayiotakopoulos G Tsiodras S Full-genome evolutionary analysis of the novel corona virus (2019-nCoV) rejects the hypothesis of emergence as a result of a recent recombination event. Infect Genet Evol. 2020; (published online Jan 29.)	A BLAST search of 2019-nCoV middle fragment revealed no considerable similarity with any of the previously characterized corona viruses. Bat_SARS-like coronavirus sequences cluster in different positions in the tree, suggesting that they are recombinants, and thus that the 2019-nCoV and RaTG13 are not recombinants. Codon usage analyses can resolve	The middle segment with no similarity to other corona viruses is about 40% of the entire genome. I agree SARS-CoV-2 is not a recombinant of RaTG13. I agree, codon usage analysis here supports the furin binding site insertion as having been invented de novo. A recent recombination event is not necessary for a

	<p>the origin of proteins with deep ancestry and insufficient phylogenetic signal or <u>invented de novo</u>. Our study rejects the hypothesis of emergence as a result of a recent recombination event. Notably, the new coronavirus provides a new lineage for almost half of its genome, with no close genetic relationships to other viruses within the subgenus of sarbecovirus. This genomic part comprises half of the spike region encoding a multifunctional protein responsible also for virus entry into host cells</p>	<p>laboratory derived theory of origin. Statements do not advance a zoonotic origin.</p>
<p>7. Benvenuto D Giovanetti M Ciccozzi A Spoto S Angeletti S Ciccozzi M The 2019-new coronavirus epidemic: evidence for virus evolution. J Med Virol. 2020; (published online Jan 29.)</p>	<p>The epidemic originated in Wuhan, China. A phylogenetic tree has been built using the 15 available whole genome sequences of 2019-nCoV, 12 whole genome sequences of 2019-nCoV, and 12 highly similar whole genome sequences available in gene bank (five from the severe acute respiratory syndrome, two from Middle East respiratory syndrome, and five from bat SARS-like coronavirus). >97% maximum likelihood match to Bat SARS-like virus 2015 (Fig 1) is noted. The SARS and MERS viruses are excluded as a source of SARS-CoV-2. These results do not</p>	<p>A 3% genome distance from the noted bat virus to human is about 34 years at 26 mutations per year, the in-human mutation rate. Predicted a future mutation like the D614G mutation which is more infective.</p>

	exclude the fact that further mutation due to positive selective pressure, led by the epidemic evolution, could favor an enhancement of pathogenicity and transmission of this novel virus.	
8.Wan Y Shang J Graham R Baric RS Li F Receptor recognition by novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS. J Virol. 2020; (published online Jan 29.)	Based on predicted RBD-host ACE2 receptor affinities, civet, mice, and rats are ruled out as source species. Pigs, ferrets, cats, and nonhuman primates contain largely favorable 2019-nCoV-contacting residues in their ACE2. SARS-CoV was isolated in wild palm civets near Wuhan in 2005, and its RBD had already been well adapted to civet ACE2.	The potential nonhuman primate ACE2 usage is noted. Consistent with a laboratory origin from VERO cells , a monkey kidney cell line. It expresses an ACE2 that permits SARS-CoV-2 infection, making it a possible source for the virus. A common tissue culture cell line for SARS virus research.
9.US Center for Disease Control and Prevention Coronavirus disease 2019 (COVID-19) situation summary. https://www.cdc.gov/coronavirus/2019-nCoV/summary.html Date: Feb 16, 2020 Date accessed: February 8, 2020	Rarely, animal coronaviruses can infect people and then spread between people such as with MERS-CoV, SARS-CoV, and now with this new virus, named SARS-CoV-2. The SARS-CoV-2 virus is a betacoronavirus, like MERS-CoV and SARS-CoV. All three of these viruses have their origins in bats. The sequences from U.S. patients are similar to the one that China initially posted, suggesting a likely single, recent emergence of this virus from an animal reservoir.	There are no data to support these statements about bats as the source for SARS-CoV-2.

10. Andersen KG Rambaut A Lipkin WI Holmes EC Garry RF The proximal origin of SARS-CoV-2. http://virological.org/t/the-proximal-origin-of-sars-cov-2/398 Date: Feb 16, 2020 Date accessed: February 17, 2020	See Table 2.	See Table 2.
11. Bengis R Leighton F Fischer J Artois M Morner T Tate C The role of wildlife in emerging and re-emerging zoonoses. <i>Rev Sci Tech.</i> 2004; 23: 497-512	In one pattern, actual transmission of the pathogen to humans is a rare event but, once it has occurred, human-to-human transmission maintains the infection for some period of time or permanently. Some examples of pathogens with this pattern of transmission are human immunodeficiency virus/acquired immune deficiency syndrome, influenza A, Ebola virus and severe acute respiratory syndrome.	This 2004 paper describes the pattern of rare animal-to-human transmission followed by human-to-human spread as an example of the SARS virus. It does not address the origin of SARS-CoV-2.
12. Woolhouse ME Gowtage-Sequeria S Host range and emerging and reemerging pathogens. <i>Emerg Infect Dis.</i> 2005; 11: 1842-1847	Emerging and reemerging pathogens are disproportionately viruses, with 37% being RNA viruses. Emerging and reemerging pathogens more often are those with broad host ranges that often encompass several mammalian orders and even nonmammals. For pathogens that are minimally transmissible within human populations (R_0 close to 0), outbreak size is determined largely by the number of introductions from the reservoir. For pathogens that are highly transmissible within human populations	This 2005 article has good general information about looking broadly for the reservoir species(s), identifies RNA viruses as a major source of human epidemics, predicts a large outbreak size for a high R_0 virus, but does not address the origin of SARS-CoV-2.

	($R_0 \gg 1$), outbreak size is determined largely by the size of the susceptible population.	
<p>13.NASEM The National Academies of Science Engineering and Medicine of the USA. NAS, NAE, and NAM presidents' letter to the White House Office of Science and Technology Policy.</p> <p>https://www.nationalacademies.org/incudes/NASEM%20Response%20to%20OOSTP%20re%20Coronavirus_February%206,%202020.pdf Date: Feb 6, 2020 Date accessed: February 7, 2020</p>	<p>The closest known relative of 2019-nCoV appears to be a coronavirus identified from bat-derived samples collected in China.⁴ The experts informed us that additional genomic sequence data from geographically- and temporally-diverse viral samples are needed to determine the origin and evolution of the virus. Samples collected as early as possible in the outbreak in Wuhan and samples from wildlife would be particularly valuable. Understanding the driving forces behind viral evolution would help facilitate the development of more effective strategies for managing the 2019-nCoV outbreak and for preventing future outbreaks.</p>	<p>Agree. If additional genomic sequence data is available from geographically- and temporally-diverse viral samples are needed to determine the origin and evolution of the virus this should be made publicly available.</p>
<p>14.WHO Director-General's remarks at the media briefing on 2019 novel coronavirus on 8 February 2020.</p> <p>https://www.who.int/dg/speeches/detail/director-general-s-remarks-at-the-media-briefing-on-2019-novel-coronavirus---8-february-2020 Date: Feb 8, 2020 Date accessed: February 18, 2020</p>	<p>A general statement about the emerging pandemic without reference to the origin of SARS-CoV-2</p>	<p>There is no data about the origin of the pandemic.</p>

In November 2020 the Watchdog group, US Right-to-Know, reported the following with respect to the *Lancet* article:⁵⁹

“Emails obtained by U.S. Right to Know show that a statement in *The Lancet* authored by 27 prominent public health scientists condemning “conspiracy theories suggesting that COVID-19 does not have a natural origin” was organized by employees of EcoHealth Alliance, a non-profit group that has received millions of dollars of U.S. taxpayer funding to genetically manipulate coronaviruses with scientists at the Wuhan Institute of Virology.”

“The emails obtained via public records requests show that EcoHealth Alliance President Peter Daszak drafted the *Lancet* statement, and that he intended it to “not be identifiable as coming from any one organization or person” but rather to be seen as “simply a letter from leading scientists”. Daszak wrote that he wanted “to avoid the appearance of a political statement.”

A separate, worrisome article entitled, “Peter Daszak’s EcoHealth Alliance Has Hidden Almost \$40 Million In Pentagon Funding And Militarized Pandemic Science,”⁶⁰ seems to indicate a serious conflict of interest with respect to Dr. Daszak’s participation in any investigations on the origin of SARS-CoV-2.

Paper 3: The March 17, 2020 article in *Nature Medicine* entitled “The proximal origin of SARS-CoV-2” by Andersen et al.^{61, 62}

According to the journal, this article is in the 99th percentile (ranked 2nd) of the 312,683 tracked articles of a similar age in all journals and the 99th percentile (ranked 1st) of the 147 tracked articles of a similar age in *Nature Medicine*. The metrics also indicate it has been accessed over five million times. It is clearly the most cited paper and since its title and topic are the origin of the pandemic it clearly has an outsized influence on the topic.

The following statements form the evidence in the article of the natural origin of CoV-2:

- “While the analyses above suggest that SARS-CoV-2 may bind human ACE2 with high affinity, computational analyses **predict that the interaction is not ideal** and that the RBD sequence is different from those shown in SARS-CoV to be optimal for receptor binding. Thus, the high-affinity binding of the SARS-CoV-2 spike protein to human ACE2 is **most likely the result of natural selection on a human or human-like ACE2** that permits another optimal binding solution to arise. **This is strong evidence that SARS-CoV-2 is not the product of purposeful manipulation.**” [emphasis added.]

⁵⁹ <https://usrtk.org/biohazards-blog/ecohealth-alliance-orchestrated-key-scientists-statement-on-natural-origin-of-sars-cov-2/>

⁶⁰ <https://www.independentsciencenews.org/news/peter-daszaks-ecohealth-alliance-has-hidden-almost-40-million-in-pentagon-funding/>

⁶¹ <https://www.nature.com/articles/s41591-020-0820-9>

⁶² Two non-peer reviewed analyses are included here because they provide a nearly line-by-line analysis. They unfortunately include occasional colorful language but the content is worth noting: <https://harvardtothebighouse.com/2020/03/19/china-owns-nature-magazines-ass-debunking-the-proximal-origin-of-sars-cov-2-claiming-covid-19-wasnt-from-a-lab/> ; <https://www.youtube.com/watch?v=HmSCMb8Nds4>

- A later analysis of over 3800 possible substitutions of amino acids in a 200 amino acid receptor binding region, much larger than the small, selective region referred to in this paper, shows that CoV-2 is 99.5% optimized for binding to the ACE-2 receptor. This near perfect binding has never been seen before in a recent interspecies transmission jump.
- “Polybasic cleavage sites have not been observed in related ‘lineage B’ betacoronaviruses, although other human betacoronaviruses, including HKU1 (lineage A), have those sites and predicted O-linked glycans. Given the level of genetic variation in the spike, **it is likely that SARS-CoV-2-like viruses with partial or full polybasic cleavage sites will be discovered in other species.**” [emphasis added.]
 - As of the writing of this manuscript no other lineage B (sarbecovirus) has been found to have a furin site. In addition, the furin site of CoV-2 has the unusual -CGG-CGG- codon dimer, which has never been seen in an analysis of 58 other sarbecoviruses, that is, 580,000 codons. Since recombination between subgenera of beta coronaviruses is rare, or unknown, there is no source for the CGG-CGG dimer via a natural recombination event.
- “The acquisition of polybasic cleavage sites by HA has also been observed after repeated passage in cell culture or through animals.”
 - It is curious why the above statement did not lead to a hypothesis somewhere in the article about a similar mechanism on CoV-2, a clear indication of a laboratory origin.
- “It is improbable that SARS-CoV-2 emerged through laboratory manipulation of a related SARS-CoV-like coronavirus.”
 - This conclusory statement is unsupported by evidence.
- “Furthermore, if genetic manipulation had been performed, one of the several reverse-genetic systems available for betacoronaviruses would **probably have been used.** However, the genetic data irrefutably show that SARS-CoV-2 is not derived from any previously used virus backbone.” [emphasis added.]
 - There is no explanation for why a prior backbone would necessarily be used. All synthetic biology chimera coronaviruses created in the past as published in prior papers have each used a unique backbone with no particular pattern in backbone selection. Each backbone was selected for the particular needs of those current experiments. This non-repeating prior pattern of reverse-genetic systems makes the above statement untenable. And with 16,000+ reported coronavirus specimens at the WIV it is entirely reasonable a non-published virus could have been used.

- “Natural selection in an animal host before zoonotic transfer. For a precursor virus to acquire both the polybasic cleavage site and mutations in the spike protein suitable for binding to human ACE2, **an animal host would probably have to have a high population density (to allow natural selection to proceed efficiently)** and an ACE2-encoding gene that is similar to the human ortholog.” [emphasis added.]
 - The paragraph discusses the pangolin as the possible intermediate host but at the time of this manuscript the coronavirus data from pangolins has been discredited. This author agrees with statement that selection of the two unique features of CoV-2 require a high population density of the animal host. Of course, in the laboratory the animal hosts for either *in vitro* cell culture experiments or in animal experiments are a single species at high density.
- Natural selection in humans following zoonotic transfer. “It is possible that a progenitor of SARS-CoV-2 jumped into humans, acquiring the genomic features described above through adaptation during **undetected human-to-human transmission**. Once acquired, these adaptations would enable the pandemic to take off and produce a sufficiently large cluster of cases to trigger the surveillance system that detected it.” [emphasis added.]
- “Studies of banked human samples could provide information on whether such cryptic spread has occurred. Further serological studies should be conducted to determine the extent of prior human exposure to SARS-CoV-2.”
 - As will be shown in later sections, this prior undetected human-to-human transmission would be evident in archived specimens from before the fall of 2019. In both SARS-CoV-1 and MERS, this prior seroconversion averaged about 0.6% with almost 5% among workers exposed to the intermediate hosts. At the time of the writing of this manuscript, in limited sampling of archived specimens there has been no seroconversion detected. The author believes there are thousands of archived specimens from Wuhan taken in the fall of 2019 and these should be immediately examined for evidence of seroconversion. Since finding seroconversion among these specimens would be strong evidence for a zoonotic origin and not a laboratory accident, the absence of any information from China on this important evidence is hard to understand.
- Selection during passage. “Basic research involving passage of bat SARS-CoV-like coronaviruses in cell culture and/or animal models has been ongoing for many years in biosafety level 2 laboratories across the world, and there are documented instances of laboratory escapes of SARS-CoV. We must therefore examine the possibility of an inadvertent laboratory release of SARS-CoV-2.”

- “In theory, it is possible that SARS-CoV-2 acquired RBD mutations during adaptation to passage in cell culture, as has been observed in studies of SARS-CoV.”
- “New polybasic cleavage sites have been observed only after prolonged passage of low-pathogenicity avian influenza virus in vitro or in vivo. Furthermore, a hypothetical generation of SARS-CoV-2 by cell culture or animal passage would have required prior isolation of a progenitor virus with very high genetic similarity, **which has not been described**. Subsequent generation of a polybasic cleavage site would have then required repeated passage in cell culture or animals with ACE2 receptors similar to those of humans, **but such work has also not previously been described**.” [emphasis added.]
 - The authors correctly describe a method for CoV-2 to have been generated in the laboratory and then dismiss it because the work has not been published previously. As active scientists themselves, the authors must know how disingenuous this sounds. Almost by definition elite scientists, like Dr. Shi of the WIV, work in secret until the publication of any given line of research. As the say, the absence of evidence cannot be used as evidence of its absence.
 - A peer-reviewed paper⁶³ entitled, “Might SARS-CoV-2 Have Arisen via Serial Passage through an Animal Host or Cell Culture? A potential explanation for much of the novel coronavirus’ distinctive genome,” provides a compelling argument that serial passage in the laboratory might indeed have been the manner in which CoV-2 acquired many of its devastating traits.
- “Although the **evidence shows that SARS-CoV-2 is not a purposefully manipulated virus**, it is currently impossible to prove or disprove the other theories of its origin described here. However, **since we observed all notable SARS-CoV-2 features, including the optimized RBD and polybasic cleavage site, in related coronaviruses in nature, we do not believe that any type of laboratory-based scenario is plausible**.” [emphasis added.]
 - This author could identify no prior evidence in the paper to warrant saying it is not a purposefully manipulated virus. There is also no evidence that would point to a purposely manipulated virus.
 - The evidence in the paper shows that no prior zoonotic interspecies transmission has ever had an RBD as optimized as the CoV-2 RBD for the human ACE2. The evidence also shows that there is no natural source for the polybasic cleavage site (PCS). No other member of the subgenera to which CoV-2 belongs has a PCS. Since these are the only coronaviruses from which recombination could supply a polybasic cleavage site, the data in this paper refutes the natural origin.

⁶³ <https://onlinelibrary.wiley.com/doi/full/10.1002/bies.202000091>

- The belief statement concerning a laboratory-based scenario would be closer to the evidence if it was professed with, “despite evidence which is consistent with a laboratory-based scenario.”

Based on the author’s analysis of the paper, the following email was sent to the lead author:



Soon after this email was written Dr. Andersen blocked the author from following his Twitter account. A reply to the above email was never received.

Conclusion. Three high visibility papers were published between January and May 202 which purported to settle the question of the origin of SARS-CoV-2 as a zoonotic transmission and not a laboratory accident. The analysis above concludes that these papers are not persuasive. The

author has elected to not use evidence within these papers to change the prior likelihood of a zoonotic versus laboratory origin. They are presented here as neutral evidence that supports neither theory.

Likelihood from initial state is unchanged following this evidence analysis:

Zoonotic origin (98.8%) and laboratory origin (1.2%)

Evidence. SARS-like infections among employees of the Wuhan Institute of Virology in the fall of 2019

The State Department of the United States issued the following statement on January 15, 2021⁶⁴:

“1. Illnesses inside the Wuhan Institute of Virology (WIV):

- The U.S. government has reason to believe that several researchers inside the WIV became sick in autumn 2019, before the first identified case of the outbreak, with symptoms consistent with both COVID-19 and common seasonal illnesses. This raises questions about the credibility of WIV senior researcher Shi Zhengli’s public claim that there was “zero infection” among the WIV’s staff and students of SARS-CoV-2 or SARS-related viruses.”

There is no additional evidence to support either parties position in the above statement. The U.S. Government statement would be considered hearsay in a court of law and probably not admissible. The veracity of Dr. Shi’s statement above could be called into question due to other inconsistencies in some of her testimony, as reported elsewhere in this document.

At this time, the above evidence cannot be used to change the likelihood of either theory about the origin of SARS-CoV-2. The statement is kept within this analysis with the hope that in the future new information will come to light that could make this evidence a useful addition to the overall analysis.

Likelihood from initial state is unchanged following this evidence analysis:

Zoonotic origin (98.8%) and laboratory origin (1.2%)

⁶⁴ <https://2017-2021.state.gov/fact-sheet-activity-at-the-wuhan-institute-of-virology//index.html>

Evidence. A Bayesian Analysis of one aspect of the SARS-CoV-2 origin, where the first recorded outbreak occurred, increases the probability of a laboratory origin.

Introduction. The two competing hypotheses of the origin of SARS-CoV-2 as a natural, zoonotic spillover event versus a laboratory-acquired infection (LAI) or other laboratory accident each had supporting evidence from the very beginning of the pandemic.

On the one hand, about 40% of early patients with COVID-19 had an association with the Hunan Seafood Market in Wuhan. Since this mirrored SARS-CoV-1, where markets selling civet cats were determined to be the origin of that human epidemic, the natural origin hypothesis seemed logical. The Chinese CDC have now ruled out the market as a source for the outbreak.

On the other hand, the laboratory origin hypothesis also had an early beginning with the fact that the outbreak began adjacent to the only high security, BSL-4 laboratory in all of China, and one of the top coronavirus research centers in the world, was the Wuhan Institute of Virology (WIV). The hospitals of the first COVID patients were very close to the WIV.

This evidence statement is taken from an article applying a Bayesian analysis to the hypothesis that the proximal origin of SARS-CoV-2 was an uncontrolled⁶⁵ release from a laboratory using, as evidence, one aspect of the SARS-CoV-2 origin story — where the first recorded outbreak occurred.⁶⁶

Hypothesis: The first recorded outbreak of SARS-CoV-2 in the human population occurred in a city that is also home to a virology laboratory that actively performs research on closely related viruses.

In this case, the city is Wuhan, and the virology laboratory is run by the Wuhan Institute of Virology.

Analysis. This analysis set the likelihood of a laboratory escape (the prior probability the hypothesis was true) at three values, 0.01%, 0.1%, and 1.0%. The second term was the conditional probability of the evidence, given that the hypothesis is actually false. This was set at 0.01. Finally, the third term was the conditional probability of the evidence, given the hypothesis is true. This was set, biasing to the natural origin, at 0.71.

Results. The paper provides the three-by-three cube of results for the three parameters of interest.

The ardent sceptic's probability begins at 0.01% and the revised estimate is no more than 0.05% or 5/10000. It applies to someone who was initially very skeptical about a lab origin (0.01% probability), who believes there is no more than 51% chance that an uncontrolled release of a highly contagious disease would lead to a local outbreak, and who thinks there was at least a

⁶⁵ By using the term uncontrolled release, the author was specifically excluding from consideration the possibility that the pathogen was deliberately released from the laboratory.

⁶⁶ <https://jonseymour.medium.com/a-bayesian-analysis-of-one-aspect-of-the-sars-cov-2-origin-story-where-the-first-recorded-1fbdcbea0a2b>

10% chance that a natural outbreak of a virus native to Yunnan would have occurred in Wuhan before any place else.

On the other extreme, is the ardent believer who started with at least a 1% belief in a laboratory outbreak, is 100% certain that an uncontrolled laboratory release would result in a local outbreak and believes that the probability that a natural outbreak of a virus native to Yunnan would occur in Wuhan before any place else is less than 0.1%. The ardent believer's revised belief is that the probability that the Wuhan outbreak was caused by an uncontrolled laboratory release changes from 1% to at least 91%.

In the center, is the so-called "central" observer who accepts that the central values for each of the parameter ranges are reasonable estimates of the true values of the probability being estimated. The central observer started with an initially skeptical belief in the hypothesis of 0.1%, believes that average citizen in Wuhan was as likely as any other citizen of China to be the initial vector of the virus into the human population and believes that there is no more or less than a 71% chance that an uncontrolled release from a laboratory of a highly contagious pathogen such as SARS-CoV-2 would result in a local outbreak as opposed to an outbreak in some other location. The central observer's revised belief in the hypothesis is 6.8%. If the central observer began with a 1% belief in a laboratory origin, this analysis would change that to 41.8%.

Conclusion. For purposes of this analysis and to be as conservative as possible, the assumptions will be that there is at least a 1% prior belief in a laboratory outbreak (because that was our starting probabilities), but there is no more than a 51% chance that an uncontrolled release of a highly contagious disease would lead to a local outbreak, and that there was at least a 10% chance that a natural outbreak of a virus native to Yunnan would have occurred in Wuhan before any place else. Using these assumptions, the initial likelihood of a 1% laboratory origin changes to 4.9%.

Starting likelihood from initial state: Zoonotic origin (98.8%) and laboratory origin (1.2%)

Adjusted likelihood: Zoonotic origin (95.1%) and laboratory origin (4.9%)


Evidence: Lack of seroconversion in Wuhan and Shanghai. Summary of evidence:


- A hallmark of zoonotic infections (vertebrate animal host-to-human microbial infection) is repeated, abortive jumps into humans over time until sufficient 'human-adapted' mutations permit efficient human-to-human spread and further evolution

Summary


- A hallmark of zoonotic infections (vertebrate animal host-to-human microbial infection) is repeated, abortive jumps into humans over time until sufficient 'human-adapted' mutations permit efficient human-to-human spread and further evolution
- A record of these abortive jumps can be found in archived specimens of either healthy individuals or patients with an influenza-like illness that are examined for residual virus, by PCR, or seroconversion, by antibody tests
- This permits the classification of an epidemic as a zoonotic event without having to find a viral host
- Four studies of SARS-CoV-1 and MERS in a total of 12,700 human specimens shows an average seroconversion prevalence of 0.6%
- Two studies, one in Wuhan (n=520) looking for seroconversion and one in Shanghai (n=1271), using both PCR and seroconversion, found no SARS-CoV-2 positive specimen before the first week of January
- Using the combined prevalence (0.6%) of SARS-CoV-1 and MERS, both known zoonotic epidemics, and the sensitivity of the PCR assay used (94.4%), the negative predictive value of these results is $\geq 91\%$

- A record of these abortive jumps can be found in archived specimens of either healthy individuals or patients with an influenza-like illness that are examined for residual virus, by PCR, or seroconversion, by antibody tests





- This permits the classification of an epidemic as a zoonotic event without having to find a viral host
- A laboratory accident is a situation in which there are no prior exposures within the human population as shown in the Figure below:



- Four studies of SARS-CoV-1 and MERS in a total of 12,700 human specimens shows an average seroconversion prevalence of 0.6%

SARS-related Virus Predating SARS Outbreak, Hong Kong

SARS-CoV-1 began in fall of 2002 in southern China

Patient Population	Serum samples collected in May 2001 from 938 healthy adults in Hong Kong	48 confirmed SARS patients diagnosed in February and March 2003 in Guangdong
Civet CoV > SARS-CoV-1 Seropositivity	13	0
SARS-CoV-1 > Civet CoV Seropositivity	4	48
Total	17 out of 938 = 1.8%	48 out of 48 = 100%

Pre-epidemic seroprevalence in the adult community

Prevalence is 0.6% for SARS-CoV-1 and MERS in 12,700 specimens

Epidemic	Nature of the Study	Seropositivity	Reference
SARS-CoV-1	Archived specimens from healthy adults in Hong Kong collected two years before CoV-1 were tested for Ab to civet or human CoV	17/938	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3322899/
MERS	Archived human sera collected in 2011 was tested for MERS-CoV S1-specific antibodies by ELISA	1/90	https://www.sciencedirect.com/science/article/pii/S1876034120300010#fig0010
SARS-CoV-1	Serum specimens collected from military recruits from the People's Republic of China in 2002 were tested for SARS-CoV-1 antibodies.	11/1621	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1074388/
MERS	Between Dec 1, 2012, and Dec 1, 2013, 10,009 individual serum samples were tested for anti-MERS-CoV antibodies in regions without cases.	15/10,009	https://pubmed.ncbi.nlm.nih.gov/25863564/
SARS-CoV-1	Serum samples that were collected from 42 individuals during 2001-2002, before the SARS outbreak, and tested for IgG antibody against SARS-CoV.	28/42	https://arxiv.org/ftp/arxiv/papers/1305/1305.2659.pdf

Pre-epidemic seroprevalence in MERS shepherds and slaughterhouse workers is higher

Prevalence is 2.3% (2/87) in shepherds and 3.6% (5/140) in slaughterhouse workers

Reference: <https://pubmed.ncbi.nlm.nih.gov/25863564/>

- Two studies, one in Wuhan (n=520) looking for seroconversion and one in Shanghai (n=1271), using both PCR and seroconversion, found no SARS-CoV-2 positive specimen before the first week of January

Pre-epidemic seroconversion has never been seen for SARS-CoV-2

Epidemic	Nature of the Study	Seropositivity	References
SARS-CoV-2	RNA PCR from 1271 nasopharyngeal swab samples, as well as the prevalence of IgM, IgG, and total antibodies against SARS-CoV-2 in 357 matched serum samples collected from hospitalized patients with influenza-like illness between 1 December 2018 and 31 March 2020 in Shanghai Ruijin Hospital. First positive was January 25, 2020.	0/1271	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7473166/pdf/TEMI_9_1785952.pdf
SARS-CoV-2	Re-analysed 5200 throat swabs collected from patients in Wuhan with influenza-like-illness from 6 October 2019 to week one January 2020 and found no positive specimens for SARS-CoV-2 RNA by quantitative PCR.	0/520	https://www.nature.com/articles/s41564-020-0713-1
CoV-2 Studies Combined		0/1791	Probability is one in 14,881

- Using the combined prevalence (0.6%) of SARS-CoV-1 and MERS, both known zoonotic epidemics, and the sensitivity of the PCR assay used (94.4%), the negative predictive value of these results is $\geq 91\%$

Negative Predictive Value of SARS-CoV-2 PCR Test	
BioGerm PCR Test has a sensitivity of 94.4%	
SARS & MERS Seroconversion	0.60%
PCR Sensitivity	94.40%
Negative Predictive Value Calculation	$<0.6/(0.6 + 0.054)$
Negative Predictive Value	$\geq 91\%$

Here, the negative predictive value (NPV) represents the probability that a CoV-2 is not a zoonosis, given the negative seroconversion findings.

Subjective Discount Factor: 90% (a one in 10 chance this is wrong). This is a subjective value.

The change in origin likelihoods from this evidence and the calculations are shown in the Text-Table below.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin
Starting likelihood	0.951	0.049
Negative predictive value of lack of seroconversion	0.91	
Reduced by 90% Subjective Discount Factor	$0.91 \times 0.9 = 0.82$	
Impact of this evidence	Reduces the likelihood of ZO by 82/18 or 4.6-fold. For every 100 tests, a true ZO would be seen 18 times and a non-ZO would be seen 82 times	
Impact of evidence calculation	$0.951/4.6 = 0.207$	
Normalize this step of analysis	$0.207/(0.207 + 0.049) = 0.809$	$0.049/(0.207 + 0.049) = 0.191$

Adjusted likelihood: Zoonotic origin (80.9%) and laboratory origin (19.1%)

Evidence: Lack of posterior diversity for SARS-CoV-2 compared to MERS and SARS-CoV-1

- The earliest stages of human CoV-1 and MERS infections were characterized by viral genome base diversity as expected for multiple, independent jumps from a large and diverse intermediate host population into humans.
- Combining MERS and CoV-1 studies, out of the earliest 255 human infections in which virus genome sequences are available, 137 could not be rooted in a prior human-to-human infection and so are attributed to an independent intermediate host-to-human infection.⁶⁷
- That is about 54% non-human-to-human transmission.
- On the other hand, Ralph Baric has written⁶⁸ that CoV-2 is different: “SARS-CoV-2 probably emerged from bats, and early strains identified in Wuhan, China, showed limited genetic diversity, which suggests that the virus **may have been introduced from a single source.**” [emphasis added.]
- With CoV-2, there are 249 viral genomes in GISAID from Hubei province, where Wuhan is located, collected between Dec 24, 2019 and Mar 29, 2020.
- From Dec 24, 2019 to November 2020, there are 1001 genomes sequenced from all of China and 198,862 worldwide.
- For CoV-2, every single genome sequence is rooted in the first sequence from the PLA Hospital in Wuhan.
- Not one case of posterior diversity.
- Using the frequency of non-rooted genome diversity seen with MERS and CoV-1, about 50:50 or a coin toss, the probability that CoV-2 is a zoonotic pandemic with 0/249 genomes is the chance of tossing a coin 249 times and getting heads every time!
- Mathematically that is nonexistent; specifically, one in 10 with 84 zeros.
- Since Wuhan had approximately 500,000 cases during the time interval of this sampling, the potential sampling error of testing only 249/500,000 or 0.05% is significant. This sampling error, while large, is unable to obliterate the overwhelming odds that this did not arise from an intermediate host in Wuhan.
- Therefore, to permit continued evidence analysis, this finding will be set at the boundary of customary statistical significance, a p-value of 0.05 or a 1 in 20 likelihood that this is zoonotic.

⁶⁷ <https://elifesciences.org/articles/31257#abstract> ;
https://www.researchgate.net/publication/225726653_Molecular_phylogeny_of_coronaviruses_including_human_SARS-CoV ; <https://science.sciencemag.org/content/300/5624/1394/tab-pdf> ;
<https://pubmed.ncbi.nlm.nih.gov/14585636/> ;
<https://www.microbiologyresearch.org/content/journal/jgv/10.1099/vir.0.016378-0?crawler=true> ;
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7118731/>

⁶⁸ <https://www.nejm.org/doi/10.1056/NEJMcibr2032888>

Detailed explanation

A fundamental difference between a laboratory and a non-laboratory acquired zoonotic disease, the imprint of phylogenetic diversity through pre-human spread within the source population, can be examined by the posterior diversity of human cases with no *a priori* knowledge of an intermediate host.

MERS. The MERS epidemic has been documented to have arisen from the initial jump from bats to camels, a three-to-five-year expansion within the camel population in which mutational diversity arose by random mistakes, and then a jump into humans. This model of spread predicts that there would, at some point, be additional jumps from other camels into other patients, and a pattern of “posterior diversity,” would be found in the human specimens. If the COVID-19 pandemic arose by a similar mechanism the same pattern would be seen. The following Text-Table contains such data.

Phylogenetic Feature	MERS	SARS-CoV-2
Posteriority Diversity	28/30 (93%)	0
No Posteriority Diversity	2/30 (7%)	7666
Time from first patient to first example of posterior diversity	About 60 days	None at >120 days
Depth of posterior diversity to first patient	>365 days	None

The study of MERS noted above was published in 2013 in Lancet⁶⁹ in an article entitled, “Transmission and evolution of the Middle East respiratory syndrome coronavirus in Saudi Arabia: a descriptive genomic study.” Thirty specimens were used in the analysis. The features of a camel-to-human zoonotic epidemic are easily identified. Specimens taken within sixty days of the first patient, “Patient Zero,” began to show a background diversity that could not be traced back through Patient Zero. The analysis of all thirty, in fact, documented that 93% were transmitted directly from the camel intermediate reservoir. And looking only at the “background” diversity permitted a calculation of the last common ancestor for the spread within the camel population of over 365 days.

A study of SARS-CoV-2⁷⁰ available May 5, 2020 and entitled, “Emergence of genomic diversity and recurrent mutations in SARS-CoV-2,” looked at 7666 patient specimens from around the world for phylogenetic diversity. The authors state: “There is a robust temporal signal in the data, captured by a statistically significant correlation between sampling dates and ‘root-to-tip’ distances for the 7666 SARS-CoV-2 ($R^2 = 0.20$, $p < .001$). Such positive association between sampling time and evolution is expected to arise in the presence of measurable evolution over the timeframe over which the genetic data was collected.” This conclusion also argues against a

⁶⁹ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3898949/>

⁷⁰ <https://www.sciencedirect.com/science/article/pii/S1567134820301829>

MERS-like pattern of posterior diversity. In fact, the 95% upper bound for the probability of no posterior diversity being seen in SARS-CoV-2, given the data in MERS, is 3.9×10^{-4} .



The finding of posterior diversity in MERS was seen quickly, that is, within 60 days of the first patient and in only 30 specimens. In this study of COVID-19 the cutoff date of the 7666 specimens was April 19, 2020 or approximately 140 days after the first documented case. The lack of posterior diversity in COVID-19 at a much later date than what was seen with MERS also argues against a non-laboratory source for this pandemic.

A useful avenue of future research for those working to find an animal source for COVID-19 would be new mathematical models or statistical methods that might find a “hidden” signal of posterior diversity in the current data set which shows none. And given access to the unprecedented quantity of human data for COVID-19 which can be mined via bioinformatics, efforts to find the “missing link” in the wild through search and sample should be a second priority to mining the human specimen data set.


SARS-CoV-1. A similar pattern of clinical cases that do not show a common ancestor in the human population but instead is evidence of posterior diversity is shown in the Text-Table on the left for SARS-CoV-1⁷¹ compared to CoV-2 on the right⁷². SARS-CoV-1 shows clusters of cases in humans that are connected only by phylogenetic branches that reach back in time (all of the branches inside the purple box. This is because of the extensive mutational background created while being in the intermediate host, the civet. With CoV-2 on the right, every clinical case descends from the first clinical case, in the 19A clade. There are no background mutations to account for. I will show elsewhere that the first Clade A patient was at the PLA Hospital about 3 km from the WIV.

⁷¹ <https://pubmed.ncbi.nlm.nih.gov/14585636/>

⁷² <https://nextstrain.org/>



Given the rate of mutations of 22.8 per year for CoV-2 as shown in the Nextstrain graph below and a sequencing accuracy of about two calls per genome, CoV-2 could not have spent more than a few weeks in an intermediate host before a pattern of background mutations would be identified as posterior diversity. In the laboratory a pure culture on a single genome is used and the CoV-2 pattern is most consistent with a single pure culture infection a first human.




Non-zoonotic evolution. In a hypothetical in which there was a singular event in which one genetically pure virus infected one person and then the epidemic grew the development of the genetic diversity would have a clear, identifiable pattern: every new mutation would only appear on a background of the previous mutations.

The mutations in this virus are literally a personal tag. The general mutation rate leads to one mutation per patient. So, by definition, Patient Zero will have just one mutation. And then the 2-4 people that patient passes it to will have that mutation and then will add a new one, and so on. As time goes by two things happen: each patient gets a new mutation of their own and they pass on all the mutations of the past.

Since the virus has 29,900 nt and the mutation rate, as shown in this graph prepared by NextStrain is 26 mutations per year, there is very little chance a mutation will appear and then later get undone. By carefully going back in time, it is possible to literally name each person at each generation by the one (on average) new mutation they have and all of those that went before.

This graph of mutations on the Y-axis shows them gradually increasing and the color coding shows where they came from. In this infection, they only came from a previous patient and from the next previous patient and so on.



A NextStrain graphic.

How is that different from MERS, which was passed from camels to humans in a true zoonotic process?


In a true zoonotic spread to humans there is usually an initiating species (in MERS it is bats), and then an intermediate species (in MERS it is camels), and then it moves to humans, either because of a new “enabling mutation” or for a non-domestic species, a chance encounter, and Source Zero and Patient Zero meet, and a cross species event occurs. But “Source Zero” doesn’t stop there with one infection in one human; the virus also transmits itself vertically into the intermediate species. Source Zero also creates a vertical infection in the camels. Whether it is mild or not doesn’t matter. The new human jumping gene is moving into a very diverse population of viruses, who have themselves been evolving since the first bat to camel transmission.

What is the outcome in terms of a test to show this is happening?

The diversity of the virus in humans becomes great, and the spots where the mutations occur don’t match up to MERS Patient Zero like they do in COVID-19. In MERS, the virus in Patient Zero and the virus in a later infection are not direct descendants but cousins and only descended from an earlier virus that spent time in another camel population, collecting random mutations until it got the one it needed to infect humans, and then it begins again.

The chart below, from Lancet. 2013 Dec 14; 382(9909): 1993–2002, shows just how this works. The patient at Bisha is the earliest case in this chart (Patient Zero in the red circle). But notice, no other case comes from that patient. The viruses have such a diverse genetic background they appear to only be related to the Bisha virus with a posterior timeline of about one year. Their background is in the green boxes and it skips Patient Zero.

.



Even without knowing that camels are the zoonotic source for MERS, this data, from clinical sample only and without any field work in cave or camels, is all you need to know that this arose in the wild.

A paper just appeared with this analysis for a region of China and the posterior genomic diversity indicated a single starting point on December 1, 2019 for all cases. There was no posterior diversity. At this point with over 322,000 full genomes sequenced⁷³ and all showing an additive pattern of mutations and with none showing background diversity before the known appearance in Wuhan, the only conclusion is that there is no reservoir of genetic diversity.

On January 26, 2020 in an article in *Science* written by Jon Cohen, Kristian Andersen, an evolutionary biologist at the Scripps Research Institute who had analyzed sequences of CoV-2 to try to clarify its origin said: “The scenario of somebody being infected outside the market and then later bringing it to the market is one of the three scenarios we have considered that is still consistent with the data. It’s entirely plausible given our current data and knowledge.”

The negative predictive value of finding no posterior diversity in CoV-2 with 322,000 total infections sequenced, over 1000 in China, is 95%

Subjective Discount Factor: 95% (a one in 20 chance this is wrong)

⁷³ <https://www.gisaid.org/>

Below is the impact of the pack of posterior diversity on the likelihood of a zoonotic versus laboratory origin

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin
Starting likelihood	0.809	0.191
Negative predictive value of lack of posterior diversity	0.95	
Reduced by 95% Subjective Discount Factor	$0.95 \times 0.95 = 0.90$	
Impact of this evidence	Reduces the likelihood of ZO by 90/10 or 9-fold. For every 100 tests, a true ZO would be seen 10 times and a non-ZO would be seen 90 times	
Impact of evidence calculation	$0.809/9 = 0.085$	
Normalize this step of analysis	$0.085/(0.085 + 0.191) = 0.308$	$0.191/(0.085 + 0.191) = 0.692$

Adjusted likelihood: Zoonotic origin (30.8%) and laboratory origin (69.2%)

Evidence: Opportunity.

The Wuhan Institute of Virology has publicly disclosed that by 2017 it had developed the techniques to collect novel coronaviruses, systematically modify the receptor binding domain to improve binding or alter zoonotic tropism and transmission, insert a furin site to permit human cell infection, make chimera and synthetic viruses, perform experiments in humanized mice, and optimize the ORF8 gene to increase human cell death (apoptosis).

Wuhan Institute of Virology scientists maps RBD and then takes a civet coronavirus that won't infect human cells, changes two amino acids in the receptor binding domain & it infects human cells.⁷⁴



THE JOURNAL OF BIOLOGICAL CHEMISTRY
© 2005 by The American Society for Biochemistry and Molecular Biology, Inc.

<http://www.paper.edu.cn>
Vol. 280, No. 33, Issue of August 19, pp. 29588–29595, 2005
Printed in U.S.A.

Identification of Two Critical Amino Acid Residues of the Severe Acute Respiratory Syndrome Coronavirus Spike Protein for Its Variation in Zoonotic Tropism Transition via a Double Substitution Strategy*

Received for publication, January 19, 2005, and in revised form, June 16, 2005
Published, JBC Papers in Press, June 24, 2005, DOI 10.1074/jbc.M500662200

**Xiu-Xia Qu,^{a,b} Pei Hao,^{b,c} Xi-Jun Song,^{a,b} Si-Ming Jiang,^{a,b} Yan-Xia Liu,^a Pei-Gang Wang,^a
Xi Rao,^a Huai-Dong Song,^d Sheng-Yue Wang,^e Yu Zuo,^a Ai-Hua Zheng,^a Min Luo,^a
Hua-Lin Wang,^f Fei Deng,^f Han-Zhong Wang,^f Zhi-Hong Hu,^f Ming-Xiao Ding,^a
Guo-Ping Zhao,^{c,e,g,h} and Hong-Kui Deng^{a,i}**

From the ^aDepartment of Cell Biology and Genetics, College of Life Sciences, Peking University, Beijing 100871, the ^bBioinformation Center/institute of Plant Physiology and Ecology/Health Science Center, Shanghai institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, the ^dState Key Laboratory for Medical Genomics/Pôle Sino-Français de Recherche en Sciences du Vivant et Génomique, Ruijin Hospital Affiliated with the Shanghai Second Medical University, Shanghai 200025, the ^eChinese National Human Genome Center, 250 Bi Bo Road, Zhang Jiang High Tech Park, Shanghai 201203, the ^fState Key Laboratory of Virology, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan, 430071, and the ^gState Key Laboratory of Genetic Engineering/Department of Microbiology, School of Life Science, Fudan University, Shanghai 200433, China

Baric & Shi at WIV take bat coronavirus that won't infect human cells, change S746R to add an ARG at S1/S2 site to make furin-like cleavage site, & the new coronavirus infects human cells.⁷⁵

Baric & Shi of WIV create completely synthetic coronavirus from bat spike & mouse adapted backbone that no treatment, monoclonal antibody, or vaccine will touch.⁷⁶

- “Using the SARS-CoV reverse genetics system², we generated and characterized a chimeric virus expressing the spike of bat coronavirus SHC014 in a mouse-adapted SARS-CoV backbone.
- The results indicate that group 2b viruses encoding the SHC014 spike in a wild-type backbone can efficiently use multiple orthologs of the SARS receptor human angiotensin

⁷⁴ <http://www.paper.edu.cn/scholar/showpdf/NUT2kN0INTT0gxeQh>

⁷⁵ <https://jvi.asm.org/content/jvi/89/17/9119.full.pdf>

⁷⁶ <https://pubmed.ncbi.nlm.nih.gov/26552008/>

converting enzyme II (ACE2), replicate efficiently in primary human airway cells and achieve in vitro titers equivalent to epidemic strains of SARS-CoV.

- Additionally, in vivo experiments demonstrate replication of the chimeric virus in **mouse lung with notable pathogenesis**.
- Evaluation of available SARS-based immune-therapeutic and prophylactic modalities revealed poor efficacy; both monoclonal antibody and vaccine approaches failed to neutralize and protect from infection with CoVs using the novel spike protein.
- On the basis of these findings, we **synthetically re-derived an infectious full-length SHC014** recombinant virus and demonstrate robust viral replication both in vitro and in vivo.”

This study was conducted, with permission, during the gain of function moratorium put in place by NIH in 2014:

“These studies were initiated before the US Government Deliberative Process Research Funding Pause on Selected Gain-of-Function Research Involving Influenza, MERS and SARS Viruses (<http://www.phe.gov/s3/dualuse/Documents/gain-of-function.pdf>). This paper has been reviewed by the funding agency, the NIH. Continuation of these studies was requested, and this has been approved by the NIH.”

Drs. Daszak and Shi becomes world's expert on ORF8 induced apoptosis by CoVs in human cells (HeLa) & maximizing lethality.⁷⁷

The full-length **ORF8** protein of SARS-CoV is a luminal endoplasmic reticulum (ER) membrane-associated protein that induces the activation of ATF6, an ER stress-regulated transcription factor that activates the transcription of ER chaperones involved in protein folding [35]. We amplified the **ORF8** genes of Rf1, Rf4092 and WIV1, which represent three different genotypes of bat SARSr-CoV **ORF8** (**S3C Fig**), and constructed the expression plasmids. All of the three **ORF8** proteins transiently expressed in HeLa cells can stimulate the ATF6-dependent transcription. Among them, the WIV1 **ORF8**, which is highly divergent from the SARS-CoV **ORF8**, exhibited the strongest activation. The results indicate that the variants of bat SARSr-CoV **ORF8** proteins may play a role in modulating ER stress by activating the ATF6 pathway. In addition, the **ORF8a** protein of SARS-CoV from the later phase has been demonstrated to induce apoptosis [28]. In this study, we have found that the **ORF8a** protein of the newly identified SARSr-CoV Rs4084, which contained an 8-aa insertion compared with the SARS-CoV **ORF8a**, significantly triggered apoptosis in 293T cells as well.

This paper also demonstrates the collection of 64 novel bat coronaviruses from caves in southern China, including Yunnan where Dr. Shi has said is the location of the bat ancestor of CoV-2.

This evidence is necessary for a laboratory origin hypothesis in which genetic manipulation to create CoV-2 is a precursor to a laboratory accident. However, it does not per se, provide

⁷⁷ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5708621/>

increased weight in favor of a laboratory origin. It is however provided here to be a guide for the kinds of investigations to be conducted if access to the WIV records is ever provided.

Likelihood from prior state is unchanged following this evidence analysis:

Zoonotic origin (30.8%) and laboratory origin (69.2%)

Evidence and Motive for laboratory furin site insertion:

A key to infectivity of coronaviruses is the addition, in nature or the laboratory, of a furin cleavage site (FCS) at the S1/S2 junction of the Spike Protein.

Furin cleavage sites (FCS) have been widely understood to be important for many viral infections, including HIV, influenza, and others. It has also been widely understood before now that lineage B coronaviruses do not have FCS.

It was therefore surprising when an examination of SARS-CoV-2 Spike Protein found an insertion of a 12-nt, 4-AA sequence near the junction of the S1/S2 subunits which creates a furin site that is essential to human infectivity and transmission. As expected from previous work, no lineage B (sarbecovirus) coronavirus has this feature. This is the most difficult “molecular fingerprint” of SARS-CoV-2 to explain having been acquired in the wild and for that reason there are no even passingly feasible theories.

One database of whole genome sequences of 386 coronaviruses was devoid of furin cleavage sites.⁷⁸ Another database of 2956 genomes of sarbecovirus strains sequences shows that none have a furin site.⁷⁹ This is a highly significant finding with a probability that sarbecovirus has a furin site in the wild of one in about 985.⁸⁰

It has been known since 1994 that viral glycoproteins can be cleaved by secreted proteases, including furin.⁸¹ Even before that, in 1992, it was known the peptide sequence R-X-K/R-R in surface glycoproteins was required for avian influenza viruses of Serotype H7 pathogenesis.⁸² The first paper using furin inhibitors to define a role for an FCS in coronavirus-cell fusion was published in 2004.⁸³

Since that time, it has become common practice to insert FCS during laboratory gain-of-function experiments to increase infectivity. The following Text-Table illustrates the scope of just a few of the experiments conducted, with the hyperlink to the paper in column one.

URL for Paper	Title of Paper
One	Characterization of a panel of insertion mutants in human cytomegalovirus glycoprotein B.
Two	Insertion of the two cleavage sites of the respiratory syncytial virus fusion protein in Sendai virus fusion protein leads to enhanced cell-cell fusion and a decreased dependency on the HN attachment protein for activity.

⁷⁸ <https://academic.oup.com/bioinformatics/article/36/11/3552/5766118>

⁷⁹ <https://academic.oup.com/database/advance-article/doi/10.1093/database/baaa070/5909701>

⁸⁰ When a series of samples are taken and none produce the result expected, the probability that this is a false negative finding can be estimated by taking the number of samples and dividing by three. Here, 2956 sarbecoviruses without a single furin site is a probability of one in 2956/3 or 985.

⁸¹ <https://www.ncbi.nlm.nih.gov/pubmed/8162439>

⁸² <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7172898/pdf/main.pdf>

⁸³ <https://www.ncbi.nlm.nih.gov/pubmed/15141003>


Three	Recombinant Sendai viruses expressing fusion proteins with two furin cleavage sites mimic the syncytial and receptor-independent infection properties of respiratory syncytial virus.
Four	Amino acid substitutions and an insertion in the spike glycoprotein extend the host range of the murine coronavirus MHV-A59
Five	Induction of IL-8 release in lung cells via activator protein-1 by recombinant baculovirus displaying severe acute respiratory syndrome-coronavirus spike proteins: identification of two functional regions.
Six	Coronaviruses as vectors: stability of foreign gene expression.
Seven	Experimental infection of a US spike-insertion deletion porcine epidemic diarrhea virus in conventional nursing piglets and cross-protection to the original US PEDV infection.
Eight	Minimum Determinants of Transmissible Gastroenteritis Virus Enteric Tropism Are Located in the N-Terminus of Spike Protein.
Nine	Reverse genetics with a full-length infectious cDNA of the Middle East respiratory syndrome coronavirus.
Ten	Construction of a non-infectious SARS coronavirus replicon for application in drug screening and analysis of viral protein function
Eleven	A severe acute respiratory syndrome coronavirus that lacks the E gene is attenuated in vitro and in vivo.

The creation in the wild of a coronavirus FCS that is used as an example of what might have happened in SARS-CoV-2 is uninformative. In this case, a strain of influenza, in which a new polybasic site appears spontaneously leads to increased infectivity and lethality,⁸⁴ was reported by Tse *et al.* 2014. The mechanism of the FCS acquisition in this paper is an RNA polymerase dependent stuttering at a small, constrained loop in which one or more A nt were inserted, removing the strain in the loop and inserting an AAA codon which represents the basic amino acid lysine. No such method exists for the insertion of arginine, the amino acid in the CoV-2 furin site that needs to be created.

The insert generates a canonical 20 AA furin site sequence. In 2011 Tian et al.⁸⁵ published an analysis of 126 furin cleavage sites from three species: mammals, bacteria and viruses. The analysis showed that when the furin sites are recorded as a 20-residue motif, a canonical structure emerges. It includes one core cationic region (eight amino acids, P6–P2') and two flanking solvent accessible regions (eight amino acids, P7–P14, and four amino acids, P3'–P6').

⁸⁴ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3911587/>

⁸⁵ <https://www.nature.com/articles/srep00261>



This figure above shows the 20-AA of the furin motif in SARS-CoV-2 (in green) with the P14 to P6' AA positions marked with the cleavage site being the amide bond between P1-R and the P1' residue. The motif is color coded with the requirements (in most cases, except for the positively charged AA requirements, most position requirements can be relaxed).

With the insertion, all 20 residues obey the rules as established by Tian. Since there are 20^4 different 4-AA peptides or 160,000 choices, it is remarkable that the 4 AA insert created a sequence that contained a small or cationic AA (8 AA/20 qualify), a cationic AA (3/20), another cationic AA (3/20), and a small AA (5/20) in that order. In fact, there are only 360 or the total or about 0.2% of all four amino acid inserts that would be expected to follow the exact rules for furin substrates. Of course, given the increase in infectivity SARS-CoV-2 has over other coronaviruses that do not have a well-designed furin cleavage site, selection pressure would drive this rare mutational event once it happened randomly. It would also be a likely choice for a laboratory designed furin cleavage site created *de novo*.

Based on the evidence that there are no furin cleavage sites in 2956 sarbecovirus (beta coronavirus) genome sequences⁸⁶, the likelihood that CoV-2 acquired the furin site from a wild sarbecovirus is one in 985 or 0.001. Because this is highly significant, we will use the conservative rule established in the beginning and use a likelihood of 0.05 for this evidence.

Subjective Discount Factor. 95% confidence (only a one in 20 chance this is wrong). Below is the calculation of the Bayesian adjustment.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin
Starting likelihood	0.308	0.692
Negative predictive value of a lack of furin sites in sarbecovirus genomes	0.95	
Reduced by 95% Subjective Discount Factor	$0.95 \times 0.95 = 0.90$	
Impact of this evidence	Reduces the likelihood of ZO by 90/10 or 9-fold. For every 100 tests, a true ZO would be seen 10 times and a non-ZO would be seen 90 times	
Impact of evidence calculation	$0.308/9 = 0.034$	
Normalize this step of analysis	$0.034/(0.034 + 0.692) = 0.047$	$0.692/(0.692 + 0.034) = 0.953$

Adjusted likelihood. Zoonotic origin (4.7%), laboratory origin (95.3%).

⁸⁶ <https://academic.oup.com/database/advance-article/doi/10.1093/database/baaa070/5909701>

Evidence: Codon usage can distinguish insertion events in the wild from those created in the laboratory.

Not only is the insertion of an FCS peptide unique among lineage B coronaviruses, the nt sequence used for the process is more broadly unique among coronaviruses in general, regardless of lineage:

-CCT-CGG-CGG-GCA-

I will now use synonymous codon bias methods to try to inform the question of the origin of SARS-CoV-2.

Because of the redundancy of the genetic code, more than one 3-nt sequence specifies any given amino acid. For example, there are six codons that specify arginine, R. The frequencies with which such synonymous codons are used are unequal and have coevolved with the cell's translation machinery to avoid excessive use of suboptimal codons that often correspond to rare or otherwise disadvantaged tRNAs. This results in a phenomenon termed "synonymous codon bias," which varies greatly between evolutionarily distant species and possibly even between different tissues in the same species.

Decades of research has identified that all life forms, viruses, bacteria, and humans alike, use the codons in a signature pattern of frequency which can be used to identify a particular sequence of RNA or DNA as human or non-human; viral or non-viral.


In this way, viruses in nature and scientists in the laboratory, with different goals and motivations, make distinguishing codon usage decisions which can sometimes provide a fingerprint of their source.

The Text-Table below contains the arginine codon usage for two populations, pooled data for SARS-CoV 2003 and related viruses and 13 Sars-CoV-2 human specimens from widely dispersed locations.

Codon	SARS-CoV 2003 and ten other evolutionary related viruses in the Nidovirales	SARS-CoV-2 from 13 Geo-locations
CGG	0.09	0.09
CGA	0.44	0.37
CGC	0.72	0.37
AGG	0.9	1.07
CGU	1.77	1.63
AGA	2.08	2.48

Since these values are of a type of multiplicative scale, they were fit using a log-normal distribution, which appears appropriate (although the sample size is small). Using the log mean and standard deviation and this distribution, the probability of finding a CGG codon is about 0.024. Assuming they are independent the probability of finding a CCG-CCG codon pair is effectively 0.024^2 or 0.00058. This is a likelihood of about one in 1700.

The following Figure shows the RSCU for the amino acids that comprise the new furin cleavage site in SARS-CoV-2. As one can see, the RSCU values are similar to each other with the exception of the RR dimer insert, which have a very low RSCU of 0.09.



The RSCU value for the CGG codon for R of 0.09 was taken from a 2004 paper of the RSCU for SARS-CoV 2003 and ten other evolutionary related viruses in the *Nidovirales* and is confirmed by 13 SARS-CoV-2 specimens obtained from diverse geographic locations. If one assumes that the RSCU observations are independent and that the probability distribution of these measurements is Gaussian (normal; a reasonable assumption), then one can calculate the probability of obtaining a result as small as 0.09. Removing the two 0.09 values, then the mean and standard deviation of the remaining values are 1.275 and 0.4992, respectively. Then the probability of a single 0.09 value is 0.0088. However, there are two 0.09 values. If we assume that these are independent findings, then the probability of both values being seen is 0.0088^2 or 7.7×10^{-5} . Using the RSCU of 0.2 from the Table above does not change the immense improbability of the usage of a CGGCGG codon pair in the wild.

Single Arginine CGG codon usage analysis suggests this will not be found in the wild.

The codon usage for SARS-CoV-2, like most coronaviruses studied, has a bias toward AT and away from GC nucleotides. The frequency of third position G use in CoV-2, for example, is 13%, 21%, 17%, and 16% for the spike protein, envelope, membrane, and nucleocapsid protein, respectively.

In that context, the scarcity of the CGG genome in SARS-CoV-2 and related coronaviruses, the relative synonymous codon usage, determined by the method of Behura and Severson,⁸⁷ was calculated and tabulated below. The color coding is blue for underutilized codons (RSCU < 1.0) and red for overutilized codons (RSCU > 1.0); light blue for RSCU values of 0.60 to 0.99 and

⁸⁷ <https://www.ncbi.nlm.nih.gov/pubmed/22889422>

light red for RSCU of 1.01 to 1.60. The highest RSCU usage of CGG is 1.21 in the membrane protein in the MERS virus but zero in SARS-CoV-2.

RSCU	SARS-CoV-2	Beta CoV Pangolin	SARS CoV	Bat SARS CoV	MERS CoV
Spike	0.29	0	0.19	0.08	0.25
Envelope	0	0	0	0	0
Membrane	0	0.35	0.74	0.24	1.21
Nucleocapsid	0.41	0.16	0.03	0.04	0.8

Looking at these five coronaviruses:

The largest structural protein of the coronaviruses is the spike protein, with 1273 amino acids. In SARS-CoV-2 there are 42 R residues, with only one RR dimer, the one in the insert that created SARS-CoV-2.

As a reminder none of these related coronaviruses have the 12-nucleotide insertion that forms the putative furin site in CoV-2. Interestingly, the pangolin coronavirus has no CGG residues in the spike protein. The significance of this is it makes the acquisition of this insert from pangolin by recombination impossible.

The smallest structural protein, the envelope protein, has 75 amino acids, including three R residues, but has no CGG codons in any of the related coronaviruses examined.

The SARS-CoV-2 membrane protein has 441 amino acids, 14 R residues and no CGG codons. Among related coronaviruses, this is the most unique finding of the four proteins for SARS-CoV-2 since the other four coronaviruses all utilize CGG to some extent in this protein. In the case of the MERS virus, this protein is the only occurrence in which this codon is overutilized.

The nucleocapsid protein has 418 amino acids and is responsible for packing the RNA genome. As expected for the role of R in protein-RNA interactions, it has 29 R residues and four RR dimers. None of the dimers use the CGGCGG sequence.

The nt usage of the 12-nt insert which forms the FCS cleavage site has a probability this sequence was selected for in the wild of one in 129,870.

A blast search was performed for the 12-nt inserted sequence and adjacent extensions and only the SARS-CoV-2 sequences were identified.

Shortening the search to just the two CGG-CGG codons was only slightly more fruitful. The Text-Table below shows the frequency of the middle half of the insert, CGGCGG, across the genomes of all seven known human coronaviruses, as well as a specimen bovine coronavirus and the bat and pangolin coronaviruses with greatest homology to SARS-CoV-2. Only a single example, outside of the Spike Protein gene, has been found.

Furin PBCS sequence	Beta Coronavirus		Total Arginine Dimers Anywhere	CGGCGG in Spike Protein *	CGGCGG Anywhere in genome *	CCGCCG Anywhere in genome
SRRKRRS	Human CoV-HKU1	GenBank: KF686346.1	12	0	0	0
KRRSRRA	Bovine CoV-Quebec	GenBank: AF220295.1	12	0	0	0
PRRARSV	SARS-CoV-2 Wuhan reference sequence GenBank: NC_045512.2		16	1; nt 23,606	0	0
PRSVRS	MERS-CoV	NCBI Reference Sequence: NC_019843.3	21	0	0	0
NRRSRGA	Human CoV-OC43	London/2011 GenBank: KU131570.1	16	0	0	0
None	Human CoV-229E	GeneBank: KF514433.1	15	0	0	0
None	Human CoV NL63	NCBI Reference Sequence: NC_005831.2	9	0	0	0
None	SARS-CoV 2003 ZJ0301 from China	GenBank: DQ182595.1	17	0	0	0
None	Bat coronavirus RaTG13	GeneBank: MN996532.1	11	0	1; nt 9394	0
None	Pangolin PCoV_GX-P4L	GenBank: MT040333.1	10	0	0	0
Total			139	1	0	0
* - Includes both in phase codons as well as out of phase, frameshift codons.						

To understand what this means for the search for the zoonotic source for SARS-CoV-2, a statistical approach was taken. Using the data from the nine viruses other than SARS-COV-2 there was a single incidence of the CGGCGG found in the bat coronavirus. Assuming 10,000 codons per genome, the frequency of CGGCGG in coronaviruses can be estimated at 2 per 45,000 codons or 4×10^{-5} . Therefore, the frequency of finding the center half of the SARS-CoV-2 insert is very small. This is consistent with the strong bias in all coronaviruses to place an A/U nt in the third codon position.

The last column above, the presence of -CCG-CCG- in these coronaviruses was included because it is the hybridization sequence partner for the negative strand sequence, which arises during genome replication. This eliminates the possibility of a strand jumping event to generate a CGGCGG codon dimer.

A similar analysis for the spike protein gene can be done. Since there are no instances of CGGCGG in the spike protein genome, and the gene is 3819 nucleotides long, there are 636 pairs of codons. Thus, over the 9 other viruses, there are 5724 pairs of codons and no cases of the CGGCGG pair. To calculate the upper bound on the probability of such a pair from these data, one can use the Poisson “Rule of Three”, which yields a value of $3/5724$ or 0.00052 with 95% confidence. Now examining the SARS-COV-2 genome, there was one instance of the pair in question out of 636 pairs. The probability of this happening if the true rate of this occurrence for a beta coronavirus is 0.00052 is 0.044. Obviously for smaller assumed rates of this occurrence, this would result in probabilities less than 0.044.

Since the 12-nt insert has been found nowhere in the coronavirus genomic universe, examining over 300,000 sequences and using the Poisson “Rule of Three” again, the upper bound on the frequency that it exists in nature is less than one in 100,000 with 95% confidence.

This observation in conjunction with the lack of finding the 12-nt sequence in any candidate zoonotic species makes unlikely a natural source for the virus. One line of investigation to establish a wild source for this infection would be to find a coronavirus strain with the 12-nt sequence somewhere in nature. The fact that 10 of the 12 nts are either G or C coupled, the documented bias against GC suggests this search would be futile.

Based on these analyses that demonstrate that the finding of a -CGG-CGG- codon pair in the furin site of CoV-2 is a highly improbable event, and using the conservative value of a one in 20 chance (the value for a p-value of 0.05), one can recalculate the likelihood of the choice between a zoonotic origin and a laboratory origin.

Subjective Discount Factor. 95% confidence (only a one in 20 chance this is wrong). Below is the calculation of the Bayesian adjustment.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin
Starting likelihood	0.047	0.953
Negative predictive value of the absence of the -CGG-CGG- pair in any coronavirus in nature	0.95	
Reduced by 95% Subjective Discount Factor	$0.95 \times 0.95 = 0.90$	
Impact of this evidence	Reduces the likelihood of ZO by 90/10 or 9-fold. For every 100 tests, a true ZO would be seen 10 times and a non-ZO would be seen 90 times	
Impact of evidence calculation	$0.047/9 = 0.005$	
Normalize this step of analysis	$0.005/(0.005 + 0.953) = 0.005$	$0.953/(0.953 + 0.005) = 0.995$

Adjusted likelihood. Zoonotic origin (0.5%), laboratory origin (99.5%).

Codon optimization by recombinant methods (that is, to bring a gene's synonymous codon use into correspondence with the host cell's codon bias) has been widely used to improve cross-species expression of protein.

Though the opposite objective of reducing expression by intentional introduction of suboptimal synonymous codons has not been extensively investigated, isolated reports indicate that replacement of natural codons by rare codons can reduce the level of gene expression in different organisms. For example, one approach to vaccine development is to create an attenuated virus which comprises a modified viral genome containing nucleotide substitutions engineered in multiple locations in the genome, wherein the substitutions introduce synonymous de-optimized codons.

In US Patent 9,476,032⁸⁸ titled, “Attenuated viruses useful for vaccines,” they state: “In one high-priority redesigned virus, most or all Arg codons are changed to CGC or **CGG** (the top two frequent human codons). This does not negatively affect translation.” The patent contains numerous codon usages optimized for vaccine production, including the SARS-CoV virus, and in fact they use the CGG-CGG codon pair 45 times.

Beginning with a paper in 2004,⁸⁹ one motivation for codon-optimized SARS genomes is stated here: “The gene encoding the S protein of SARS-CoV contains many codons used infrequently in mammalian genes for efficiently expressed proteins. We therefore generated a codon-optimized form of the S-protein gene and compared its expression with the S-protein gene of the native viral sequence. S protein was readily detected in HEK293T cells transfected with a plasmid encoding the codon-optimized S protein.”

Since that time, human optimized codons have been frequently used for coronavirus research, mostly in gain-of-function experiments. In that context the “molecular fingerprint” of CGG for R is one of those common laboratory reagent gene manipulators.

Other examples:

Examples of the use of CGG codon for arginine in coronavirus research	Reference
SARS was genetically modified to improve ACE2 binding using "human optimized" codons, like CGG for arginine, to grow better in the laboratory. The strains were more infective.Preparation of SARS-CoV S protein pseudotyped virus. “The full-length cDNA of	Wu, K. et al. Mechanisms of Host Receptor Adaptation by Severe Acute Respiratory Syndrome

⁸⁸ <http://patft.uspto.gov/netacgi/nph-Parser?Sect1=PTO1&Sect2=HITOFF&d=PALL&p=1&u=%2Fnetacgi%2FPTO%2Fsrchnum.htm&r=1&f=G&l=50&s1=9476032.PN.&OS=PN/9476032&RS=PN/9476032>

⁸⁹ <https://www.ncbi.nlm.nih.gov/pubmed/15367630>

the SARS-CoV S gene was optimized according to human codon usage and cloned into the pCDNA3.1(+) vector (Invitrogen). The resulting “humanized” S sequence was identical with that of strain BJ01 at the amino acid level.”	Coronavirus. J Biol Chem. 2012 Mar 16; 287(12): 8904–8911.
Predictions of future evolution of a virus are a difficult, if not completely impossible, task. However, our detailed structural analysis of the host receptor adaptation mutations in SARS-CoV RBD has allowed us to predict, design, and test optimized SARS-CoV RBDs that may resemble future evolved forms of the virus. "RBD might evolve into the human-optimized form by acquiring two mutations at the 442 and 472 position." SARS-CoV-2 acquired the mutation at position 472.	Fang Li. Receptor recognition and cross-species infections of SARS coronavirus. Antiviral Res. 2013 Oct; 100(1): 246–254.
Plasmid encoding a codon-optimized form of the SARS-CoV S protein of the TOR2 i	Wenhui Li, Chengsheng Z, et al., Receptor and viral determinants of SARS-coronavirus adaptation to human ACE2. EMBO J. 2005 Apr 20; 24(8): 1634–1643.
The gene encoding the S protein of SARS-CoV contains many codons used infrequently in mammalian genes for efficiently expressed proteins. We therefore generated a codon-optimized form of the S-protein gene and compared its expression with the S-protein gene of the native viral sequence. S protein was readily detected in HEK293T cells transfected with a plasmid encoding the codon-optimized S protein (Fig. (Fig.1).1). No S protein was detected in cells transfected with a plasmid encoding the native S-protein gene.	Moore, MJ, Dorfman, T. Retroviruses Pseudotyped with the Severe Acute Respiratory Syndrome Coronavirus Spike Protein Efficiently Infect Cells Expressing Angiotensin-Converting Enzyme 2. J Virol. 2004 Oct; 78(19): 10628–10635.
Published in 2019 by Dr. Zhengli-Li Shi , entitled "Origin and evolution of pathogenic coronaviruses," reviews genetic optimized SARS viruses using human codons.	Cui, J, Fang, L. Origin and evolution of pathogenic coronaviruses. Nat Rev Microbiol. 2019; 17(3): 181–192.
In 2006, Montana scientists put a synthetic furin cleavage site into a SARS coronavirus by adding an R residue at position R667. They write: "We show that furin cleavage at the modified R667 position generates discrete S1 and S2 subunits and potentiates membrane fusion activity." Mutations were introduced by using	Follis, KE, York, J, Nunberg, JH. Furin cleavage of the SARS coronavirus spike glycoprotein enhances cell–cell fusion but does not affect virion entry. Virology 350 (2006) 358–369

QuikChange mutagenesis (Stratagene) ⁹⁰	
Identification of murine CD8 T cell epitopes in codon-optimized SARS-associated coronavirus spike protein is the title of a paper that shows that the expression of spike protein in vitro was greatly increased by expression cassette optimization.	Zhia, Y, Kobinger, GP, Jordan, H, et al. Identification of murine CD8 T cell epitopes in codon-optimized SARS-associated coronavirus spike protein
As for the human clec4C_1 and mouse clec14A, they showed very similar profiles with spike genes, especially with bat SARS-CoV, in the arginine coding groups, showing the high RSCU values over 2.50 in AGA.	Ahn, I, Jeong, B-J, Son, HS. Comparative study of synonymous codon usage variations between the nucleocapsid and spike genes of coronavirus, and C-type lectin domain genes of human and mouse. Experimental & Molecular Medicine volume 41, pages 746–756, 2009.

One relevant paper,⁹¹ in which arginine residues were being inserted into bovine herpesvirus-1, used primers to create RR dimers with nine separate -CGG-CGG- codon pairs. as testament to their broad use in the Wuhan Institute of Virology laboratory.

Scientists from the Wuhan Institute of Virology provided the scientific community with a technical bulletin on how to make genetic inserts in coronaviruses and proposed using the very tool that would insert this CGGCGG codon.

A Technical Appendix⁹² entitled, “Detailed methods and primer sequences used in a study of genetically diverse filoviruses in Rousettus and Eonycteris spp. bats, China, 2009 and 2015, by Yang, Xinglou & Zhang, Yunzhi & Jiang, Ren-Di & Guo, Hua & Zhang, Wei & Li, Bei & Wang, Ning & Wang, Li & Rumberia, Cecilia & Zhou, Ji-Hua & Li, Shi-Yue & **Daszak, Peter** & Wang, Lin-Fa & **Shi, Zheng-Li.** (2017), from the Wuhan Institute of Virology identifies primer sequences for doing genetic experiments in coronaviruses and identifies CGG containing primers when a R amino acid is being inserted.

⁹⁰ Since the codon usage here was not reported I contacted Professor Nunberg to inquire which arginine codons were used. He replied: “Unfortunately, those files have all been archived and access to the nt sequences would involve considerable digging. If it is useful to you, I typically choose codons that are more frequent in highly expressed human proteins.”

⁹¹ From the Wuhan Institute of Virology; <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7125963/>

⁹² <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5382765/>

Given that there are two codons of six possibilities that are used in codon optimization, CGG and CGC, the finding of a CGG pair would have a likelihood of happening by chance of (2/6) times (2/6) or one in nine.

Subjective Discount Factor: 80% (this has a probability of being wrong one in five times). This is arbitrary. The calculation to make this adjustment in likelihood is shown here:

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.005	0.995
This is the outcome expected 8 of 9 times if this is codon optimization		0.88
Reduced by 80% confidence		$0.88 \times 0.8 = 0.704$
Impact of this evidence		Increases the likelihood of LO by 70.4 divided by 29.6 or 2.378.
Impact of evidence calculation		$0.995 \times 2.378 = 2.37$
Normalize this step of analysis	$0.005 / (2.37 + 0.005) = 0.002$	$2.37 / (0.005 + 2.37) = 0.998$

Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: SARS-CoV-2 Spike Protein is Highly Optimized for ACE2 Binding and Human Cell Infectivity, a Finding that is Inconsistent with Natural Selection but is Consistent with Laboratory Creation

Summary:

- Andersen et al.⁹³ hypothesized that if the CoV-2 interaction with the human ACE2 was apparently “not ideal,” it was evidence that CoV-2 arose by natural selection.
- The alternative hypothesis would be that a finding that CoV-2 was optimized for ACE2 binding and human infection from the initial infection would be evidence of laboratory creation.
- Andersen relied on a paper for the “not ideal” interaction that relied on a computer algorithm rather than laboratory data, was qualitative in nature, sampled only five amino acids or 0.45% of the interaction region, and was over-interpreted.
- The analysis of the Baric et al. paper cited by Andersen as evidence the interaction was not ideal was reexamined, and it was concluded that Andersen had over-interpreted the paper. The paper was a computer simulation study of only 5 of 201 amino acids in the CoV-2-ACE2 interaction region. Only one of the five amino acids discussed was said to be inferior to the equivalent amino acid in SARS-CoV-1; the remainder were either positive or neutral with respect to binding.
- More recently, Baric has clarified his thoughts concerning the CoV-2 ACE2 receptor binding interaction. In a December 31, 2020 *New England Journal of Medicine* paper⁵⁷ he wrote: “Early zoonotic variants in the novel coronavirus SARS-CoV that emerged in 2003 affected the receptor-binding domain (RBD) of the spike protein and thereby enhanced virus docking and entry through the human angiotensin-converting-enzyme 2 (hACE2) receptor. **In contrast, the spike-protein RBD of early SARS-CoV-2 strains was shown to interact efficiently with hACE2 receptors early on.**” [emphasis added.]
- A comprehensive, laboratory-based, and quantitative paper by Starr et al. of all 201 amino acids in the receptor binding region, not just five amino acids, was examined. Fully 99.6% of all of the possible 3819⁹⁴ amino acid substitutions were tested for their effect on CoV-2 binding to ACE2. Only 21 substitutions of the 3819 improved ACE2 binding. Therefore, CoV-2 has been optimized for human ACE2 binding in 99.45% of the possible amino acids in its Spike Protein interaction region.

⁹³ <https://www.nature.com/articles/s41591-020-0820-9>

⁹⁴ There are 201 amino acids in the residue 331 to 531 interaction region and so 201 times the 19 possible alternative amino acids not found in CoV-2 equals 3819.

- To support this finding, Starr also made an examination of 31,570 CoV-2 sequences from human infections, looking for the 21 substitutions that had been shown to improve CoV-2 binding in the above in vitro laboratory experiments. Among the 31, 570 CoV-2 cases, they failed to find even a single case in which there was an amino acid substitution that improved binding at the time of writing this analysis.⁹⁵
- Based on Andersen's hypothesis and its alternative, SARS-CoV-2 is fully optimized for interaction with the human ACE2 receptor and was at the time of the first patient. There is no evidence of an evolving SP binding region, as was seen with SARS-CoV-1. This is consistent with a laboratory optimized coronavirus which entered the human population fully evolved.

Analysis

Quote from Andersen: "While the analyses above suggest that SARS-CoV-2 may bind human ACE2 with high affinity, computational analyses predict that the interaction is not ideal (reference 7) and that the RBD sequence is different from those shown in SARS-CoV to be optimal for receptor binding (references 7,11).

Thus, the high-affinity binding of the SARS-CoV-2 spike protein to human ACE2 is most likely the result of natural selection on a human or human-like ACE2 that permits another optimal binding solution to arise. This is strong evidence that SARS-CoV-2 is not the product of purposeful manipulation."

The apparent **hypothesis** for the above conclusion is:

"If the SARS-CoV-2 (CoV-2) Spike Protein interaction with the ACE2 receptor is not maximized, then it is evidence that the interaction is the product of natural selection and not purposeful (laboratory) manipulation."


This would lead to an **alternative hypothesis**:

"If the CoV-2 Spike Protein interaction with the ACE2 receptor is maximized, then it is evidence that the interaction *was* the product of purposeful (laboratory) manipulation."

Background.

The Spike Protein (SP) structure and its functional domains are shown in this Figure. The S1 subunit is the initial host interaction portion while the S2 is the post-binding portion responsible for initiating host cell entry, with HR1, HR2, and TM being responsible for breaching the host cell membrane. Allowing viral RNA to enter the cell.

⁹⁵ The recent finding of the N501Y variant, first in the UK, and now spreading globally, is evidence of the power of this analysis. N501Y is one of only five potential substitutions in the Starr analysis that had a major effect in improving ACE2 binding.



The interaction of the SP portions which interact with the ACE2 of the host cell, which begins the internalization, infectious process, are contained in the Receptor Binding Domain (RBD) and to a lesser extent the Receptor Binding Motif (RBM), specifically residues 331 to 531. Herein, residues 331 to 531 are called the “interaction region.”

Evidence given by Andersen:

Reference 7 in the Andersen paper above is a Ralph Baric paper⁹⁶ from early in the pandemic (submitted January 22, 2020) and examines five key residues in the receptor binding domain of the Spike Protein (SP) and whether they are “ideal” for interacting with the ACE2 of human cells. The entire paper is based on computer calculations or prior laboratory work but importantly does not do any new “wet” lab work with CoV-2.

Baric et al. had previously identified five amino acid residues that are important for SP-ACE2 interaction. Using the amino acid numbers of CoV-2, these amino acids are: 455, 486, 493, 494, and 501. Baric opines that the most critical residues are 493 and 501 and the next most important residues are 455, 486, and 494. The authors then discuss each amino acid in turn:

Residue 493: “Gln493 in 2019-nCoV RBD is compatible with hot spot 31, suggesting that 2019-nCoV is capable of recognizing human ACE2 and infecting human cells.” In this analysis, 4 of the 20 amino acids are probed.

Residue 501: “This analysis suggests that 2019-nCoV recognizes human ACE2 less efficiently than human SARS-CoV (year 2002) but more efficiently than human SARS-CoV (year 2003). Hence, at least when considering the ACE2-RBD interactions, 2019-nCoV has gained some capability to transmit from human to human.”

Direct binding evidence has shown that this statement is misleading, and CoV-2 binds the ACE2 receptor about ten-times better than SARS-CoV (year 2002).⁹⁷ In this analysis 3 of the 20 amino acids are probed.

Residues 455, 486, and 494: First, Baric et al. state: “Leu455 of 2019-nCoV RBD provides favorable interactions with hot spot 31, hence enhancing viral binding to human ACE2.”

Next, they state: “Phe486 of 2019-nCoV RBD provides even more support for hot spot 31, hence also enhancing viral binding to human ACE2.” Importantly, they also talk about their own laboratory work on an “optimized” receptor binding domain and state: “Leu472 of human and

⁹⁶ <https://jvi.asm.org/content/94/7/e00127-20>

⁹⁷ <https://www.cell.com/action/showPdf?pii=S0092-8674%2820%2931003-5> ;
<https://www.nature.com/articles/s41586-020-2179-y> ;
<https://www.sciencedirect.com/science/article/pii/S0092867420302622> ;
<https://science.sciencemag.org/content/367/6483/1260>

civet SARS-CoV RBDs provides favorable support for hot spot 31 on human ACE2 through hydrophobic interactions with ACE2 residue Met82 and several other hydrophobic residues (**this residue has been mutated to Phe472 in the optimized RBD**).” [emphasis added.]

Finally, they state: Ser494 in 2019-nCoV RBD still provides positive support for hot spot 353, but the support is not as favorable as that provided by Asp480. Overall, Leu455, Phe486, and Ser494 of 2019-nCoV RBD support the idea that 2019-nCoV recognizes human ACE2 and infects human cells.”

In this analysis they probe 3 of 20 amino acid residues for position 480, 4 of 20 for position 486, and 4 of 20 for position 442.

As shown in the Figure below from the Baric paper, the in vitro designed, optimized human SP (red arrow) had the amino acid residues F, F, N, D, and T at these five key residues. Since CoV-2 was identical in only one of these five it was not “optimal” and, according to Andersen, it therefore was not laboratory derived.

B

Virus	Year	442	472	479	480	487
SARS - human	2002	Y	L	N	D	T
SARS - civet	2002	Y	L	K	D	S
SARS - human/civet	2003	Y	P	N	G	S
SARS - civet	2005	Y	P	R	G	S
SARS - human	2008	F	F	N	D	S
Viral adaption to human ACE2		F > Y	F > L > P	N = R >>> K	D > G	T >>> S
Optimized - human	In vitro design	F	F	N	D	T
Viral adaption to civet ACE2		Y > F	P = L > F	R > K = N	G > D	T > S
Optimized - civet	In vitro design	Y	P	R	G	T
SARS - bat	2013	S	F	N	D	N
2019-nCoV – human	2019	L (455)	F (486)	Q (493)	S (494)	N (501)

Conclusion from the above paper: by examining five amino acid residues of the 200 residues encompassing the interaction region, and calculating the expected interaction of a total of 18 of the 4000 possible residues or 0.45% of all possibilities, they conclude CoV-2 can infect human cells, but is not optimized to do so. This data was twisted by Andersen to show ‘strong evidence’ of natural selection.

An alternative and comprehensive analysis in another paper:⁹⁸


The receptor binding domain (RBD) of the CoV-2 SP is included in residues 331 to 531, a 201 amino acid sequence, of the SP. To examine the effect of each and every amino acid in each and every position, all 19 different amino acids were changed into all 201 positions of the RBD to the extent possible. Out of a total potential of 3819 different single amino acid variants, the scientists

⁹⁸ <https://www.cell.com/action/showPdf?pii=S0092-8674%2820%2931003-5>

were able to create 3804 of the potential variants or 99.6% of the possible variants. It is probable that the variants with the 0.4% amino acid substitutions could not be made for one reason or another. These 3804 were then tested for binding to the human ACE2. Finally, the RBD from SARS-CoV-1 also was tested.

The Figure below is the result of the experiment. Starting with amino acid 331 and ending with amino acid 531, the amino acids that were changed are in vertical columns and are color coded. Shades of brown are amino acid substitutions that reduce ACE2 binding affinity and blue are amino acid substitutions that improve binding, in all cases compared to the 'native' CoV-2 SP sequence. White is the color of a neutral substitution which neither enhances nor diminishes binding. Only the dark blue substitutions provide a strong improvement in ACE2 binding. There is a black square along the top row that denotes amino acids in the SP that interact with the ACE2 protein. Unlike in the Baric analysis above, in which only five amino acids were considered, this group of 19 amino acids provide a more complete interaction picture.

The first overarching observation is that most amino acid substitutions among the 201 amino acids are negative; while a large number are neutral. The fact that the vast majority of amino acid substitutions do not provide an improved ACE2 interaction is clear evidence that the CoV-2 SP interaction region is not newly evolved to the human ACE2 but arrived in the first patient having been “trained” to invade and kill human cells.




There are three levels of improved binding as designated by dark blue, medium blue, and pale blue. Out of the 3804 variants tested, there are 4 dark blue substitutions or 0.11% and 17 medium blue or 0.45%. According to the paper, the binding effect of the light blue could not be measured as different from the native sequence.

The conclusion of this comprehensive work is the demonstration that for 99.45% of the amino acids in the 201 amino acid interaction region, the CoV-2 choice is optimized, where any substitution is either detrimental or, at best, neutral with respect to the first step of CoV-2 entry to human cells, the binding step to the ACE2 receptor.

How much could CoV-2 binding be improved or made worse by substitutions during the human-to-human transmission of the pandemic?

The Figure 4 below, taken from the paper, shows that the three best amino acid substitutions have only a slight effect on the binding curve (Black is wildtype; curves to the left are better binding; curves to the right are worse binding). This is further evidence that CoV-2 is an optimized form of the original virus.



The authors also concluded that Anderson et al. was wrong: “An initially surprising feature of SARS-CoV-2 was that its RBD tightly binds ACE2 despite differing in sequence from SARS-CoV-1 at many residues that had been defined as important for ACE2 binding by that virus (Andersen et al., 2020; Wan et al., 2020).”

In fact, multiple studies have shown that CoV-2 binds ACE2 better than SARS-CoV-1, contradicting Andersen.


Is there evidence that CoV-2 in human circulation has mutations that enhance ACE2 binding?

Another measure of whether CoV-2 is optimized for human infection is to see if Spike Protein mutations have arisen during the pandemic that improve binding of the virus to the ACE2 receptor or if the SP amino acids are ideal from the very first human patient.

The Starr paper addressed this issue as well. A total of 31,570 human sequences were analyzed to see if any of the 21 amino acid substitutions from the binding experiments (or any other for that matter) were being selected for. That is, if there is any evidence of evolutionary pressure to improve SARS-CoV-2 infectivity.

Below is Figure 8 of the Starr paper. Of the 31,570 sequences, all mutations in the receptor interaction region were analyzed for their effect on ACE2 binding. The data below are for all examples of a single nt mutation (1192), two mutations (98), 3-5 mutations (42), and six or more (13) and the effect the mutation would have on ACE2 binding. The logarithmic scale has the

wildtype CoV-2 as 0 and each negative integer is a 10-fold reduction in affinity. Shockingly, there is not a single mutation that is above the 0 line, which would be an improved affinity for the ACE2 receptor. All of the mutations lower the receptor affinity.



Here are the results, in the words of Starr:

“Our discovery of multiple strong affinity-enhancing mutations to the SARS-CoV-2 RBD raises the question of whether positive selection will favor such mutations, since the relationship between receptor affinity and fitness can be complex for viruses that are well-adapted to their hosts (Callaway et al., 2018; Hensley et al., 2009; Lang et al., 2020). Strong affinity-enhancing mutations are accessible via single-nucleotide mutation from SARS-CoV-2 (Figure S8C), but **none are observed among circulating viral sequences in GISAID** (Figure 8A), and **there is no significant trend for actual observed mutations to enhance ACE2 affinity more than randomly drawn samples of all single nucleotide mutations** (see permutation tests in Figure S8D). **Taken together, we see no clear evidence of selection for stronger ACE2 binding, consistent with SARS-CoV-2 already possessing adequate ACE2 affinity at the beginning of the pandemic.**” [emphasis added.]

It is striking that the authors, in observing the complete absence of any evidence for stronger ACE2 binding in over thirty thousand cases, would describe this as evidence of “adequate ACE2 affinity” and not as an exceptional finding of “optimized ACE2 affinity.” Of course, calling the SP affinity exceptional from the beginning of the pandemic would beg the question of a laboratory derived virus.

Returning to the initial hypotheses, since the 3804 possible amino acids at the receptor interaction region of CoV-2 are 99.45% optimized for ACE2 binding, and there is not a single

example in 31,570 human CoV-2 genomes of a substitution that enhances ACE2 binding, the CoV-2 interaction with ACE-2 was maximized from the get-go.

Therefore, the hypothesis, “If the SARS-CoV-2 (CoV-2) Spike Protein interaction with the ACE2 receptor is not maximized, then it is evidence that the interaction is the product of natural selection and not purposeful (laboratory) manipulation,” is **rejected**.

The alternative hypothesis, “If the CoV-2 Spike Protein interaction with the ACE2 receptor is maximized, then it is evidence that the interaction was the product of purposeful (laboratory) manipulation,” is thus **accepted**.

At the time of this writing, a new RBD mutant N501Y has been observed. It is one of the five potential mutations that could be expected to increase RBD-ACE2 affinity.

This is the first example of evidence that will not be statistically quantified but treated as a 51%.49% preponderance of the evidence adjustment. The evidence is more consistent with having been optimized by various methods used in the laboratory than with the slow natural process as seen with SARS-CoV-1, and so the conservative rule that this is consistent with a laboratory origin (51%) versus zoonotic origin (49%) will be used. There will be no confidence adjustment.

The adjusted likelihoods are shown in the following table.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.002	0.998
This is the outcome favors LO over ZO at 51% versus 49%		0.51
Impact of this evidence		Increases the likelihood of LO by $51/49 = 1.041$
Impact of evidence calculation		$1.041 \times 0.998 = 1.039$
Normalize this step of analysis	$0.002/(0.002 + 1.039) = 0.002$	$1.039/(0.002 + 1.039) = 0.998$


Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence. Whole genome comparison of human adaption of CoV-2 compared to SARS-CoV-1 is consistent with a “pre-adaption” of CoV-2 to the human host

A paper⁹⁹ entitled, “SARS-CoV-2 is well adapted for humans. What does this mean for re-emergence?” by Shing Hei Zhan, Benjamin E. Deverman, and Yujia Alina Chan states in the abstract:

“In a side-by-side comparison of evolutionary dynamics between the 2019/2020 SARS-CoV-2 and the 2003 SARS-CoV, we were surprised to find that SARS-CoV-2 resembles SARS-CoV in the late phase of the 2003 epidemic, after SARS-CoV had developed several advantageous adaptations for human transmission. Our observations suggest that **by the time SARS-CoV-2 was first detected in late 2019, it was already pre-adapted to human transmission to an extent similar to late epidemic SARS-CoV. However, no precursors or branches of evolution stemming from a less human-adapted SARS-CoV-2-like virus have been detected.** The sudden appearance of a highly infectious SARS-CoV-2 presents a major cause for concern that should motivate stronger international efforts to identify the source and prevent re-emergence in the near future. [Emphasis added.]

The following Figure from the paper best illustrates the relative SNV adaption for SARS-CoV-1 versus CoV-2.



The paper also makes a tangential comment about posterior diversity: “It would be curious if no precursors or branches of SARS-CoV-2 evolution are discovered in humans or animals.”

This is another example of evidence that will not be statistically quantified. The evidence is more consistent with having been adapted by various known methods used in a laboratory than with the slow natural process as seen with SARS-CoV-1, and so the conservative rule that this is consistent with a laboratory origin (51%) versus zoonotic origin (49%) will be used. There will be no confidence adjustment.

⁹⁹ <https://www.biorxiv.org/content/10.1101/2020.05.01.073262v1>

The adjusted likelihoods are shown in the following table.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.002	0.998
This is the outcome favors LO over ZO at 51% versus 49%		0.51
Impact of this evidence		Increases the likelihood of LO by $51/49 = 1.041$
Impact of evidence calculation		$1.041 \times 0.998 = 1.039$
Normalize this step of analysis	$0.002 / (0.002 + 1.039) = 0.002$	$1.039 / (0.002 + 1.039) = 0.998$


Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: Evidence of CoV-2 during early 2019 in wastewater from Barcelona, Spain is a false positive artifact

A paper entitled “Sentinel surveillance of SARS-CoV-2 in wastewater anticipates the occurrence of COVID-19 cases”¹⁰⁰ claims CoV-2 was present in Barcelona, Spain in March 2019. Specifically, they state:

“This possibility prompted us to analyze some archival WWTP samples from January 2018 to December 2019 (Figure 2). All samples came out to be negative for the presence of SARS-CoV-2 genomes with the exception of March 12, 2019, in which both IP2 and IP4 target assays were positive. This striking finding indicates circulation of the virus in Barcelona long before the report of any COVID-19 case worldwide.”

This is a false positive



As shown above from the paper, they found 43/45 runs with zero and two runs had only 600-800 CoV-2 copies/L

But the limit of detection (LoD) of their assay is 1,000,000 CoV-2/L.

According to the Promega PCR assay FDA clearance package, the Ct at the LoD is 33-34 for the N1 and N2, respectively (Table 17, page 51).¹⁰¹ Here the LoD is listed as 1 RNA/μL.

In the paper the Ct is 40 or 6-7 above the LoD.

This evidence is neutral as to origin and will not be used to adjust the likelihoods. It does reduce the credibility of some of the new origin theories coming out of China.

¹⁰⁰ <https://www.medrxiv.org/content/10.1101/2020.06.13.20129627v1.full.pdf>

¹⁰¹ https://twitter.com/quay_dr/status/1340572543548227585/photo/1

Evidence: WHO and Dr. Shi have spoken of the singular nature of the beginning of COVID-19

On January 23, 2020 Dr. Shi wrote in the draft of her paper: “The almost identical sequences of this virus in different patients imply a probably recent introduction in humans...”¹⁰² By February 3, 2020, when the final version of this paper was published, this sentence had been **deleted**.¹⁰³

On April 23, 2020 the WHO stated: “All the published genetic sequences of SARS-CoV-2 isolated from human cases are very similar. This suggests that the start of the outbreak resulted from a single point introduction in the human population around the time that the virus was first reported in humans in Wuhan, China in December 2019.”¹⁰⁴

The evidence, like the lack of posterior diversity and seroconversion reported earlier, is more consistent with a single introduction in a laboratory accident. This evidence will not be used to adjust probabilities but is included because it could be a form of party admissions of unfavorable facts.

¹⁰² [RaTG13 paper as a preprint](#)

¹⁰³ [RaTG13 final Nature paper](#)

¹⁰⁴ [WHO document page 2 of 12](#)

Evidence. As documented by Drs. Daszak, Humes, and Shi, mammalian biodiversity and bat species differences between Yunnan and Hubei Province are significant and do not support a zoonotic origin

Summary. SARS-CoV-2 is most closely related to bat coronaviruses from Yunnan, a rural province in South West China. Wuhan, where the pandemic began, is a large urban city of 11 million inhabitants in north central China. These two areas are approximately 1900 km apart.

This is the US equivalent of the difference between New York City (population 8.4 million) and the Everglades in Florida, 2000 km away. The incongruent image of a bat or intermediate host in the Everglades somehow finding its way to New York City is a clear demonstration of the difficulty in this hypothetical transmission process. Nonetheless, a strict literature-based analysis will be conducted.

If COVID-19 is a zoonotic disease it must have travelled from bats to humans or from bats to an intermediate species to humans. Therefore, an examination of mammalian biodiversity differences and commonalities between Yunnan and Wuhan might provide useful information about the intermediate host or the particular bat species.

Peter Daszak, Zhengli-li Shi and colleagues published an August 2020 paper entitled, “Origin and cross-species transmission of bat coronaviruses in China,”¹⁰⁵ in which they make a number of observations that are relevant to this analysis. It should be remembered that both lead authors have made multiple, strong, public statements over many months where they assert that SARS-CoV-2 is a natural virus of zoonotic origin.

Yunnan and Hubei Provinces have very dissimilar mammalian diversity

Quoting from the Methods section of the Daszak, Shi paper:

“Defining zoogeographic regions in China:


Hierarchical clustering was used to define zoogeographic regions within China by clustering provinces with similar mammalian diversity. Hierarchical cluster analysis classifies several objects into small groups based on similarities between them. To do this, we created a presence/absence matrix of all extant terrestrial mammals present in China using data from the IUCN spatial database and generated a cluster dendrogram using the function *hclust* with average method of the R package stats. Hong Kong and Macau were included within the neighboring Guangdong province. We then visually identified geographically contiguous clusters of provinces for which CoV sequences are available (Fig. 1 and Supplementary Fig. 1).

We identified six zoogeographic regions within China based on the similarity of the mammal community in these provinces: **SW (Yunnan province)**, **NO** (Xizang, Gansu, Jilin, Anhui, Henan, Shandong, Shaanxi, Hebei, and Shanxi provinces and Beijing municipality), **CN (Sichuan and Hubei provinces)**, **CE** (Guangxi, Guizhou, Hunan, Jiangxi, and Zhejiang provinces), **SO** (Guangdong and Fujian provinces, Hong Kong, Macau, and Taiwan), and **HI**.

¹⁰⁵ <https://www.nature.com/articles/s41467-020-17687-3#Sec19>

Hunan and Jiangxi, clustering with the SO provinces in our dendrogram, were included within the central region to create a geographically contiguous Central cluster (Supplementary Fig. 1). These six zoogeographic regions are very similar to the biogeographic regions traditionally recognized in China. The three β -CoV sequences from HI were included in the SO region to avoid creating a cluster with a very small number of sequences.”

Below is a cluster dendrogram of Chinese provinces based on similarities between their mammalian diversity (hierarchical clustering). Provinces with CoV sequences available in this study are highlighted in bold.



The y-axis height is a measure of the biodiversity with 1.0 being complete similarity and 0.0 being no similarity. As expected for the geography and location of the two provinces, Yunnan (red arrow above) and Hubei (green arrow above) have a height score of about 0.1, with seven branches and six nodes separating them. This is close to the biggest different in mammalian biodiversity of any two locations in all of China.

In conclusion, Daszak and Shi et al. demonstrate that the mammalian biodiversity between Yunnan and Hubei is very significant, reducing the options for a common intermediate host to be the natural conduit between bats and humans.

Shi, Humes, and Daszak statement: “SARS-CoV-2 is likely derived from a clade of viruses originating in horseshoe bats (*Rhinolophus* spp.). The geographic location of this origin appears to be Yunnan province.”

This evidence will not be statistically quantified. The evidence reduces the biodiversity overlap needed to create a common intermediate species between the two provinces, and so the conservative rule that this is consistent with a laboratory origin (51%) versus zoonotic origin (49%) will be used. There will be no subjective discount factor adjustment.


Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.002	0.998
This is the outcome favors LO over ZO at 51% versus 49%		0.51
Impact of this evidence		Increases the likelihood of LO by $51/49 = 1.041$
Impact of evidence calculation		$1.041 \times 0.998 = 1.039$
Normalize this step of analysis	$0.002/(0.002 + 1.039) = 0.002$	$1.039/(0.002 + 1.039) = 0.998$

Because of the rule on the use of significant figures, the likelihood does not change.

Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: The ancestor of SARS-CoV-2 can hypothetically only obtain a furin site by recombination outside of the sarbecovirus subgenera but there is strong evidence that coronavirus recombination is largely limited to the clade level, with limited evidence of subgenera or genera recombination

- SARS-CoV-2 is a beta coronavirus, subgenera sarbecovirus and is the only sarbecovirus with a furin site.¹⁰⁶
- Furin sites can be found in either alpha or gamma coronaviruses or the other beta coronavirus subgenera. The following Figure from reference 66 shows examples of such coronaviruses (furin containing viruses are shown in red):




- To acquire a furin site in nature would require a co-infection between the CoV-2 sarbecovirus ancestor and a furin-containing non-sarbecovirus as shown above.
- However, there is no evidence of recombination in coronaviruses at either the genus level or the subgenus level; only at the clade level.¹⁰⁷¹⁰⁸
- There is also evidence from Daszak and Shi that within the subgenera of the beta coronaviruses, there is bat host specificity. So, each subgenera of coronaviruses has a preferred bat host species. This reduces the opportunities for a co-host event to permit recombination.¹⁰⁹ The phylogeny below shows the problem of host incompatibility for beta coronaviruses:

¹⁰⁶ <https://www.sciencedirect.com/science/article/pii/S1873506120304165#f0015>


¹⁰⁷ <file:///C:/Users/Steven%20Quay/Desktop/journal.pgen.1009272.pdf>

¹⁰⁸ <https://academic.oup.com/mbe/advance-article/doi/10.1093/molbev/msaa281/5955840>


¹⁰⁹ <https://www.nature.com/articles/s41467-020-17687-3#Sec2>




- Daszak and Shi also identified preferred directions of host switching. Since RaTG13, the closest coronavirus to SARS-CoV-2, is most closely related to viruses with bat hosts from the family, Rhinolophidae, it would be reasonable to expect furin-containing viruses from other bat hosts to migrate into Rhinolophidae, recombine by methods which have not been identified, and then the furin-containing sarbecovirus could evolve into the ancestor of SARS-CoV-2. Unexpectedly, Daszak et al. found host migration for the Rhinolophidae bats only outward and not inward, as required by the above, admittedly, convoluted process. The data Figure is shown here:



- Daszak and Shi also observed outward host switches from *Rhinolophus* at the genera level as well, also against a hypothesis for furin-site acquisition:



- Finally, this paper by Daszak and Shi states: “We used our Bayesian discrete phylogeographic model with zoogeographic regions as character states to reconstruct the spatiotemporal dynamics of CoV dispersal in China.” If SARS-CoV-2 began in Yunnan and first crossed over into humans in Wuhan, this analysis should support a northerly spatiotemporal dispersal of beta coronaviruses. Unfortunately, Daszak and Shi cannot catch a break; their own data do not support the expected route of dispersion:



As shown in the above Figure the only dispersal routes into Wuhan, which is in the CN region, are from the northern region. And the northern region has no inward dispersals from the SW, southwest region, where Yunnan and the origin of the ancestor of SARS-CoV-2, is located.

- Independent evidence documents that Hubei province does not have the bat species needed for SARS-CoV-2 reservoir host¹¹⁰

While statistical models of this data could be interesting and informative for general research about future spillovers, this is evidence will not be statistically quantified for this analysis. The evidence reduces the opportunities for subgenera co-infection and furin-site recombination into the CoV-2 ancestor and so the conservative rule that this is less consistent with a zoonotic origin (49%) versus laboratory origin (51%) will be used. There will be no subjective discount factor adjustment.

The results from the calculations are shown below.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.002	0.998
This is the outcome favors LO over ZO at 51% versus 49%		0.51
Impact of this evidence		Increases the likelihood of LO by 51/49 = 1.041
Impact of evidence calculation		1.041 x 0.998 = 1.039
Normalize this step of analysis	$0.002 / (0.002 + 1.039) = 0.002$	$1.039 / (0.002 + 1.039) = 0.998$

Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹¹⁰ <file:///C:/Users/Steven%20Quay/Desktop/Zhangetal2009.pdf>

Evidence: Of 410 vertebrate species tested for affinity to CoV-2 Spike Protein binding domain, primate ACE2 receptor, including human and VERO monkey cells, are the best at binding and bat species ACE2 are the worse, making direct bat-to-human host jumping extremely unlikely

- An examination of the ACE2 receptor binding domain amino acid sequences and their suitability for interacting with SARS-CoV-2 was performed in 410 vertebrates, including 252 mammals.¹¹¹
- A five-category binding score was developed based on the conservation properties of 25 amino acids important for the binding between ACE2 and the SARS-CoV-2 spike protein.
- Only mammals fell into the medium to very high categories and only primates scored 25/25 for binding.
- This implies that SARS-CoV-2 is optimized for human ACE2-bearing cells from the first introduction into the human population, an observation that contradicts a zoonotic origin.
- It also suggests that other primates may be the proximate species from which SARS-CoV-2 entered the human population.
- Both VERO monkey kidney cells and ACE2 humanized mice would qualify as an intermediate species by this criterion.
- Surprisingly, “all chiropterans (bats) scored low ($n = 8$) or very low ($n = 29$), including the Chinese rufous horseshoe bat, from which a coronavirus (SARSr-CoV ZC45) related to SARS-CoV-2 was identified.”
- This is evidence that bats are probably not a reservoir host for SARS-CoV-2.
- A separate study observed: “Severe acute respiratory syndrome coronavirus 2 did not replicate efficiently in 13 bat cell lines.”¹¹²
- The following two Tables are taken from the paper and are organized according to ACE2 SARS-CoV-2 affinity, from highest to lowest:

¹¹¹ <https://www.pnas.org/content/117/36/22311>

¹¹² https://wwwnc.cdc.gov/eid/article/26/12/20-2308_article

Steven C. Quay, MD, PhD

29 January 2021

	ID	S19	D24	F28	D30	K31+	E37	D38	C41	L45	N43+	I79	Y83	N89+	N32P	K33P	G344	D355	B393	
VERY HIGH																				
<i>Homo sapiens</i> (Human)	25
<i>Gorilla gorilla gorilla</i> (Western lowland gorilla)	25
<i>Nomascus leucogenys</i> (Northern white-cheeked gibbon)	25
<i>Pongo abelii</i> (Sumatran orangutan)	25
<i>Macaca fascicularis</i> (Crab-eating macaque)	25
<i>Mandrillus leucophaeus</i> (Drill)	25
<i>Nasalis larvatus</i> (Proboscis monkey)	25
<i>Pan paniscus</i> (Bonobo)	25
<i>Pan troglodytes</i> (Chimpanzee)	25
<i>Ptilocolobus tephrosceles</i> (Ugandan red colobus)	25
<i>Pygathrix nemaeus</i> (Red-shanked douc)	25
<i>Rhinopithecus roxellana</i> (Golden snub-nosed monkey)	25
<i>Chlorocebus sabaeus</i> (Green monkey)	25
<i>Erythrocebus patas</i> (Patas monkey)	25
<i>Macaca mulatta</i> (Rhesus macaque)	25
<i>Papio anubis</i> (Olive baboon)	25
<i>Theropithecus gelada</i> (Gelada)	25
<i>Cercocebus atys</i> (Sooty mangabey)	25
<i>Macaca nemestrina</i> (Southern pig-tailed macaque)	25
HIGH																				
<i>Colobus angolensis</i> (Angola colobus)	24	R
<i>Propithecus coquereli</i> (Coquerel's sifaka)	24	T
<i>Cricetomys gambianus</i> (Gambian pouched rat)	22	.	A	.	.	Q	N
<i>Cricetulus griseus</i> (Chinese hamster)	22	Q	N	.	H
<i>Ctenodactylus gundi</i> (Common gundi)	22	Q	N	.	T
<i>Delphinapterus leucas</i> (Beluga whale)	22	Q	I	T
<i>Eulemur flavifrons</i> (Blue-eyed black lemur)	22	.	E	A	T
<i>Indri indri</i> (Indri)	22	N	Q	N
<i>Monodon monoceros</i> (Narwhal)	22	Q	I	T
<i>Neophocaena asiaeorientalis</i> (Narrow-ridged finless porpoise)	22	Q	I	T
<i>Phocoena phocoena</i> (Harbour porpoise)	22	Q	I	T
<i>Balaenoptera acutorostrata scammoni</i> (Minke whale)	21	R	.	I	T
<i>Balaenoptera bonaerensis</i> (Antarctic minke whale)	21	Q	R	.	.	I	T
<i>Eschrichtius robustus</i> (Gray whale)	21	Q	E	.	.	I	T
<i>Nannospalax galili</i> (Spalax)	21	Q	K	.	D	Q	.	.	.
<i>Odocoileus virginianus texanus</i> (White-tailed deer)	21	E	M	T	.	H	.	.	.
<i>Rangifer tarandus</i> (Reindeer)	21	E	M	T	.	H	.	.	.
<i>Tamandua tetradactyle</i> (Southern tamandua)	21	E	Q	I	T
<i>Dipodomys stephensi</i> (Stephens's kangaroo rat)	20	.	L	.	.	N	Q	I	.	K	.	.	.
<i>Elaphurus davidianus</i> (Pere David's deer)	20	E	N	M	T	.	H	.	.	.
<i>Ellobius lutescens</i> (Transcaucasian mole vole)	20	.	D	.	.	.	Q	S	.	H	.	D	.	.
<i>Globicephala melas</i> (Long-finned pilot whale)	20	.	R	.	.	Q	.	R	I	T
<i>Lagenorhynchus obliquidens</i> (Pacific white-sided dolphin)	20	.	R	.	.	Q	.	R	I	T
<i>Lipotes vexillifer</i> (Baiji)	20	.	R	.	.	Q	I	T	F
<i>Myrmecophaga tridactyla</i> (Giant anteater)	20	E	Q	.	.	N	I	T
<i>Ondatra zibethicus</i> (Muskrat)	20	N	.	Q	S	.	H	.	D	.	.
<i>Orcinus orca</i> (Killer whale)	20	.	R	.	.	Q	.	R	I	T
<i>Tursiops truncatus</i> (Common bottlenose dolphin)	20	.	R	.	.	Q	.	R	I	T
MEDIUM																				
<i>Daubentonia madagascariensis</i> (Aye-aye)	23	F	T
<i>Cheirogaleus medius</i> (Fat-tailed dwarf lemur)	22	K	Q	.	.	.
<i>ICTidomys tridecemlineatus</i> (Thirteen-lined ground squirrel)	22	.	L	.	.	.	Q	A
<i>Marmota flaviventris</i> (Yellow-bellied marmot)	22	.	L	.	.	.	Q	A
<i>Marmota marmota marmota</i> (Alpine marmot)	22	.	L	.	.	.	Q	A
<i>Mesocricetus auratus</i> (Golden hamster)	22	Q	N	.	Y	.	.	.
<i>Physeter catodon</i> (Sperm whale)	22	Q	T	T	.	.	.
<i>Spermophilus dauricus</i> (Daurian ground squirrel)	22	.	L	.	.	.	Q	A
<i>Allactaga bullata</i> (Gobi jerboa)	21	.	T	Q	T	.	H	.	.
<i>Ammotragus lervia</i> (Barbary sheep)	21	E	M	T	.	Y	.	.
<i>Antilocapra americana</i> (Pronghorn)	21	E	M	T	.	Y	.	.
<i>Aotus nancymae</i> (Nancy Ma's night monkey)	21	H	E	Q	.
<i>Beatragus hunteri</i> (Hirola)	21	E	.	Q	M	T	.	Y	.	.
<i>Bison bison bison</i> (American bison)	21	E	M	T	.	Y	.	.
<i>Bos indicus</i> (Zebu)	21	E	M	T	.	Y	.	.
<i>Bos mutus</i> (Wild yak)	21	K	M	T	.	Y	.	.

		S19	S24	F28	D30	H34	H34	E37	E37	V41	Q42	L45	L79	M82	Y83	N322	N330	K359	D365	R367
MEDIUM (continued)		ID																		
	<i>Bos taurus</i> (Cattle)	21	.	.	E	M	T	Y
	<i>Bubalus bubalis</i> (Water buffalo)	21	M	T	Y
	<i>Callicebus donacophilus</i> (White-eared tili)	21	H	E	.	.	T	.	.	Q
	<i>Callithrix jacchus</i> (Common marmoset)	21	H	E	.	.	T	.	.	Q
	<i>Capra aegagrus</i> (Wild goat)	21	.	.	E	M	T	Y
	<i>Capra hircus</i> (Goat)	21	.	.	E	M	T	Y
	<i>Cebus capucinus imitator</i> (Panamanian white-faced capuchin)	21	H	E	.	.	T	.	.	Q
	<i>Felis catus</i> (Cat)	21	.	L	.	E	.	.	E	T
	<i>Giraffa tippelskirchi</i> (Masai giraffe)	21	.	.	E	M	T	Y
	<i>Hemitragus hylociurus</i> (Nilgiri tahr)	21	M	T	Y
	<i>Lynx canadensis</i> (Canadian lynx)	21	.	L	.	E	.	.	E
	<i>Mirza coquereli</i> (Coquerel's giant mouse lemur)	21	.	I	.	.	N	T	.	K
	<i>Moschus moschiferus</i> (Siberian musk deer)	21	.	.	E	M	T	Y
	<i>Neofelis diardi</i> (Sunda clouded leopard)	21	.	L	.	E	.	.	E
	<i>Neofelis nebulosa</i> (Clouded leopard)	21	.	L	.	E	.	.	E
	<i>Okapia johnstoni</i> (Okapi)	21	.	.	E	M	T	Y
	<i>Ovis aries</i> (Sheep)	21	.	.	E	M	T	Y
	<i>Panthera onca</i> (Jaguar)	21	.	L	.	E	.	.	E	T
	<i>Panthera pardus</i> (Leopard)	21	.	L	.	E	.	.	E	T
	<i>Panthera tigris altaica</i> (Siberian tiger)	21	.	L	.	E	.	.	E	T
	<i>Pantholops hodgsonii</i> (Tibetan antelope)	21	.	.	E	M	T	Y
	<i>Perognathus longimembris</i> (Little pocket mouse)	21	.	V	.	.	Q	T	.	K	.	.	.
	<i>Peromyscus maniculatus bairdi</i> (Deer mouse)	21	.	I	.	.	Q	N	.	H	.	.	.
	<i>Pithecia pithecia</i> (White-faced saki)	21	H	E	.	.	T	.	.	Q	.	.	.
	<i>Puma concolor</i> (Cougar)	21	.	L	.	E	.	.	E	T
	<i>Saimiri boliviensis boliviensis</i> (Black-capped squirrel monkey)	21	H	E	.	.	T	.	.	Q	.	.	.
	<i>Sapajus apella</i> (Tufted capuchin)	21	H	E	.	.	T	.	.	Q	.	.	.
	<i>Urocyon parvulus</i> (Arctic ground squirrel)	21	.	L	.	.	Q	.	.	H	.	.	.	D
	<i>Bos indicus</i> x <i>Bos taurus</i>	21	.	.	E	M	T	Y
	<i>Acinonyx jubatus</i> (Cheetah)	20	.	L	.	E	.	.	E	T	.	K	.	.	.
	<i>Alouatta palliata</i> (Mantled howler)	20	.	.	E	H	E	.	.	T	.	.	Q	.	.	.
	<i>Ateles geoffroyi</i> (Geoffroy's spider monkey)	20	A	H	E	.	.	T	.	.	Q	.	.	.
	<i>Fukomys damarensis</i> (Damaraland mole-rat)	20	L	.	.	.	Q	A	.	H	.	N	.	.
	<i>Heterocephalus glaber</i> (Naked mole-rat)	20	L	A	.	H	.	D	.	.
	<i>Hippopotamus amphibius</i> (Hippopotamus)	20	.	.	L	.	D	.	.	.	A	.	F	D
	<i>Lepus americanus</i> (Snowshoe hare)	20	.	L	.	E	.	Q	T	.	S
	<i>Nanger dama</i> (Dama gazelle)	20	.	.	E	M	T	F	Y
	<i>Oryctolagus cuniculus</i> (European rabbit)	20	.	.	L	.	E	.	Y	T	.	S
	<i>Oryx dammah</i> (Scimitar oryx)	20	.	.	E	M	T	Y
	<i>Saguinus imperator</i> (Emperor tamarin)	20	.	A	H	E	.	.	.	T	.	.	Q	.	.	.
	<i>Vicugna pacos</i> (Alpaca)	20	.	L	.	K	E	A	.	I
LOW																				
	<i>Ceratotherium simum cottoni</i> (Northern white rhinoceros)	21	.	L	.	E	.	P	T
	<i>Ceratotherium simum simum</i> (Southern white rhinoceros)	21	.	L	.	E	.	P
	<i>Diceros bicornis</i> (Black rhinoceros)	21	.	L	.	E	.	P
	<i>Galeopterus variegatus</i> (Sunda flying lemur)	21	F	L	E	.	.	N
	<i>Peromyscus leucopus</i> (White-footed mouse)	21	.	I	.	.	Q	N	.	P
	<i>Ailuropoda melanoleuca</i> (Giant panda)	20	.	L	.	E	.	Y	H	T
	<i>Camelus bactrianus</i> (Bactrian camel)	20	.	L	.	E	E	T
	<i>Camelus dromedarius</i> (Dromedary)	20	.	L	.	E	E	T
	<i>Camelus ferus</i> (Wild bactrian camel)	20	.	L	.	E	E	T
	<i>Dicorhinus sumatrensis</i> (Sumatran rhinoceros)	20	.	L	.	E	.	P	.	H	.	.	.	T
	<i>Graphiurus murinus</i> (Woodland dormouse)	20	.	T	.	.	N	R	H	N
	<i>Tapirus indicus</i> (Malayan tapir)	20	.	L	.	E	.	P	.	H	T
	<i>Tapirus terrestris</i> (South American tapir)	20	.	L	.	E	.	P	.	H	T
	<i>Ursus arctos horribilis</i> (Grizzly bear)	20	.	L	.	E	.	Y	H	T
	<i>Ursus maritimus</i> (Polar bear)	20	.	L	.	E	.	Y	H	T
	<i>Canis lupus dingo</i> (Dingo)	19	.	L	.	E	.	Y	.	E	T	.	D	.	.	.
	<i>Canis lupus familiaris</i> (Dog)	19	.	L	.	E	.	Y	.	E
	<i>Chinchilla lanigera</i> (Long-tailed chinchilla)	19	.	.	.	H	E	K	A	.	H	.	D	.
	<i>Chrysocyon brachyurus</i> (Maned wolf)	19	.	L	.	E	.	Y	.	E	T	.	D	.	.	.
	<i>Dipodomys ordii</i> (Ord's kangaroo rat)	19	Q	L	.	.	N	Q	I	.	K
	<i>Eonycteris spelaea</i> (Lesser dawn bat)	19	.	L	.	E	.	.	T	T	.	D	.	K	.
	<i>Equus asinus</i> (Donkey)	19	.	L	.	E	.	S	.	E	H	.	.	.	H
	<i>Equus caballus</i> (Horse)	19	.	L	.	E	.	S	.	E	H	.	.	.	H

Bayesian Analysis of SARS-CoV-2 Origin

Steven C. Quay, MD, PhD

29 January 2021

	D	S19	Q24	Q27	F28	K31*	H34	E39*	D38	Y41	L45	N53*	L79	Y83	N89*	N93*	K353*	G354	R357	R358
LOW (continued)																				
<i>Equus przewalskii</i> (Przewalski's horse)	19	L	E	S	E	H														
<i>Hydrocoerus hydrochaeris</i> (Capybara)	19			E	L	K														
<i>Hystrix cristata</i> (Crested porcupine)	19	I			Q								F	A	H	N				
<i>Megaderma lyra</i> (Indian false vampire)	19			E	L	E			H	F										
<i>Microtus ochrogaster</i> (Prairie vole)	19	D	A		Q								S	H	D					
<i>Rhinolophus pearsonii</i> (Pearson's horseshoe bat)	19	I			R		H	E												
<i>Rhinolophus sinicus</i> (Chinese rufous horseshoe bat)	19	F	R		E	F			N											
<i>Rousettus aegyptiacus</i> (Egyptian rousette)	19	L	E	T									T	D	K					
<i>Speothos venaticus</i> (Bush dog)	19	L	E	Y		E														
<i>Sus scrofa</i> (Pig)	19	L	E	L									I	T	T					
<i>Tragulus javanicus</i> (Java mouse-deer)	19	I	E	L						M	T	H								
<i>Vulpes lagopus</i> (Arctic fox)	19	L	E	Y		E							T	D						
<i>Vulpes vulpes</i> (Red fox)	19	L	E	Y		E							T	D						
<i>Balaena mysticetus</i> (Bowhead whale)	18			Q	E	R			N				T	T	H					
<i>Carlito syrichta</i> (Philippine tarsier)	18	Q			Q		H		I	S		N	S							
<i>Dasypus punctata</i> (Central American agouti)	18	F			E	Q	K					A	P	N						
<i>Dolichotis patagonum</i> (Pantagonian mara)	18	F			E	L	K					A	H	N						
<i>Eidolon helvum</i> (Straw-colored fruit bat)	18	L	E	T						T	D	K		K						
<i>Loxodonta africana</i> (African elephant)	18	L		T	Q					D	F	S	P							
<i>Microtus murinus</i> (Gray mouse lemur)	18	Q		E	N	N		H		T		K								
<i>Ochotona princeps</i> (American pika)	18	L	E	K			N			T	S	D								
<i>Octodon degus</i> (Common degu)	18	F			N	Q	K					A	H	N						
<i>Procavia capensis</i> (Rock hyrax)	18	L		T	Q					S	F	S								
<i>Pteropus alecto</i> (Black flying fox)	18	L	E	T						A	D	K		K						
<i>Pteropus vampyrus</i> (Large flying fox)	18	L	E	T						A	D	K		K						
<i>Trichechus manatus latirostris</i> (West Indian manatee)	18	L		T	Q					N	F	S								
VERY LOW																				
<i>Catagonus wagneri</i> (Chacoan peccary)	20	L	E	L						T	T									
<i>Jaculus jaculus</i> (Lesser Egyptian jerboa)	19	M			Q					Y	T	P	N							
<i>Cavia porcellus</i> (Guinea pig)	18	F			E	L	K					A	P	N						
<i>Cavia tschudii</i> (Montane guinea pig)	18	F			E	L	K					A	P	N						
<i>Hipposideros armiger</i> (Great roundleaf bat)	18	L	E		T		H	L		R	D									
<i>Hipposideros pratti</i> (Pratt's roundleaf bat)	18	L	E		T		H	L		R	D									
<i>Mesopodisma bidens</i> (Sowerby's beaked whale)	18	P	K	I	Q							T	S							
<i>Spilogale gracilis</i> (Western spotted skunk)	18	L	I	E	Y		E			E	T									
<i>Zapus hudsonius</i> (Meadow jumping mouse)	18	V	D		I	Q				R	T	P								
<i>Ctenomys sociabilis</i> (Social tuco-tuco)	17	F	I		N	Q	K				A	H	N							
<i>Cynopterus brachyotis</i> (Lesser short-nosed fruit bat)	17	L	E	T						T	H	D	K	H						
<i>Cynopterus sphinx</i> (Greater short-nosed fruit bat)	17	L	E	T						T	H	D	K	H						
<i>Enhydra lutris kenyoni</i> (Sea otter)	17	P	E	Y		E			H	T	D	R								
<i>Eumetopias jubatus</i> (Steller sea lion)	17	L	E	S		E			Q	T	D	H								
<i>Grammomys surdaster</i> (African woodland thicklet rat)	17	E			Q					T	N	F	T	Y	H					
<i>Gulo gulo</i> (Wolverine)	17	L	E			E				Q	T	D	H							
<i>Heterohyrax brucei</i> (Yellow-spotted rock hyrax)	17	L		T	Q		E			S	F	S								
<i>Macroglossus sobrinus</i> (Long-tongued fruit bat)	17	L	E	T			E			N	D	K		K						
<i>Manis javanica</i> (Sunda pangolin)	17	E		E	S		E			I	N	K	H							
<i>Manis pentadactyla</i> (Chinese pangolin)	17	E		E	S		E			I	N	K	H							
<i>Mellivora capensis</i> (Honey badger)	17	L	E	Y		E				Q	T	D	R							
<i>Mus pahari</i> (Gaidner's shrewmouse)	17	N		N	Q					T	N	F	H	H						
<i>Mustela erminea</i> (Stoat)	17	L	E	Y		E				H	T	D	R							
<i>Mustela lutreola</i> (European mink)	17	L	E	Y		E				H	T	D	R							
<i>Mustela nigripes</i> (Black-footed ferret)	17	L	E	Y		E				H	T	D	R							
<i>Mustela putorius furo</i> (Ferret)	17	L	E	Y		E				H	T	D	R							
<i>Neomonachus schauinslandi</i> (Hawaiian monk seal)	17	L	E	Y		E				Q	T	D	H							
<i>Petromus typicus</i> (Dassie rat)	17	L		T	Q					A	H	D								
<i>Phoca vitulina</i> (Harbor seal)	17	L	E	Y		E				Q	T	D	R							
<i>Pteronura brasiliensis</i> (Giant otter)	17	L	E	Y		E				H	T	D	R							
<i>Rhinolophus ferrumequinum</i> (Greater horseshoe bat)	17	L	K		D	S		N	H		N	F								
<i>Taxidea taxus</i> (American badger)	17	L	E	Y		E				H	T	D	H							
<i>Thryonomys swinderianus</i> (Greater cane rat)	17	L	L		T	Q				A	R	D								
<i>Zalophus californianus</i> (California sea lion)	17	L	E	S		E				Q	T	D	H							
<i>Acomys cahirinus</i> (Cairo spiny mouse)	16	L	E		S	Q	K			S	N	F	H							
<i>Anoura caudifer</i> (Tailed tailless bat)	16	T	E		E	N	T			A	D									

	ID	S19	Q24	F28	K31*	H34	E39*	D38	Y41	L45	N53*	L79	Y83	N89*	N93*	K353*	G354	R357	R358
VERY LOW (continued)																			
<i>Artibeus jamaicensis</i> (Jamaican fruit-eating bat)	16	A	D		E	T		E		E		A	D		N				
<i>Callorhinus ursinus</i> (Northern fur seal)	16	L		E	S		E		F	Q	T	D		H					
<i>Choloepus hoffmanni</i> (Hoffmann's two-toed sloth)	16	L			T	Q	Q		H		I	T	F	K					
<i>Condylura cristata</i> (Star-nosed mole)	16				E	T	R		E	N		D	R	F	D				
<i>Cryptoprocta ferox</i> (Fossa)	16	L		E	Y		Q	E	L		T	S							K
<i>Dasypus novemcinctus</i> (Nine-banded armadillo)	16				E	T	Q	Q	E	H			M	N	F				
<i>Hipposideros galeritus</i> (Cantor's roundleaf bat)	16		S			I		T	D	E	H			D	D	K			
<i>Hyena hyaena</i> (Striped hyena)	16	L		E	Y		Q	E	L		T	D							K
<i>Miniopterus natalensis</i> (Natal long-fingered bat)	16		K		E	Q	S	Q		F	E								
<i>Miniopterus schreibersii</i> (Schreibers' long-fingered bat)	16		K		I		E	N	S	Q		F	K						
<i>Mirounga angustirostris</i> (Northern elephant seal)	16	L		K	E		Y		E			Q	T	D		H			
<i>Mus caroli</i> (Ryukyu mouse)	16		N		N		N		Q				T	S	F	T	H		
<i>Mus musculus</i> (House mouse)	16		N		N		N		Q					T	S	F	T	H	
<i>Mus spretus</i> (Algerian mouse)	16		N		N		S		Q					T	S	F	T	H	
<i>Myocastor coypus</i> (Coypu)	16	L	A			N	Q	K				F	A		H	N			
<i>Myotis davidii</i> (David's myotis)	16		K		I		N	S	K		H	E		T	S				
<i>Myotis myotis</i> (Greater mouse-eared bat)	16		K		I		N	S	K		H	E		T	S				
<i>Noctilio leporinus</i> (Greater bulldog bat)	16	N		A		E	N	S	K		E			A	D				
<i>Odobenus rosmarus divergens</i> (Walrus)	16	L		E	Y		E		F	Q	T	D		H					
<i>Otlemur gametii</i> (Northern greater galago)	16	Q			N	R		E	H			I	T	E	D				
<i>Paguma larvata</i> (Masked palm civet)	16	L		E	T	Y		Q	E		V	T	D						
<i>Phataginus tricuspis</i> (White-bellied pangolin)	16	A	E		E	S		E			I	N	K		H				
<i>Psammomys obesus</i> (Fat sand rat)	16				E	Q	K					I	N	F	T	H	Q		
<i>Rattus norvegicus</i> (Brown rat)	16		K	S		N		Q				I	N	F	Q	H			
<i>Sarcophilus harrisii</i> (Tasmanian devil)	16	L	M	G		E	N	K	A				A	A					
<i>Ailurus fulgens styani</i> (Red panda)	15				E	T	N	Q	N		E	H	T		H	D			
<i>Carollia perspicillata</i> (Seba's short-tailed bat)	15	T	E		E	T		E	H	E		A	D		N				
<i>Chrysochloris asiatica</i> (Cape golden mole)	15	L	A		N	N	Q		N	H		K	F		D				
<i>Elephantulus edwardii</i> (Cape elephant shrew)	15	P	A		E	Q	Q			V	N	F	D						
<i>Eplesicus fuscus</i> (Big brown bat)	15	N	I		E	N	S		H	E		T	S	N					
<i>Helogale parvula</i> (Common dwarf mongoose)	15	L		E	Q		Q	E		L	V	R	A	S					
<i>Mastomys coucha</i> (Southern multimammate mouse)	15	Q	N		N	Q					I	N	F	T	H				
<i>Meriones unguiculatus</i> (Mongolian gerbil)	15				E	Q	K					I	N	F	T	H	K	Q	
<i>Monodelphis domestica</i> (Gray short-tailed opossum)	15	N	D			D	A	K	E	H		I	T		N				
<i>Mungos mungo</i> (Banded mongoose)	15	L		E	Q		Q	E		L	V	R	A	S					
<i>Murina fese</i> (Little tube-nosed bat)	15	K	A		E	T	S	K		H	E		T	S					
<i>Myotis brandtii</i> (Brandt's bat)	15	K	I		E	N	S	K		H	E		T	S					
<i>Myotis lucifugus</i> (Little brown bat)	15	K	I		E	N	S	K		H	E		T	S					
<i>Orycteropus afer</i> afer (Aardvark)	15	A	L		E	Q		N			I	S	F	P	K				
<i>Paradoxurus hermaphroditus</i> (Asian palm civet)	15	L		E	T	Y		Q	E		V	T	D		D				
<i>Phyllostomus discolor</i> (Pale spear-nosed bat)	15	T	D	K		E	N	N		E		N	D		K				
<i>Scalopus aquaticus</i> (Eastern mole)	15	L		E	N	L	K	N	E			Q	D		N				
<i>Sorex araneus</i> (Common shrew)	15	N	K		N	Q	D					I	T	F	D				
<i>Suricata suricatta</i> (Meerkat)	15	L		E	Q		Q	E		L	V	R	A	S					
<i>Tadarida brasiliensis</i> (Brazilian free-tailed bat)	15	E	I		Q	R	T		E	H		H	R	D					
<i>Tonatia saurophila</i> (Stripe-headed round-eared bat)	15	T			E	N	T	K	E	H		T	D		K				
<i>Microgale talazaci</i> (Talazac's shrew tenrec)	14		Q		E	Q						N	F	D	S	F	N		
<i>Molossus molossus</i> (Velvety free-tailed bat)	14	K	I		N	I	R	E	H		Q	D		N	N				
<i>Mormoops blainvilliei</i> (Antillean ghost-faced bat)	14	I	E		I	N	S		K	H		T	D		N	N			
<i>Neovison vison</i> (American mink)	14	L		E	Y		E			H	T	D		H					
Phascolarctos cinereus (Koala)																			
<i>Pteronotus parnellii</i> (Parnell's mustached bat)	14	F	R	E	E	T	K		E		I	T	F	D					
<i>Solenodon paradoxus</i> (Hispaniolan solenodon)	14	N	K	E		E	L		K	H			E	F		N	N		
<i>Vombatus ursinus</i> (Common wombat)	14	I	E	S	E	S	K	Q		E		K	D		N				
	14	F	R	E	E	T	K		E			I	T	F	D				
<i>Desmodus rotundus</i> (Common vampire bat)	13	T	E		E	N	T		E			I	T	D	S	N	K		
<i>Echinops telfairi</i> (Lesser hedgehog tenrec)	13	S		T	T	N					K	F	D	P	K	L	N		
<i>Erinaceus europaeus</i> (European hedgehog)	13	T	E	K		D	R	Q	N	E		T	N		S	F	N		
<i>Miconycteris hirsuta</i> (Hairy big-eared bat)	13	T	E		E	N	T	K		E	H		K	D		N	K		
<i>Onithorhynchus anatinus</i> (Platypus)	13	K	E	Q		T	Q	K				N	K	F	D		N		
<i>Pipistrellus kuhlii</i> (Kuhl's pipistrelle)	13	E			E	S	N		N	H	E		A	F	D	K	D		
<i>Pipistrellus pipistrellus</i> (Common pipistrelle)	13	E			E	S	N		N	H	E		R	A	F	S	E	D	
<i>Tupaia chinensis</i> (Northern treeshrew)	13	T	E	V		N		I		E	H		K	R	D	D	K	N	
<i>Uropsilus gracilis</i> (Gracile shrew mole)	11	E		E		E	N	R	N	N	K		I	Q	F	D	S	M	K

WIV in GoF research. This will contribute a 51%/49% contribution in favor of laboratory compared to zoonotic origin. There will be no subjective discount factor adjustment.

The results from the calculations are shown below.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.002	0.998
This is the outcome favors LO over ZO at 51% versus 49%		0.51
Impact of this evidence		Increases the likelihood of LO by $51/49 = 1.041$
Impact of evidence calculation		$1.041 \times 0.998 = 1.039$
Normalize this step of analysis	$0.002 / (0.002 + 1.039) = 0.002$	$1.039 / (0.002 + 1.039) = 0.998$

Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: Did a Review of Samples Collected from a Mineshaft Cause the COVID-19 Pandemic?¹¹³

Abstract. The origin of the COVID-19 pandemic caused by SARS-CoV-2 has been hotly debated. Proponents of the natural spillover theory allege that the virus jumped species, possibly via an intermediary host, to cross over to humans via the wildlife trade or by other means. Proponents of a rival theory claim that the virus escaped from a laboratory in Wuhan. This research presents circumstantial evidence of a transmission route via a late 2019 review of samples collected from a mineshaft in Mojiang, Yunnan Province, China. It examines the activity at the Wuhan Institute of Virology in late 2019, when samples from a mineshaft associated with a suspected SARS outbreak were being reviewed. It proposes that spillover occurred during this review of samples including of a virus (BtCoV/4991) only 1% different to SARS-CoV-2 in its RNA-dependent RNA polymerase (RdRp).

It is a meticulous sourced analysis. It purposely avoids the question of whether SARS-CoV-2 was being grown or manipulated in the laboratory, but only addresses the evidence that events in the fall of 2019 are consistent with a laboratory accident.

This will not be used to adjust the likelihoods.

Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹¹³ https://zenodo.org/record/4029545#.X-x_f9gzbOg. Author anonymous. A meticulously documented analysis that concludes an accident occurred at the Wuhan Institute of Virology during the fall of 2019. Includes many primary documents from Mandarin. No direct evidence of 'what' was the nature of the accident or if it was SARS-CoV-2.

Evidence: The Huanan market was not the source of SARS-CoV-2

From the WHO Terms of Reference for the investigation of the origin of SARS-CoV-2:¹¹⁴

“The Huanan wholesale market is a large market (653 stalls and more than 1180 employees) mainly supplying seafood products but also fresh fruits and vegetables, meat, and live animals. In late December 2019, 10 stall operators were trading live wild animals including chipmunks, foxes, racoons, wild boar, giant salamanders, hedgehogs, sika deer, and many others. Farmed, wild and domestic animals were also traded at the market including snakes, frogs, quails, bamboo rats, rabbits, crocodiles, and badgers. The market was closed on 1 January 2020, and several investigations followed, including environmental sampling, as well as sampling of frozen animal carcasses at the market. **Of the 336 samples collected from animals, none were PCR positive for SARS-CoV-2**, whereas 69 out of 842 environmental samples were positive by PCR for SARS-CoV-2. Sixty- one of those (88%) were from the western wing of the market. Of these, 22 samples were from 8 different drains and sewage, and 3 viruses were isolated, sequenced and shared on GISAID. These were virtually identical to the patient samples collected at the same time (>99.9 % homology).”

For contrast, with SARS-CoV-1 91 civets & 15 raccoon dogs in wet markets were tested with 106/106, 100% positive.¹¹⁵

This will not be used to adjust the likelihoods.

Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹¹⁴ <https://drive.google.com/file/d/1rx0W2efbE0R1Ag-lALWTqD22VsWbTIO-/view>

¹¹⁵ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1212604/>

Evidence: Analysis of the hospital of admission for COVID-19 patients during December 2019 places “ground zero” for the outbreak somewhere along Line 2 of the Wuhan Metro System.

Line 2 carries one million people per day and services the Wuhan Institute of Virology, the Hunan Seafood Market, the high-speed rail system, and the Wuhan International Airport

A preprint manuscript¹¹⁶ reported that the earliest genomic cluster of SARS-CoV-2 patients is a group of four individuals associated with the General Hospital of Central Theater Command of People's Liberation Army (PLA) of China in Wuhan. This cluster contains the “Founder Patients” of both Clade A and Clade B, from which every SARS-CoV-2 coronavirus that has infected every patient with COVID-19 anywhere in the world has arisen.

The PLA Hospital is about one mile from the Wuhan Institute of Virology (WIV) and the closest hospital to WIV. Both the PLA Hospital and WIV are serviced by Line 2 of the Wuhan Metro System. The Hunan Seafood Market is also located adjacent to Line 2. All patients between December 1st, 2019 and early January 2020 were first seen at hospitals that also are serviced by Line 2 of the Metro system.

With 40 hospitals located near seven of the nine Metro Lines, the likelihood that all early patients were seen at hospitals only near Line 2 by chance is about 1 in 68,500 (p-value = 0.0000146). The inference then would be that the early spread of SARS-CoV-2 was through human-to human transmission on Line 2.

Line 2 carries one million passengers per day and assuming most are round trip business workers going to and from work in the morning and evening, represents 500,000 riders or about 5% of the Wuhan population. A very recent publication determined that, in fact, 500,000 residents of Wuhan contracted COVID-19, a ten-fold upper estimate.¹¹⁷ The coincidence of my prediction that 500,000 riders on Line 2 were likely exposed to SARS-CoV-2 in late 2019 and the recent admission from Chinese CDC that Wuhan had 500,000 COVID-19 cases is duly noted!

Line 2 connects to all eight other lines of the Wuhan Metro System (1, 3, 4, 6, 7, 8, 11, and Yanglu) facilitating rapid spread in Wuhan and Hubei Province, and also services both the high-speed rail station (Hankou Railway Station), facilitating rapid spread throughout China, and the Wuhan International Airport (Tianhe International Airport), facilitating rapid spread throughout Asia, Europe, and to the United States. In fact, direct human-to-human spread from the Reference Sequence patient to patients around the world is suggested by an unexpectedly reduced genome base substitution rate seen in patient specimens in cities with direct flights from Wuhan.


¹¹⁶ <https://zenodo.org/record/4119263#.X-rszNgzbOg>

¹¹⁷ https://mp.weixin.qq.com/s/LXTfDmsQLf3qZnu_S_MxcA ;
<https://thehill.com/policy/international/china/531935-study-shows-wuhan-coronavirus-cases-may-have-been-10-times-higher>

In a separate paper by Quay and Dr. Martin Lee, Adjunct Professor of Statistics, UCLA, from May 2020, now accepted for publication in *Epidemics*,¹¹⁸ the authors provide evidence that COVID-19 was appearing in California as early as the first week of 2020. This is likely due to direct flights connecting Line 2 to the Wuhan airport and then to San Francisco.

In conclusion, Line 2 of the Wuhan Metro System services the PLA Hospital with the first genomic cluster of patients with COVID-19, the hospitals where patients first went in December 2019 and early January 2020 and is the likely conduit for human-to-human spread throughout Wuhan, China, and the world.

The following slide overview provides a visual analysis of this evidence:



Zoonotic Origin	Laboratory Origin
Hunan Seafood Market	Wuhan Institute of Virology (WIV); Wuhan Center for Disease Control and Prevention (CDC)

How did COVID start? • After 10 months we still don't know!

¹¹⁸ https://www.researchgate.net/publication/341742303_COVID-19_May_Have_Have_Reached_United_States_in_January_2020_05272020




View detail	
Virus name:	NCV-19Wuhan/WHU10219
Accession ID:	EPI_ISL_406790
Type:	Betacoronavirus
Lineage (GISAID):	B.1.1
Clade:	Original
Passage details/history:	
Source and collection date:	2019-12-20
Location:	Asia / China / Hubei / Wuhan
Host:	Human
Additional location information:	
Gender:	male
Patient age:	44
Patient status:	suspected
Specimen source:	Bronchoalveolar Lavage NCCT-C51913
Additional host information:	
Outbreak:	
Last vaccinated:	
Treatment:	DNB08Q
Sequencing technology:	
Assembly method:	SPAdes v.3.12.0
Coverage:	85x
Comment:	
Institute information:	General Hospital of Central Theater Command of People's Liberation Army of China
Originating lab:	NO.627 Wuhan Road, Wuchang District, Wuhan, China
Address:	



GISAID Database

Earliest cases at the PLA Hospital




PLA Hospital is part of the Joint Logistic Support Force Complex

Position in RS	Bat-SL-CoVZC45	Bat-SL-CoVZXC21	RaTG13	PLA-4	PLA-3	PLA-2	Hu-1 Ref seq	PLA-1	GISAID #1
5' UTR	1-5 missing	1-5 missing	1-15 missing	1-16 missing	1-20 missing	1-36 missing	Intact	1-25 missing	Intact
3778	A	A	A	A	A	A	A	A	G
6968	T	T	C	C	C	C	C	A	C
8782	T	T	T	T	C	C	C	C	C
8987	T	T	T	T	T	T	T	T	A
11764	T	T	T	T	T	NA - Note 1	T	A	T
28144	C	C	C	C	T	T	T	T	T
3' UTR	last 4 poly-A missing	last 4 poly-A missing	last 13 poly-A missing	last 15 poly-A missing	last 15 poly-A missing	NA - Note 1	Intact	last 12 poly-A missing	last 4 poly-A missing
Genome length	29802	29732	29855	29872	29868	NA - Note 1	29903	29866	29899
Clade A SNPs Clade B SNPs Non-RaTG13 DNP									
Note 1 - GISAID record: "Long stretches of NNNs (34.45% of overall sequence). Gap of 13 nucleotide(s) found at refpos 26171 (FRAMESHIFT). Gap of 13 nucleotides when compared to the reference sequence. 0.40% Unique Mutations."									

The PLA patient cluster

- PLA-4 is genetically the closest human infection to the three closest bat viruses
- The four PLA patients have the close sequence pattern usually seen only in family transmissions

The Wuhan Metro System



Methodology: Metro station line for intersection of hospital catchment zone and residential living district zone was identified


**All COVID
patients from
Dec 1 to early
Jan were
admitted to
hospitals on
Metro Line 2**

This had the likelihood
of happening by chance
of one in 68,500



Feature	Relationship to Pandemic
Line 2 carried 1 MM passengers a day before COVID	Assuming 2 trips/d for commuters, about 5% of the Wuhan population uses this Line, making it an efficient transmission route for all of Wuhan as well as Hubei Province. A single patient can leave a droplet/aerosol cloud for hours to infect others.
Line 2 shares stations with every other Metro Line	Permits human-to-human spread to every part of Wuhan at the stations shared with Line 2
Line 2, Hankou Railway Station	Connects Wuhan to all of China by high speed rail
Line 2, Tianhe International Airport	International destinations: New York City, San Francisco, London, Tokyo, Rome, Istanbul, Dubai, Paris, Sydney, Bali, Bangkok, Moscow, Osaka, Seoul, and Singapore.

The Line 2 COVID Conduit



The Line 2 COVID Conduit

- Line 2 Hankou High Speed Railway Station to all of China
- Line 2 Tianhe International Airport Station to the world

The Hunan Seafood Market, Wuhan Institute of Virology, and the Wuhan CDC, all locations suggested to be the possible source of SARS-CoV-2 in Wuhan, are also all serviced by Line 2 of the Metro system, suggesting this public transit line should become the focus for further investigations into the origin of this pandemic.


Given that the Hunan Seafood Market has been removed as a source for the origin of CoV-2, this evidence will contribute a 51%/49% contribution in favor of laboratory compared to zoonotic origin. There will be no Subjective Discount Factor adjustment.

The results from the calculations are shown below.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.002	0.998
This is the outcome favors LO over ZO at 51% versus 49%		0.51
Impact of this evidence		Increases the likelihood of LO by 51/49 = 1.041
Impact of evidence calculation		$1.041 \times 0.998 = 1.039$
Normalize this step of analysis	$0.002 / (0.002 + 1.039) = 0.002$	$1.039 / (0.002 + 1.039) = 0.998$

Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: SARS-CoV-2 infection, based on antibody seroconversion, was not found in 39 archived specimens taken from cats (1/3 feral) between March and May 2019¹¹⁹



Based on these results, the prevalence of SARS-CoV-2 in domestic and feral cats prior to January 2020 is less than 8% with a 90% confidence interval.


This will not be used to adjust the likelihoods.

Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹¹⁹ <https://www.tandfonline.com/doi/full/10.1080/22221751.2020.1817796>

Evidence: The extraordinary pre-adaption of SARS-CoV-2 for human cells is demonstrated by a paper looking at a tRNA adaption index.¹²⁰

“The proteome of SARS-CoV-2 is mainly composed of the replicase polyprotein (ORF1ab) and of structural proteins: the spike glycoprotein, the membrane and envelope proteins, and the nucleoprotein [41]. Based on the genomic codon usage of each of the possible host species, we compute the codon adaptation index (CAI) and the tRNA adaptation index (tAI) to estimate the translational efficiency of SARS-CoV-2 proteins in each host (Fig 3A and 3B and S2 Table). Humans are among the top three species whose CAIs are mostly over 0.70, together with ducks and chickens. In terms of the tAI, humans show the highest translational adaptation among all others, followed by chickens, and, to some extent, mice and rats. On the other hand, cats, ferrets, pigs, and dogs are less translationally adapted than humans both by CAI and tAI.”



As shown in panel B above, the tRNA Adaption Index is highest, by far, for humans (blue arrow) followed by the red junglefowl. This is additional evidence of the extraordinary adaption of SARS-CoV-2 to humans from the very beginning. This also is the first evidence of a reasonable intermediate host but based only on these *in silico* data.

This will not be used to adjust the likelihoods.

Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹²⁰ <https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1008450#pcbi.1008450.s004>

Evidence: Evidence of Lax procedures and disregard of laboratory safety protocols and regulations in China, including the Wuhan Institute of Virology

A collection¹²¹ from the Chinese Q&A website, <https://www.zhihu.com/>, of first-hand documentation of laboratory safety breaches and incidents within a large number of laboratories with diverse research subjects and purposes in the People's Republic of China (PRC) is provided. The laboratories involved include Chemistry labs, Biolabs, Computer labs as well as Physics and Engineering labs.

From this first-hand documentation, we obtained evidence of relaxed safety regulations and frequent breaches of such regulations, with reasons ranging from poor training/education on lab safety and chronic ignorance of safety rules, to intentional breaches of protocols for purposes other than the research projects of the lab(s) of which the breach was documented in.

Such breaches often resulted in safety accidents ranging from physical injury, chemical burns, chemical leaks, and damage to property, to lab-acquired infection and escape of in-lab pathogens. With consequences ranging from personal-level to institution-wide impacts.

Here is the reference to the State Department cables concerning safety concerns at the WIV.¹²²

The following document shows that in June 2019, the Chinese CDC was soliciting for the removal of 25-years-worth of solid and liquid medical waste. The total weight is close to two tons including three kg of highly toxic waste.

This is a Google translation of a Mandarin-original website shot from June 27, 2019. The URL highlighted above will lead to the original, which now has been removed from the internet. Having 25 years of toxic waste on site shows a staggering level of disregard for lab safety.

I do not think this is directly linked to CoV-2 origin, but it is a statement about the Chinese CDC. As a reminder, this facility is about 300 meters west of the Seafood market where CoV-2 was first thought to have originated.

¹²¹ <https://zenodo.org/record/4307879#.X-yUo9gzbOh>


¹²² <https://foia.state.gov/Search/Results.aspx?caseNumber=F-2020-05255>

11/22/2020 City Center for Disease Control and Prevention Laboratory Hazardous Chemical Waste Disposal Procurement


<https://www.whcdc.org/index.php/view/11147.html> Go APR 2019

Introduction Party building Business social service scientific Policies and Regulations

homepage guidance research regulations

 **武汉市疾病预防控制中心**
Wuhan Center For Disease Control & Prevention

<http://web.archive.org/web/20200510182006/https://www.whcdc.org/index.php/view/11147.html>



Wuhan Centers For Disease Prevention & Control

NEWS
News topic

our current location: Home

Disease Control News

Municipal Center for Disease Control and Prevention Laboratory Hazardous Chemical Waste Disposal Procurement Project Announcement on Single Source Procurement

Method

Publication unit: Publication time: 2019-06-27 12:27:56 Font size: small , medium and large

The hazardous chemical waste (including solid, liquid, and a small amount of highly toxic drugs) generated in the scientific research process of our center laboratory has not been effectively treated from 1994 to 2019. The total amount of solid and liquid waste of medical waste in the center is The total amount is close to 2 tons, of which nearly 3 kg of highly toxic chemicals are contained, which poses a certain safety hazard to the working environment of the center. In order to eliminate potential safety hazards, it is planned to conduct a one-time disposal of hazardous chemical wastes accumulated in the center.

The center conducted a public bidding for the medical waste treatment project on June 12. According to the "National Hazardous Waste List", the highly toxic substances tested in our laboratory are classified as HW49. Therefore, the corresponding hazardous waste treatment company or unit must have The corresponding qualifications. As of the deadline for registration, only Hubei Zhongyou Youyi Environmental Technology Co., Ltd. has met the qualification response.

Medical waste treatment is closely related to biosafety, environmental safety, public health safety and other aspects, and is a top priority for people's livelihood. In view of the actual situation of the bidding, it is planned to purchase the central medical waste treatment project from a single source, and it is recommended Environmental Protection Technology Co., Ltd. "HW49" qualification is publicized from a single source. The publicity period is 3 working days.

Contact number: 027-85801768.


This is a Google translation of a Mandarin-original website shot from June 27, 2019. The URL highlighted above will lead to the original, which is now removed from the internet. Having 25 years of toxic waste on site shows a level of lab safety disregard that is staggering. I do not think this is directly linked to CoV-2 origin but it is a statement Re the Chinese CDC. As a reminder, this facility is about 300 meters west of the Seafood market where CoV-2 was originally thought to originate.

This will not be used to adjust the likelihoods.

Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: The careful words of Dr. Shi do NOT say she did not have SARS-CoV-2 at the WIV.

This Figure contains quotes from an article about Dr. Shi and her reaction to the beginning of the COVID-19 pandemic.



Notice in the last frame Dr. Shi says two strange sentences:

Sentence 1: “...she frantically went through her own laboratory’s records from the past few years to check for any mishandling of experimental materials, **especially during disposal.**”

Why did she mention disposal? If you don’t know what you are looking for this, “especially during disposal,” is a bit of an odd qualifier. Other evidence from Wuhan suggests that, in fact, disposal may have been a likely source of the accidental lab release.

Sentence 2: “She breathed a sigh of relief when the results came back: none of the sequences matched those of the viruses her team had sampled from bat caves.”

If Dr. Shi had created SARS-CoV-2 as a chimera, perhaps starting with one of those cave viruses, of course you would no longer have a sequence match. This is a probably truthful statement that leaves open the question of lab creation.

This will not be used to adjust the likelihoods.

Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

Evidence: The Good, the Bad, and the Ugly: a review of SARS Lab Escapes¹²³

In 2003–04, in the wake of the SARS epidemics, there were multiple cases of laboratory acquired infection (LAI) with SARS within just a few months: first in a P3 in Singapore, then in a military P4 in Taipei and last a protracted case in a P3 in Beijing. The '[WHO SARS Risk Assessment and Preparedness Framework](#)' has a good summary of these lab accidents:

Since July 2003, there have been four occasions when SARS has reappeared. Three of these incidents [note: Singapore, Taipei and Beijing] were attributed to breaches in laboratory biosafety and resulted in one or more cases of SARS. The most recent laboratory incident [note: in Beijing] resulted in 9 cases, 7 of which were associated with one chain of transmission and with hospital spread. Two additional cases at the same laboratory with a history of illness compatible with SARS in February 2004 were detected as part of a survey of contacts at the facility.[i.1]

This article reviews some of these cases and discusses briefly some of the insights that were gained from these at the time.

Another article along the same lines is, “10 incidents discovered at the nation's biolabs”¹²⁴ This included Dr. Baric’s laboratory in which “(b)etween April 2013 and September 2014, eight individual mouse escapes were reported at the University of North Carolina-Chapel Hill. Several of the mice were infected with either SARS or the H1N1 flu virus.”

Dozens of holes in BSL-4 'spacesuits'

As a key protection against the world's most deadly pathogens, including the Ebola virus, scientists in the BSL-4 labs at the U.S. Army Medical Research Institute of Infectious Diseases (USAMRIID) at Fort Detrick in Maryland wear pressurized, full-body spacesuit-like gear and breathe purified air. Yet those suits ruptured or developed holes in at least 37 incidents during a 20-month period in 2013 and 2014, according to lab incident reports obtained by USA TODAY under the federal Freedom of Information Act.

This will contribute a 51%/49% contribution in favor of laboratory compared to zoonotic origin. There will be no confidence adjustment. The results from the calculations are shown below.

Evidence or process	Zoonotic Origin (ZO)	Laboratory Origin (LO)
Starting likelihood	0.011	0.989
The history of SARS laboratory accidents is consistent with the laboratory origin hypothesis		0.51
Impact of this evidence		Increases the likelihood of LO by 51/49 = 1.041
Impact of evidence calculation		$1.041 \times 0.989 = 1.030$
Normalize this step of analysis	$0.011 / (0.011 + 1.030) = 0.011$	$1.030 / (0.011 + 1.030) = 0.989$

Adjusted likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹²³ <https://gillesdemanuef.medium.com/the-good-the-bad-and-the-ugly-a-review-of-sars-lab-escapes-898d203d175d>

¹²⁴ <https://www.usatoday.com/story/news/2015/05/29/some-recent-us-lab-incidents/25258237/>

Evidence: Drs. Shi and Daszak use Wuhan residents as negative controls for zoonotic coronavirus seroconversion¹²⁵

"As a control, we collected 240 serum samples from random blood donors in **Wuhan >1000 km away from Jinning & where inhabitants have a much lower likelihood of contact with bats due to its urban setting**" [emphasis added]. As expected, 0/240 samples from the patients from Wuhan had a positive serological evidence of prior coronavirus infection.

"The 2.7% seropositivity for the high-risk group of residents living in close proximity to bat colonies suggests that spillover is a **relatively rare event**, however this depends on how long antibodies persist in people, since other individuals may have been exposed and antibodies waned."

In this paper from 2018, Drs. Shi and Daszak conclude that bat-to-human transfer is relatively rare for high-risk people living in close proximity to bat colonies and much less likely in Wuhan, a conclusion that does not support a hypothesis of bat-to-human transmission.

This will not be used to adjust the likelihoods.


Current likelihood: Zoonotic origin (0.2%), laboratory origin (99.8%).

¹²⁵ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6178078/>

Evidence. The Bat Coronavirus RaTG13 has the Unique Genome Sequences Necessary to be the Precursor of SARS-CoV-2 Using the ‘No See ‘Em’ Synthetic Biology Technology. *The probability that RaTG13 acquired these ‘No See ‘Em’ synthetic biology assembly sequences in nature is one in a billion.*

Summary.

- Synthetic biology techniques, like the engineered “No See ‘Em’¹²⁶ restriction enzyme-enabled insertion method,¹²⁷ have been developed that, by design, extinguish the fingerprints of the insertion when only looking at the final genome.
- The use of these techniques is revealed however, if the precursor-product genome pair of such an insertion is available for inspection.
- **Hypothesis: the unique features of the SARS-CoV-2 Spike Protein, the receptor binding domain ACE2 contact amino acid residue region and the polybasic (furin) cleavage site, are the product of a genome insertion sequence into RaTG13 using engineered Esp3I restriction enzyme sites, the so-called, ‘No See ‘Em,’ technology.**
- An example of the ‘No See’m’ Technology is shown below, taken from Baric and Sim.¹ By placing the restriction sites symmetrically on both strands of the cDNA, the resulting insertion no longer contains the identifying restriction site nts.



- According to Baric and Sims¹ “the type IIS restriction enzyme, Esp3I, recognizes an asymmetric sequence and makes a staggered cut 1 and 5 nucleotides downstream of the recognition sequence, leaving 256, mostly asymmetrical, 4-nucleotide overhangs

¹²⁶ Variably spelled ‘No See ‘Em,’ ‘No See ‘um,’ and ‘No See’m.’

¹²⁷ https://www.researchgate.net/publication/8119695_Development_of_mouse_hepatitis_virus_and_SARS-CoV_infectious_cDNA_constructs

(GCTCTCN#NNNN). As identical Esp3I sites are generated every ~1,000,000 base pairs or so in a random DNA sequence, most restricted fragments usually do not self-assemble.”

- Examination of RaTG13 identified two Esp3I cleavage sites in the Spike Protein gene, at nts 1366 and 2941 (positions 22,910 and 24,485 in the entire genome).
- As expected from the above rarity of such sites in an approximately 3800 nt gene, SARS-CoV-2 has no Esp3I sites in its SP gene. Neither do twelve other coronaviruses, including SARS-CoV-1, MERS, and other related human or bat coronaviruses.
- From all of the species other than bat RaTG13 gene source, the frequency of Esp3I sites **at any location** is 2 in 54,131 nucleotides or 0.000036947. If we assume the possibility of the occurrence of such a site at a given nucleotide is independent of any other nucleotide, then it is possible to use a binomial distribution calculation to determine the probability of 2 Esp3I sites in 3809 nucleotides for the bat RaTG13 gene. This calculation yields a probability of at least 2 sites anywhere in the Spike Protein gene of 0.009 or about one in a hundred. The probability of exactly 2 sites is 0.0086.¹²⁸
- The 5’ restriction site in RaTG13 begins at aa residue 455L, identified by [Andersen et al, Nature, 2020](#), as the start of the “receptor-binding domain ACE2 contact residues.” The downstream amino acids from this site are critical for why RaTG13 has such poor affinity for human ACE2 and the substitutions in CoV-2 are precisely why CoV-2 has such high affinity for human ACE2, why CoV-2 seems so ‘preadapted’ to human infections, etc. So this is the most important part of CoV-2 in explaining its ACE2 binding and infectivity. Further downstream is arguably the second most important site, the polybasic (furin) cleavage site.¹²⁹ Polybasic cleavage sites have not been observed in related ‘lineage B’ betacoronaviruses,’ according to [Andersen et al, Nature, 2020](#), and so there has been much speculation about how this site was acquired.
- The 3’ restriction site in RaTG13 is at residue 980L. There is no protein-based rationale for this position.
- Comparing the nt sequences between RaTG13 and CoV-2, at the 5’ restriction site, they are two codons in which only 2 of 6 nt bases are shared but, despite this low nt sequence homology, they are in fact synonymous base substitutions.
- Comparing the nt sequence between RaTG13 and CoV-2 at the 3’ restriction site, this site has 5 of 6 identical nts with a single synonymous change in CoV-2 which destroys the restriction site. This is the only such five nt site in the RaTG13 spike protein gene and so

¹²⁸ Statistical analysis provided by Dr. Martin Lee, PhD, Adjunct Professor of Statistics, UCLA Fielding School of Public Health, UCLA, Los Angeles, CA.

¹²⁹ <https://www.biorxiv.org/content/10.1101/2020.08.26.268854v1>

is the easiest site in which a one nt substitution can create or destroy an Esp3I restriction site.

- The probability of having the restriction sites at **exactly these locations** can also be calculated.² Since there are 3809 nucleotides in the RaTG13 genome then, 3807 would not have a restriction site with probability $(1-0.000036947)$, which was determined from the frequency of these restriction sites in other species. The other two sites would have this restriction site with probability 0.000036947 . So the overall probability of this configuration has a probability of: $(1-0.000036947)^{3807} \times (0.000036947)^2 = 3.343 \times 10^{-10}$. This is a frequency of these site at their exact location being here from a natural process of approximately one in a billion.
- Dr. Zhengli-Li Shi, of the Wuhan Institute of Virology, collected the bat virus RaTG13 in 2013 and sequenced it between 2014 and 2018. In 2015, Dr. Shi and colleagues have also used the 'No See 'Em' technology' with a similar restriction enzyme, BgII, in the SARS-CoV reverse genetics system to generate chimeric coronaviruses. In that paper, they inserted a spike protein gene from a bat coronavirus into a mouse-adapted coronavirus, with a 'gain-of-function' phenotypic change.¹³⁰

- **In conclusion:**

- **The bat coronavirus RaTG13 has two rare, Esp3I restriction sites strategically located to permit insertion of a genetic sequence that codes for the unique features of the SARS-CoV-2 Spike Protein, its receptor binding contact amino acids and its polybasic (furin) cleavage site, using the 'No See 'Em' synthetic biology techniques.**
- **This specific synthetic biology laboratory technique has been successfully performed previously by Wuhan Institute of Virology scientists to increase coronavirus infectivity.**
- **The probability these two sites are present and in their exact location in RaTG13 by an act of nature is one in a billion.**

¹³⁰ <https://www.nature.com/articles/nm.3985>

Text-Table. A record of the EspI restriction enzyme sites in the Spike Protein (SP) genes of fifteen coronaviruses, including RaTG13 and SARS-CoV-2. RaTG13 is unique in having two such sites, with SARS-CoV-2 and eleven other coronaviruses having no such site in the SP gene. The restriction sites were identified with the RestrictionMapper site algorithm: <http://www.restrictionmapper.org/>.

Species	Spike Protein (SP) Gene Source	Nt Size of SP Gene	Esp3I Site Location in Spike Protein Gene	Reference
Bat	Bat Coronavirus RaTG13 from WIV	3809	1366, 2941 (22910, 24485 in genome)	
Human	SARS-CoV-2 Reference Sequence	3821	None	
Bat	Rhinolophus affinis coronavirus isolate LYRa11	3779	None	Daszak and Shi paper
Bat	Bat SARS coronavirus HKU3-1	3728	None	Daszak and Shi paper
Bat	SARS-like coronavirus isolate bat-SL-CoVZC45	3740	None	Third Military University publication
Bat	SARS-like coronavirus bat-SL-CoVZXC21	3737	None	Third Military University publication
Bat	hCoV-19/bat/Yunnan/RmYN02/2019	3873	None	Wild bat coronavirus with apparent furin-like insert
Bovine	Bovine coronavirus strain Quebec	4091	None	
Human	Human coronavirus HKU1 strain	4070	3208	
Human	MERS Reference Sequence	4061	None	
Human	Human coronavirus OC43 strain	4079	None	
Human	Human coronavirus 229E strain	3512	None	
Human	Human Coronavirus NL63 Reference Sequence	4070	None	
Human	SARS 2003 coronavirus ZJ0301	3767	None	
Pangolin	Pangolin coronavirus isolate PCoV GX-P4L	3803	3351	
Human	SARS-CoV-1 Urbani	3767	None	

Figure. A comparison of the RaTG13 Spike Protein gene (Query) and the SARS-CoV-2 Reference Sequence (Sbjct) showing the only two Esp3I restriction enzyme cleavage site, both present in RaTG13 but absent in SARS-CoV-2. The restriction sites were identified with the RestrictionMapper site: <http://www.restrictionmapper.org/>. The 5' cleavage site is strategically located at the beginning of the receptor binding domain ACE2 contact residues. Despite four of six nt are different these are synonymous changes.

Query	1321	ATTGATGCAAAAGAGGGCGGTAATTTTAACTATCTTTAC	CGTCTC	TTTAGAAAAGCTAAT	1380
Sbjct	1321	CTTGATTCTAAGGTTGGTGGTAATTATAATTACCTGTATAGATTGTTTAGGAAGTCTAAT			1380

The 3' cleavage site is the only downstream -CGTCTN- sequence found in the CoV-2 Spike Protein, making it unique.

Query	2927	TCCTTTCA	CGTCTC	GACAAAGTTGAGGCTGAAGTGCAGATTGACAGGTTGATCACAGGCA	2986
Sbjct	2939	TCCTTTCACGTCTTGACAAAGTTGAGGCTGAAGTGCAAATTGATAGGTTGATCACAGGCA			2998

Figure. Comparison of Spike Protein amino acid sequence between RaTG13 (Query) and SARS-CoV-2 (Sbjct). Amino acid substitutions in CoV-2 are shown in red, single letter abbreviation. Green band; receptor binding domain. Blue band; receptor binding domain ACE2 contact residues ([Andersen et al, Nature, 2020.](#)). Purple band; polybasic (furin) cleavage site. Red brackets; Esp3I cleavage sites in RaTG13.

Score	Expect	Method	Identities	Positives	Gaps
2565 bits(6648)	0.0	Compositional matrix adjust.	1240/1273(97%)	1252/1273(98%)	4/1273(0%)
Query 1	1	MFVFLVLLPLVSSQCVNLTTRTQLPPAYTNSSTRGVVYPDKVFRSSVLHLTQDLFLPFFS			60
Sbjct 1	1F.....S.....			60
Query 61	61	NVTWFHAIHVSGTNGIKRFDNPVLPFNDGVYFASTEKSNIIRGWIFGTTLDSTQSLIV			120
Sbjct 61	61T.....			120
Query 121	121	NNATNVVIKVFCEQFCNDPFLGVVYHKNKSWMESEFRVSSANNCTFEYVSQPFLMDLE			180
Sbjct 121	121			180
Query 181	181	GKQGNFKNLREFVFKNIDGYFKIYSKHTPINLVRDLPPGFSALEPLVDLPIGINITRFQT			240
Sbjct 181	181Q.....			240
Query 241	241	LLALHRSYLTPGDSSSGNTAGAAAYVVGYLQPRTEFLKYNENGTITDAVDCALDPLSETK			300
Sbjct 241	241			300
Query 301	301	CTLKSFTVEKGIYQTSNFRVQPTDSIVRFPNITNLCPFGEVFNATTFASVYAWNKRISN			360
Sbjct 301	301E.....R.....			360
Query 361	361	CVADYSVLNYSFSTFKCYGVSPTKLNLCFTNVYADSFVITGDEVQRQIAPGQTGKIAD			420
Sbjct 361	361A.....R.....			420
Query 421	421	YNYKLPDDFTGCVIAWNSKHIDAKEGGNFNYLYRLFRKANLKPFERDISTEIQAGSKPC			480
Sbjct 421	421NNL.S.V...Y...S.....T.....			480
Query 481	481	NGQTGLNCYYPYLYRGFYPTDGVGHQPYRVVLSFELLNAPATVCGPKKSTNLVKNKCVN			540
Sbjct 481	481	..VE.F...F..QS...Q..N...Y.....H.....			540
Query 541	541	FNFNGLTGTGVLTESNKKFLPFQQFGRDIADTTDAVRDPQTLEILDITPCSFGGVSVITP			600
Sbjct 541	541			600
Query 601	601	GTNASNQVAVLYQDVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIGAHEVNNNSY			660
Sbjct 601	601	...T.....			660
Query 661	661	ECDIPIGAGICASYQTQNS----RSVASQSIIAYTMSLGAENSVAYSNNISIAIPTNFTI			716
Sbjct 661	661PRRA.....			720
Query 717	717	SVTTEILPVSMTKTSVDCTMYICGDSTECNLLQYGSFCTQLNRALTGIAVEQDKNTQE			776
Sbjct 721	721			780
Query 777	777	VFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPSKRSFIEDLLFNKVTLADAGFIKQYGDC			836
Sbjct 781	781			840
Query 837	837	LGDIAARDLICAQKFNGLTVLPLLTDEMAIQYSALLAGTITSGWTFGAGAAQIPFAM			896
Sbjct 841	841			900
Query 897	897	QMAYRFNGIGVTQNVLYENQKLIANQFNSAIGKIQDSLSTASALGKLQDVVNQNAQALN			956
Sbjct 901	901			960
Query 957	957	TLVKQLSSNFGAISSVLNDILSRIDKVEAEVQIDRLITGRLQSLQTYVTQQLIRAAEIRA			1016
Sbjct 961	961			1020
Query 1017	1017	SANLAATKMSECVLGQSKRVDFCGKGYHLMSFPQSAPHGVVFLHVTYVPAQEKNFTTAPA			1076
Sbjct 1021	1021			1080
Query 1077	1077	ICHGKAHFPREGVFSNGTHWVFTQRNFYEPQIITDNTFVSGSCDVVIGIVNNTVYDP			1136
Sbjct 1081	1081N.....			1140
Query 1137	1137	LQPELDSFKEELDKYFKNHTSPDVLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDL			1196
Sbjct 1141	1141			1200
Query 1197	1197	QELGKYEQYIKWPWYIWLGFIAGLIAIMVTIMLCMTSCCSCLKGCCSCGSCCKFDEDD			1256
Sbjct 1201	1201V.....			1260
Query 1257	1257	SEPVLLKGVKLHYT	1269		
Sbjct 1261	1261	1273		

Because it has not been established that RaTG13 was the precursor of CoV-2 this evidence statement will not be used at this time to adjust the likelihoods of the origin. If additional information is obtained at a later date this may be revisited.

Likelihood from prior state is unchanged following this evidence analysis:

Zoonotic origin (0.2%) and laboratory origin (99.8%)

Evidence. Location, location, location: Based on the distance between known SARS-CoV-1 laboratory-acquired infections and the hospital of admission of the infected personnel, the WIV is within the expected hospital catchment for a CoV-2 LAI

Hypothesis. Laboratory-acquired infections (LAI) have the property that the hospital of admission of the personnel from the laboratory with the acquired infection are close together, specifically they are within 24.64 km from the laboratory.

Prior data from SARS-CoV-1. There were four LAIs of SARS-CoV-1 that can be used to determine the distance between the laboratory where the infection occurred and the hospital of first admission. The data are here:

SARS-CoV-1 Laboratory Acquired Infection (LAI)	Hospital of admission	Distance (Google Maps)	
In September 2003, a 27-year-old student from the National University of Singapore (NUS) was infected with the SARS virus due to improper experimental procedures	Singapore General Hospital (SGH)	6.3 km	
Baiji Mountain, Sanxia, Taiwan	Taiwan Hoping Hospital, Taipei, Taiwan	27.8 km	
Ne100 Yingxin Street, Xicheng District, Beijing	Union Hospital, Beijing, China	7.3 km	
Ne100 Yingxin Street, Xicheng District, Beijing	Friendship Hospital, Beijing, China	17.6 km	
		mean = 14.75	
		SD = 10.1	
		95% Confidence Interval	14.75 ±9.887

Based on these four cases, the 95% upper confidence limit for the distance from LAI patients to the hospitals of admission is 24.6 km of the laboratory where the infection was acquired.

SARS-CoV-2. Although it is not clear which hospital the first patient was admitted to the following Text-Table contains all likely candidates.

SARS-CoV-2 Potential LAI Source	Hospital of admission	Distance (Google Maps)	Probability of being closer than the average results for SARS-CoV-1	Probability of being farther than the average results for SARS-CoV-1
Wuhan Institute of Virology, Wuhan, China	PLA Hospital, NO. 627 Wuluo Road, Wuchang District, Wuhan, China	4.8 km	0.094	0.906
Wuhan Institute of Virology, Wuhan, China	Wuhan Central Hospital, Wuhan, China	9.1 km	0.338	0.662
Wuhan Institute of Virology, Wuhan, China	Zhongnan Hospital, Wuhan, China	2.8 km	0.019	0.981
Wuhan Institute of Virology, Wuhan, China	Tongji Hospital, Wuhan, China	5.1 km	0.109	0.891
Wuhan Institute of Virology, Wuhan, China	Hubei Maternity and Child Health Care Hospital, Wuhan, China	4.4 km	0.075	0.925
Hypothesis: Given the distance from the SARS-CoV-1 laboratory where an LAI occurred to the hospital of admission for the lab workers who became infected, what is the probability that CoV-2 is also an LAI, given the distance from the hospitals where the first patients were seen to the WIV, the hypothesized source.			Probability calculations based on the use of a log-normal distribution for distances	Probability calculations based on the use of a log-normal distribution for distances

Based on the data for actual LAI for SARS-CoV-1 the distance between the WIV and the hospitals of admission for CoV-2 is consistent with the WIV being the origin for the LAI. There is no evidence the putative LAI for CoV-2 is any different than the known LAIs for CoV-1.

This evidence is not independent of other evidence that is based on location and so it cannot be used independently in the Bayesian analysis. It is included here for completeness.

Likelihood from prior state is unchanged following this evidence analysis:

Zoonotic origin (0.2%) and laboratory origin (99.8%)



Evidence. Dr. Shi successfully identifies a laboratory-acquired infection outbreak from Hanta virus in laboratory rodents.

Infection, Genetics and Evolution 10 (2010) 638–644

Contents lists available at ScienceDirect

Infection, Genetics and Evolution

journal homepage: www.elsevier.com/locate/meegid

Hantavirus outbreak associated with laboratory rats in Yunnan, China

Yunzhi Zhang^{a,b}, Hailin Zhang^{b,*}, Xingqi Dong^b, Junfa Yuan^a, Huajun Zhang^a, Xinglou Yang^a, Peng Zhou^a, Xingyi Ge^a, Yan Li^a, Lin-Fa Wang^c, **Zhengli Shi^{a,**}**

^a State Key Laboratory of Virology, **Wuhan Institute of Virology**, Chinese Academy of Sciences, Wuhan, People's Republic of China
^b Yunnan Institute of Endemic Diseases Control and Prevention, Dali, People's Republic of China
^c Commonwealth Scientific and Industrial Research Organisation Livestock Industries, Geelong, Victoria, Australia

<p>ARTICLE INFO</p> <p><i>Article history:</i> Received 16 November 2009 Received in revised form 20 February 2010 Accepted 30 March 2010 Available online 7 April 2010</p> <p><i>Keywords:</i> Hemorrhagic fever with renal syndrome Hantavirus Laboratory rats Recombination</p>	<p>ABSTRACT</p> <p>An outbreak of hemorrhagic fever with renal syndrome occurred among students in a college (College A) in Kunming, Yunnan province, China in 2003. Subsequent investigations revealed the presence of hantavirus antibodies and antigens in laboratory rats at College A and two other institutions. Hantavirus antibodies were detected in 15 additional individuals other than the index case in these three locations. Epidemiologic data indicated that the human infections were a result of zoonotic transmission of the virus from laboratory rats. A virus was isolated from rats in College A and the full-length genome sequence revealed that this was a new Hantaan virus isolate, designated strain KY. Sequence analysis of the three genome segments indicated that this new isolate is a reassortant derived from human and rat Hantaan viruses. Further sequence analysis of the medium (M) genome segment revealed that it originated from a recombination event between two rat Hantaan virus lineages.</p> <p>© 2010 Elsevier B.V. All rights reserved.</p>
---	--


The significance of this evidence is that it demonstrates the methods used by Dr. Shi and the WIV to solve a laboratory-acquired infection outbreak. The methods described herein should be applied to the WIV in order to determine if CoV-2 was also a laboratory-acquired infection.

This will not be used to directly advance the Bayesian analysis.

Likelihood from prior state is unchanged following this evidence analysis:

Zoonotic origin (0.2%) and laboratory origin (99.8%)

Evidence. Bats hibernate when the temperature is below 10.5 C;¹³¹ in Hubei province that begins in September and ends in May.



Based on this evidence, they would have been hibernating at the time of the first human outbreak in the fall of 2019. Since this evidence is cumulative to the prior evidence from Dr. Shi that the bat host species for CoV-2 does not live in Hubei Province it will not be used to change the Bayesian analysis.

Likelihood from prior state is unchanged following this evidence analysis:

Zoonotic origin (0.2%) and laboratory origin (99.8%)

¹³¹ <https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-7998.1971.tb01323.x>

Wuhan Institute of Virology analysis of lavage specimens from ICU patients at Wuhan Jinyintan Hospital in December 2019 contain both SARS-CoV-2 and adenovirus vaccine sequences consistent with a vaccine challenge trial

Summary. The most significant evidence provided herein is the finding from RNA-Seq performed by the Wuhan Institute of Virology (WIV) of lavage patient samples collected on December 30, 2019.¹³² These ICU patients were the subject of the seminal paper, entitled, “A pneumonia outbreak associated with a new coronavirus of probable bat origin,” from Dr. Zhengli Shi and colleagues that first characterized SARS-CoV-2.¹³³ This author has confirmed that the RNA-Seq of all five patients contained SARS-CoV-2 sequences.

Surprisingly the specimens also contained the adenovirus “pShuttle” vector, developed by Chinese scientists in 2005 for SARS-CoV-1.¹³⁴ Two immunogens were identified, the Spike Protein gene of SARS-CoV-2 and the synthetic construct H7N9 HA gene.¹³⁵ Hundreds of perfectly homologous (150/150) raw reads suggest this is not an artifact. Reads that cross the vector-immunogen junction are identified. While adenovirus is a common infection the wildtype viruses have low homology to the vaccine vector sequence, by design, to avoid rejection of the vaccine due to prior exposure to wildtype adenoviruses.

Two patients from the same hospital who had bronchial lavage on the same day but had their specimens sent to the Hubei CDC did not have adenovirus vaccine sequences.

Three explanations come to mind from this evidence:

1. These represent sample preparation artifacts at the WIV, such as sample spillover on the sequencer.
2. These patients were admitted with an unknown infection, were not responding to the treatment protocols for a infection of unknown origin, and they were vaccinated with an experimental vaccine in a desperate but compassionate therapeutic “Hail Mary.”
3. A clinical trial of a combination¹³⁶ influenza/SARS-CoV-2 vaccine was being conducted and an accidental release into Wuhan occurred.

Only WIV scientists and Chinese authorities can answer these questions. Until the evidence of the adenovirus sequences has been confirmed by other scientists, this author will not include this evidence in the Bayesian analysis.

Obviously if a vaccine containing the Spike Protein of SARS-CoV-2 was being administered to patients in Wuhan in December 2019 the question of laboratory origin is a settled matter.

¹³² The detailed evidence for the adenovirus vaccine sequences is given at the end of this document.

¹³³ <https://www.nature.com/articles/s41586-020-2012-7>

¹³⁴ <https://www.ncbi.nlm.nih.gov/nuccore/AY862402.1>

¹³⁵ <https://www.ncbi.nlm.nih.gov/nuccore/KY199425.1/>

¹³⁶ The proposal that this was, in fact, a combination vaccine was made by H. Lawrence Rimmel, Department of Pathology, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands.

Introduction. Following the 2003 SARS epidemic, Liu et al. developed an adenoviral expression vector of a truncated S1 subunit of SARS-CoV spike protein that resulted in specific humoral immune responses against SARS-CoV in rats.¹³⁷ This same vector was used to create the CoV-2 adenovirus vector vaccine.

In order to test the hypothesis that CoV-2 began in the PLA Hospital as a vaccine challenge clinical trial that went awry, RNA-Seq raw reads from nasopharyngeal specimens of Wuhan COVID patients (Table below) were blasted against the published genome sequence of the SARS-CoV-1 vaccine (GenBank [AY862402.1](https://www.ncbi.nlm.nih.gov/nuccore/AY862402.1)). I used the SARS-CoV-1 vaccine because the PLA CoV-2 vaccine has not been published at this time.


Adenovirus sequences detected	GenBank URL	GenBank Biosample URL	GISAID ID	CoV-2 Isolate	Sequencing Institution	Clinical Information from GISAID
>100	SRX7730879	SAMN14082200	EPI_ISL_402130	WIV07; Lineage B; mutations NSP3 D1761A, NSP4 T327I; passage original	Wuhan Institute of Virology, Chinese Academy of Sciences	56 y, male, hospitalized, ICU10G, 20 Dec 2019
>100	SRX7730880	SAMN14082196	EPI_ISL_402127	WIV02; Lineage B; mutations NSP16 D220N; passage original	Wuhan Institute of Virology, Chinese Academy of Sciences	32 y, male, hospitalized, ICU4G, outbreak 19 Dec 2019
>100	SRX7730881	SAMN14082197	EPI_ISL_402124	WIV04; Lineage B; no mutations; passage original	Wuhan Institute of Virology, Chinese Academy of Sciences	49 y, female, hospitalized, ICU-6, outbreak 27 Dec 2019, Retailer at Huanan Seafood Wholesale Market, patient alive
>100	SRX7730882	SAMN14082198	EPI_ISL_402128	WIV05; Lineage B; NSP3 G1433S, NSP16 K160R; passage original	Wuhan Institute of Virology, Chinese Academy of Sciences	52 y, female, hospitalized, ICU8G, outbreak 22 Dec 2019; recovered
>100	SRX7730883	SAMN14082199	EPI_ISL_402129	WIV06; Lineage B; no mutations; original passage	Wuhan Institute of Virology, Chinese Academy of Sciences	40 y, male, hospitalized, ICU9G, 25 Dec 2019
>100	SRX7730884	SAMN14082200	EPI_ISL_402130	WIV07; Lineage B; mutations NSP3 D1761A, NSP4 T327I; passage original	Wuhan Institute of Virology, Chinese Academy of Sciences	56 y, male, hospitalized, ICU10G, 20 Dec 2019
7 small	SRX7730885	SAMN14082196	EPI_ISL_402127	WIV02; Lineage B; mutations NSP16 D220N	Wuhan Institute of Virology, Chinese Academy of Sciences	32 y, male, hospitalized, ICU, outbreak 19 Dec 2019
1 small one	SRX7730886	SAMN14082197	EPI_ISL_402124	WIV04; Lineage B; no mutations; passage original	Wuhan Institute of Virology, Chinese Academy of Sciences	49 y, female, hospitalized, ICU-6, outbreak 27 Dec 2019, Retailer at Huanan Seafood Wholesale Market, patient alive
Very few	SRX7730887	SAMN14082199	EPI_ISL_402129	WIV06; Lineage B; no mutations; original passage	Wuhan Institute of Virology, Chinese Academy of Sciences	40 y, male, hospitalized, ICU9G, 25 Dec 2019
None	SRX8032202	SAMN14479127	EPI_ISL_412898	hCoV-19/Wuhan/HB-CDC-HB-02/2019	Hubei Provincial Center for Disease Control and Prevention	male, "traveled from Wuhan"
None	SRX8032203	SAMN14479128	EPI_ISL_402132	Wuhan HB-CDC-HB-01/2019; Lineage B; mutation Spike F32I; original passage	Hubei Provincial Center for Disease Control and Prevention	49 y, female, hospitalized

This is not related to the previous claim, now shown to be wrong, that SARS-CoV-2 itself contained adenovirus pShuttle sequences.¹³⁸

¹³⁷ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7114075/>


¹³⁸ <https://sciencefeedback.co/claimreview/2019-novel-coronavirus-2019-ncov-does-not-contain-pshuttle-sn-sequence-no-evidence-that-virus-is-man-made/>

According to Liu: “Adeno-X™ expression system (Clontech Laboratories, Inc.), comprising adenovirus type 5 genome with a deletion in the E1 and E3 regions ($\Delta E1$, 343–3465 bp; $\Delta E3$, 28,756–30,561 bp), was utilized to construct a recombinant adenovirus carrying nucleotides –45 to 1469 of Spike gene of SARS-CoV (Ad-SN) by *in vitro* ligation. This provides an immunogen which encoded a truncated S1 subunit of SARS-CoV S protein (490 N-terminal amino-acid residues),” as shown here:




The expected result would be the finding of RNA-Seq sequence raw reads that were homologous to the two Adenovirus regions but only partially homologous (about 80%) to the SARS-CoV-1 regions.

The results are shown below. All five patients have adenovirus sequences that read through the 5' junction with the immunogen but do not read through the entire gene:




As can be seen above, all five patients have significant portions of the CMV-promoter as well as almost one-half of the truncated Spike Protein gene. This is the expected result if in fact the vaccine was not the previously described SARS-CoV-1, as in that case you would expect through reads covering the entire spike protein gene.


Next, an adenovirus vaccine vector sequence was created by substituting the full CoV-2 spike protein gene into the vector cassette. The results for this construct was much greater coverage within the specimens.



For example, the sequence alignment of patient WIV-05 is shown below. The red arrow and green arrow are at the 5' and 3' junctions of the adenovirus vector sequences and the CoV-2 Spike Protein gene sequence, showing cross junction contigs.



Another surprising finding was the presence of synthetic H7N9 gene sequences, again in all five WIV sequenced patients. The contigs are shown below.



- ☐ [RNA-Seq of Homo sapiens: hCov-19 infected patients Bronchoalveolar lavage uid](#)
- 1. 1 ILLUMINA (Illumina MiSeq) run: 5.8M spots, 1.7G bases, 634.3Mb downloads
Accession: SRX8032203
- ☐ [RNA-Seq of Homo sapiens: hCov-19 infected patients Bronchoalveolar lavage uid](#)
- 2. 1 ILLUMINA (Illumina MiSeq) run: 5.2M spots, 1.6G bases, 583.4Mb downloads
Accession: SRX8032202
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 3. 1 ILLUMINA (Illumina MiSeq) run: 5.2M spots, 1.5G bases, 772.9Mb downloads
Accession: SRX7730887
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 4. 1 ILLUMINA (Illumina MiSeq) run: 5.2M spots, 1.5G bases, 768.3Mb downloads
Accession: SRX7730886
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 5. 1 ILLUMINA (Illumina MiSeq) run: 8.3M spots, 2.2G bases, 1.2Gb downloads
Accession: SRX7730885
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 6. 1 ILLUMINA (Illumina HiSeq 3000) run: 38.5M spots, 11.5G bases, 7.1Gb downloads
Accession: SRX7730884
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 7. 1 ILLUMINA (Illumina HiSeq 3000) run: 29.7M spots, 8.9G bases, 5.6Gb downloads
Accession: SRX7730883
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 8. 1 ILLUMINA (Illumina HiSeq 3000) run: 34.3M spots, 10.3G bases, 6.4Gb downloads
Accession: SRX7730882
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 9. 1 ILLUMINA (Illumina HiSeq 1000) run: 61.3M spots, 18.4G bases, 11.4Gb downloads
Accession: SRX7730881
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 10. 1 ILLUMINA (Illumina HiSeq 3000) run: 67.1M spots, 20.1G bases, 12.6Gb downloads
Accession: SRX7730880
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 11. 1 ILLUMINA (Illumina MiSeq) run: 3.6M spots, 1G bases, 548.1Mb downloads
Accession: SRX7730879

The WIV entry with the greatest read depth, Number 10 above, is described below:

SRX7730880: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid

1 ILLUMINA (Illumina HiSeq 3000) run: 67.1M spots, 20.1G bases, 12.6Gb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the MGIEasy RNA Library Prep Set (96 RXN) (Cat. No.: 1000006384). Paired-end (150 bp) sequencing of the RNA library was performed on the MGISEQ-2000RS platform.

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads

[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)

[show Abstract](#)

Sample:

[SAMN14082196](#) • [SRS6151291](#) • [All experiments](#) • [All runs](#)

Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:

Name: WIV02-2

Instrument: Illumina HiSeq 3000

Strategy: RNA-Seq

Source: METAGENOMIC

Selection: RANDOM

Layout: PAIRED

Runs: 1 run, 67.1M spots, 20.1G bases, [12.6Gb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092063	67,083,195	20.1G	12.6Gb	2020-02-16

Unexpectedly, over 100 sequences producing significant alignment were identified:

BLAST » blastn suite-SRA » results for RID-S76CAHY001R

Home Recent Results Saved Strategies Help

[< Edit Search](#) [Save Search](#) [Search Summary](#)

How to read this report? [BLAST Help Videos](#) [Back to Traditional Results Page](#)

Job Title [gb|AY862402.1](#)

RID [S76CAHY001R](#) Search expires on 10-13 07:47 am [Download All](#)

Program BLASTN [Citation](#)

Database SRA [See details](#)

Query ID [AY862402.1](#)

Description Expression vector pShuttle-SN, complete sequence

Molecule type nucleic acid

Query Length 5607

Other reports [Distance tree of results](#) [MSA viewer](#)

Descriptions Graphic Summary Alignments


Sequences producing significant alignments Download Manage Columns Show 100

☒ select all 100 sequences selected

Graphics Distance tree of results

	Description	Max Score	Total Score	Query Cover	E value	Per. Ident	Accession
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.66604450.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.66455070.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.63120099.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.63120099.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.62730385.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.61105639.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.60748776.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.60011402.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.59155252.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.59155252.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.58126883.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.67671560.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.67484454.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.56079039.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.56036194.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.55663455.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.55111963.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.53777284.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.53679813.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.52965281.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.51414708.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.51019881.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.50609371.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.50609371.1

A graphical display of the alignments shows they are not in the Spike Protein region (961 to 2507) of the adenovirus vector but outside of those regions.



An examination of individual reads show 100% homology over the entire 150 nt segments and outside of the Spike Protein region. The first set of reads are immediately downstream of the Spike Protein segment. The other read is from the region is from the 5' boundary of the Adenovirus vector with the Spike Protein region.

Download Graphics SRA				
SRX7730880				
Sequence ID: SRA:SRR11092063.66604450.1 Length: 150 Number of Matches: 1				
Range 1: 1 to 150 Graphics Next Match Previous Match				
Score	Expect	Identities	Gaps	Strand
278 bits(150)	2e-70	150/150(100%)	0/150(0%)	Plus/Plus
Query 2536	CCCGTGCCTTCCTTGACCCTGGAAGGTGCCACTCCCCTGTCCTTTCCCTAATAAAATGAG	2595		
Sbjct 1	CCCGTGCCTTCCTTGACCCTGGAAGGTGCCACTCCCCTGTCCTTTCCCTAATAAAATGAG	60		
Query 2596	GAAATTGCATCGCATTGTCTGAGTAGGTGTCATTCTATTCTgggggggtgggggtggggCAG	2655		
Sbjct 61	GAAATTGCATCGCATTGTCTGAGTAGGTGTCATTCTATTCTGGGGGGTGGGGTGGGGCAG	120		
Query 2656	GACAGCAAGGGGGAGGATTGGGAAGACAAT	2685		
Sbjct 121	GACAGCAAGGGGGAGGATTGGGAAGACAAT	150		

Download Graphics SRA				
SRX7730880				
Sequence ID: SRA:SRR11092063.66455076.2 Length: 150 Number of Matches: 1				
Range 1: 1 to 150 Graphics Next Match Previous Match				
Score	Expect	Identities	Gaps	Strand
278 bits(150)	2e-70	150/150(100%)	0/150(0%)	Plus/Minus
Query 3290	CGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCCTTATCC	3349		
Sbjct 150	CGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCCTTATCC	91		
Query 3350	GGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCC	3409		
Sbjct 90	GGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCC	31		
Query 3410	ACTGGTAACAGGATTAGCAGAGCGAGGTAT	3439		
Sbjct 30	ACTGGTAACAGGATTAGCAGAGCGAGGTAT	1		

[Download](#)
[Graphics](#)
[SRA](#)

SRX7730880

Sequence ID: [SRA:SRR11092063.50609371.2](#) Length: 150 Number of Matches: 1

Range 1: 1 to 150 [Graphics](#) [Next Match](#) [Previous Match](#)

Score	Expect	Identities	Gaps	Strand
278 bits(150)	2e-70	150/150(100%)	0/150(0%)	Plus/Plus

```

Query 703 CAAGTCTCCACCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCAAATCAACGGGACT 762
          |||
Sbjct 1 CAAGTCTCCACCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCAAATCAACGGGACT 60

Query 763 TTCCAAAATGTCGTAACAACCTCCGCCCATTTGACGCAAATGGGCGGTAGGCGTGTACGGT 822
          |||
Sbjct 61 TTCCAAAATGTCGTAACAACCTCCGCCCATTTGACGCAAATGGGCGGTAGGCGTGTACGGT 120

Query 823 GGGAGGTCTATATAAGCAGAGCTCTCTGGC 852
          |||
Sbjct 121 GGGAGGTCTATATAAGCAGAGCTCTCTGGC 150
  
```

[Download](#)
[Graphics](#)
[SRA](#)

SRX7730880

Sequence ID: [SRA:SRR11092063.50609371.1](#) Length: 150 Number of Matches: 1

Range 1: 1 to 150 [Graphics](#) [Next Match](#) [Previous Match](#)

Score	Expect	Identities	Gaps	Strand
278 bits(150)	2e-70	150/150(100%)	0/150(0%)	Plus/Minus

```

Query 784 CCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAG 843
          |||
Sbjct 150 CCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAG 91

Query 844 CTCTCTGGCTAACTAGAGAACCCACTGCTTACTGGCTTATCGAAATTAATACGACTCACT 903
          |||
Sbjct 90 CTCTCTGGCTAACTAGAGAACCCACTGCTTACTGGCTTATCGAAATTAATACGACTCACT 31


Query 904 ATAGGGAGACCCAAGCTGGCTAGCGTTTAA 933
          |||
Sbjct 30 ATAGGGAGACCCAAGCTGGCTAGCGTTTAA 1
  
```



To test if this was the actual SARS-CoV-1 vaccine vector and had been given to the patients as an desperate attempt to create immunity during an infection, the Spike Protein region of the vaccine was blasted against the above sample, looking for a near 100% homology. The only reads were a 38 nt segment of 1482-1518, with one gap, as expected. The absence of long reads for the SARS-CoV-1 Spike Protein suggests that this vaccine was not a CoV-1 vaccine.


To test if the homology seen between lavage specimens of patients in Wuhan with the CoV-1 Adenovirus vaccine was due to homology with human sequences the Expression vector was blasted against *Homo sapien* sequences, but no matches were found, as shown below.

BLAST[®] » blastn suite » results for RID-S793VKCV01R

[← Edit Search](#) [Save Search](#) [Search Summary ▼](#)

 Your results are filtered to match records that include: *Homo sapiens* (taxid:9606)

Job Title	AY862402:Expression vector pShuttle-SN, complete...		
RID	S793VKCV01R	Search expires on 10-13 08:34 am	Download All ▼
Program	 Citation ▼		
Database	nt	See details ▼	
Query ID	AY862402.1		
Description	Expression vector pShuttle-SN, complete sequence		
Molecule type	nucleic acid		
Query Length	5607		
Other reports			

 No significant similarity found. For reasons why, [click here](#)

Background. Live attenuated adenovirus vectors for vaccine or gene therapy have been under development for decades.¹³⁹ Adenovirus vaccines against SARS-CoV-1¹⁴⁰ and MERS¹⁴¹ have shown efficacy in animal models of disease. One of the earliest vaccines for CoV-2 is also an adenovirus vector vaccine, developed in collaboration with the PLA.¹⁴²


¹³⁹ <https://www.sciencedirect.com/science/article/pii/S1525001604013425>

¹⁴⁰ <https://www.sciencedirect.com/science/article/pii/S0140673603149628>


¹⁴¹ <https://www.nih.gov/news-events/news-releases/investigational-chimp-adenovirus-mers-cov-vaccine-protects-monkeys>

¹⁴² <https://www.nature.com/articles/d41586-020-02523-x> ; <https://www.nature.com/articles/s41467-020-18077-5>




Below is a blast for sequences from the patients in the same hospital who had lavage on the same day but whose specimens went to the Hubei CDC. There are no adenovirus sequences below.





Or in this specimen.







Or in these specimens.




Below begins the specimens from the WIV.









Above is a blast of Influenza A virus (A/swine/eastern China/HH24/2017(H7N9)) segment 4 hemagglutinin (HA) gene, complete cds in patient WIV-4-2 specimen

[https://www.ncbi.nlm.nih.gov/nucleotide/MG925503.1?report=genbank&log\\$=nuclalign&blast_rank=2&RID=WYG74MH9016](https://www.ncbi.nlm.nih.gov/nucleotide/MG925503.1?report=genbank&log$=nuclalign&blast_rank=2&RID=WYG74MH9016)

<https://www.ncbi.nlm.nih.gov/nuccore/AY862402.1> Expression vector pShuttle-SN, complete sequence

AY862402.1

Specimen 1

[https://www.ncbi.nlm.nih.gov/sra/SRX7730879\[accn\]](https://www.ncbi.nlm.nih.gov/sra/SRX7730879[accn])

<https://trace.ncbi.nlm.nih.gov/Traces/sra/?run=SRR11092064>

Bayesian Analysis of SARS-CoV-2 Origin

Steven C. Quay, MD, PhD

29 January 2021

SRX7730879: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid
1 ILLUMINA (Illumina MiSeq) run: 3.6M spots, 1G bases, 548.1Mb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the NEBNext Ultra II Directional RNA Kit (NEB, USA). Paired-end (150 bp) sequencing of the RNA library was performed on the Miseq platform (Illumina).

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads
[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)
[hide Abstract](#)
Discovery and characterization of a novel human coronavirus from five patients at the early stage of the Wuhan seafood market pneumonia virus outbreak .

Sample:
[SAMN14082200](#) • [SRS6151290](#) • [All experiments](#) • [All runs](#)
Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:
Name: WIV07
Instrument: Illumina MiSeq
Strategy: RNA-Seq
Source: METAGENOMIC
Selection: RANDOM
Layout: PAIRED

Runs: 1 run, 3.6M spots, 1G bases, [548.1Mb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092064	3,566,583	1G	548.1Mb	2020-02-15

ID: 10108892

BLAST[®] » blastn suite-SRA » results for RID-SBB5ZMXM01R [Home](#) [Recent Results](#) [Saved Strategies](#) [Help](#)

[< Edit Search](#) [Save Search](#) [Search Summary](#) [How to read this report?](#) [BLAST Help Videos](#) [Back to Traditional Results Page](#)

Job Title AY862402:Expression vector pShuttle-SN, complete...
RID [SBB5ZMXM01R](#) [Search expires on 10-14 21:33 pm](#) [Download All](#)
Program BLASTN [Citation](#)
Database SRA [See details](#)
Query ID [AY862402.1](#)
Description Expression vector pShuttle-SN, complete sequence
Molecule type nucleic acid
Query Length 5607
Other reports [Distance tree of results](#) [MSA viewer](#)


Filter Results
Percent Identity to
E value to
Query Coverage to
[Filter](#) [Reset](#)

Descriptions [Graphic Summary](#) [Alignments](#)

Sequences producing significant alignments [Download](#) [Manage Columns](#) [Show](#) 100 [Graphics](#) [Distance tree of results](#)

☒ select all 100 sequences selected

	Description	Max Score	Total Score	Query Cover	E value	Per. Ident	Accession
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.3512575.2
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.2917500.1
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.2878891.2
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.2878891.1
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.2655789.2
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.2415875.2
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.1494732.2
<input checked="" type="checkbox"/>	SRX7730879	279	279	2%	2e-72	100.00%	SRA:SRX11092064.1313817.2
<input checked="" type="checkbox"/>	SRX7730879	278	278	2%	9e-72	100.00%	SRA:SRX11092064.3512575.1
<input checked="" type="checkbox"/>	SRX7730879	278	278	2%	9e-72	100.00%	SRA:SRX11092064.2415875.1
<input checked="" type="checkbox"/>	SRX7730879	278	278	2%	9e-72	100.00%	SRA:SRX11092064.1313817.1
<input checked="" type="checkbox"/>	SRX7730879	276	276	2%	3e-71	100.00%	SRA:SRX11092064.2686059.2
<input checked="" type="checkbox"/>	SRX7730879	274	274	2%	1e-70	99.34%	SRA:SRX11092064.1428185.2
<input checked="" type="checkbox"/>	SRX7730879	274	274	2%	1e-70	99.34%	SRA:SRX11092064.1428185.1
<input checked="" type="checkbox"/>	SRX7730879	274	274	2%	1e-70	99.34%	SRA:SRX11092064.734472.2
<input checked="" type="checkbox"/>	SRX7730879	274	274	2%	1e-70	99.34%	SRA:SRX11092064.734472.1
<input checked="" type="checkbox"/>	SRX7730879	274	274	2%	1e-70	99.34%	SRA:SRX11092064.674542.1
<input checked="" type="checkbox"/>	SRX7730879	274	274	2%	1e-70	99.34%	SRA:SRX11092064.612514.1
<input checked="" type="checkbox"/>	SRX7730879	272	272	2%	4e-70	99.33%	SRA:SRX11092064.674542.2
<input checked="" type="checkbox"/>	SRX7730879	268	268	2%	5e-69	98.68%	SRA:SRX11092064.2917500.2
<input checked="" type="checkbox"/>	SRX7730879	268	268	2%	5e-69	98.68%	SRA:SRX11092064.2655789.1
<input checked="" type="checkbox"/>	SRX7730879	267	267	2%	2e-68	98.67%	SRA:SRX11092064.612514.2
<input checked="" type="checkbox"/>	SRX7730879	259	259	2%	3e-66	99.30%	SRA:SRX11092064.3169732.2
<input checked="" type="checkbox"/>	SRX7730879	259	259	2%	3e-66	99.30%	SRA:SRX11092064.3169732.1



SRX7730879Sequence ID: [SRA:SRR11092064.3512575.2](#) Length: **151** Number of Matches: **1**Range 1: 1 to 151 [Graphics](#)[▼ Next Match](#) [▲ Previous Match](#)

Score	Expect	Identities	Gaps	Strand
279 bits(151)	2e-72	151/151(100%)	0/151(0%)	Plus/Minus
Query 4830	ACCTGGAATGCTGTTTTCCCGGGGATCGCAGTGGTGAGTAACCATGCATCATCAGGAGTA	4889		
Sbjct 151	ACCTGGAATGCTGTTTTCCCGGGGATCGCAGTGGTGAGTAACCATGCATCATCAGGAGTA	92		
Query 4890	CGGATAAAATGCTTGATGGTCGGAAGAGGCATAAATTCGTCAGCCAGTTTGTCTGACC	4949		
Sbjct 91	CGGATAAAATGCTTGATGGTCGGAAGAGGCATAAATTCGTCAGCCAGTTTGTCTGACC	32		
Query 4950	ATCTCATCTGTAAACATCATTGGCAACGCTAC	4980		
Sbjct 31	ATCTCATCTGTAAACATCATTGGCAACGCTAC	1		

[Download](#) ▼[Graphics](#) [SRA](#)**SRX7730879**Sequence ID: [SRA:SRR11092064.2917500.1](#) Length: **151** Number of Matches: **1**Range 1: 1 to 151 [Graphics](#)[▼ Next Match](#) [▲ Previous Match](#)

Score	Expect	Identities	Gaps	Strand
279 bits(151)	2e-72	151/151(100%)	0/151(0%)	Plus/Minus
Query 3319	CCCCGTTTCAGCCCGACCGCTGCGCCTTATCCGGTAACATATCGTCTTGAGTCCAACCCGGT	3378		
Sbjct 151	CCCCGTTTCAGCCCGACCGCTGCGCCTTATCCGGTAACATATCGTCTTGAGTCCAACCCGGT	92		
Query 3379	AAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTA	3438		
Sbjct 91	AAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTA	32		
Query 3439	TGTAGGCGGTGCTACAGAGTTCTTGAAGTGG	3469		
Sbjct 31	TGTAGGCGGTGCTACAGAGTTCTTGAAGTGG	1		

[Download](#) ▼[Graphics](#) [SRA](#)**SRX7730879**Sequence ID: [SRA:SRR11092064.2878891.2](#) Length: **151** Number of Matches: **1**Range 1: 1 to 151 [Graphics](#)[▼ Next Match](#) [▲ Previous Match](#)

Score	Expect	Identities	Gaps	Strand
279 bits(151)	2e-72	151/151(100%)	0/151(0%)	Plus/Plus
Query 3059	CATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGA	3118		
Sbjct 1	CATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGA	60		
Query 3119	AACCCGACAGGACTATAAAGATACAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCCTCT	3178		
Sbjct 61	AACCCGACAGGACTATAAAGATACAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCCTCT	120		
Query 3179	CCTGTTCCGACCCTGCCGCTTACCGGATACC	3209		
Sbjct 121	CCTGTTCCGACCCTGCCGCTTACCGGATACC	151		

Specimen 2

[https://www.ncbi.nlm.nih.gov/sra/SRX7730880\[accn\]](https://www.ncbi.nlm.nih.gov/sra/SRX7730880[accn])

SRX7730880: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid

1 ILLUMINA (Illumina HiSeq 3000) run: 67.1M spots, 20.1G bases, 12.6Gb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the MGIEasy RNA Library Prep Set (96 RXN) (Cat. No.: 1000006384). Paired-end (150 bp) sequencing of the RNA library was performed on the MGISEQ-2000RS platform .

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads

[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)

[hide Abstract](#)

Discovery and characterization of a novel human coronavirus from five patients at the early stage of the Wuhan seafood market pneumonia virus outbreak .

Sample:

[SAMN14082196](#) • [SRS6151291](#) • [All experiments](#) • [All runs](#)

Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:

Name: WIV02-2

Instrument: Illumina HiSeq 3000

Strategy: RNA-Seq

Source: METAGENOMIC

Selection: RANDOM

Layout: PAIRED

Runs: 1 run, 67.1M spots, 20.1G bases, [12.6Gb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092063	67,083,195	20.1G	12.6Gb	2020-02-16

ID: 10108893

BLAST [®] » blastn suite-SRA » results for RID-SBCKMVDN01R

[Home](#)
[Recent Results](#)
[Saved Strategies](#)
[Help](#)

[< Edit Search](#)
[Save Search](#)
[Search Summary](#)

[How to read this report?](#)
[BLAST Help Videos](#)
[Back to Traditional Results Page](#)

Job Title

gb|AY862402.1|

RID

SBCKMVDN01R Search expires on 10-14 21:58 pm [Download All](#) [▼]

Program

BLASTN [?] [Citation](#) [▼]

Database

SRA [See details](#) [▼]

Query ID

AY862402.1

Description

Expression vector pShuttle-SN, complete sequence

Molecule type

nucleic acid

Query Length

5607

Other reports

[Distance tree of results](#)
[MSA viewer](#) [?]

Descriptions

Graphic Summary

Alignments

Sequences producing significant alignments

Download [▼]


Manage Columns [▼]

Show [?]

☒ select all 100 sequences selected

[Graphics](#)
[Distance tree of results](#)

	Description	Max Score	Total Score	Query Cover	E value	Per. Ident	Accession
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.66604450.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.66455076.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.63120099.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.63120099.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.62730385.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.61105639.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.60748776.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.60011402.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.59155252.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.59155252.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.58125883.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.57571550.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.57484454.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.56079039.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.56036194.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.55663455.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.55111993.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.53777284.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.53579813.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.52965281.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.51414706.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.51016881.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.50609371.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.50609371.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.49509270.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.47264810.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.45883858.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.45044544.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.42931446.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.42931446.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.41645159.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.38667194.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.38582435.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.36234753.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.34989436.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.34431070.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.32662604.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.32578887.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.32257474.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.32257474.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.31163438.2
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.30919310.1
<input checked="" type="checkbox"/>	SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.29731694.1



The above distribution of hits appears to ‘invade’ the antigenic, Spike Protein region of the vaccine, residues 961 to 2507. To determine if this was the case, the hit that contained part of the antigen section was displayed (below).

SRX7730880
Sequence ID: [SRA:SRR11092063.55111993.2](#) Length: 150 Number of Matches: 1

Range 1: 1 to 150 [Graphics](#) [Next Match](#)

Score	Expect	Identities	Gaps	Strand
270 bits(146)	3e-68	149/150(99%)	1/150(0%)	Plus/Plus
Query 2471	GTTTAAA-CCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGC	2529		
Sbjct 1	GTTTAAACCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGC	60		
Query 2530	CCCTCCCCGTCCTTCTTGACCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAA	2589		
Sbjct 61	CCCTCCCCGTCCTTCTTGACCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAA	120		
Query 2590	AATGAGGAAATTGCATCGCATTGTCTGAGT	2619		
Sbjct 121	AATGAGGAAATTGCATCGCATTGTCTGAGT	150		

[Download](#) [Graphics](#) [SRA](#)

SRX7730880
Sequence ID: [SRA:SRR11092063.54767346.1](#) Length: 150 Number of Matches: 1

Range 1: 2 to 150 [Graphics](#) [Next Match](#)

Score	Expect	Identities	Gaps	Strand
270 bits(146)	3e-68	148/149(99%)	0/149(0%)	Plus/Plus
Query 2478	CCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCTTCCCC	2537		
Sbjct 2	CCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCTTCCCC	61		
Query 2538	CGTGCTTCTTGCCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAAAATGAGGA	2597		
Sbjct 62	CGTGCTTCTTGCCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAAAATGAGGA	121		
Query 2598	AATTGCATCGCATTGTCTGAGTAGGTGTC	2626		
Sbjct 122	AATTGCATCGCATTGTCTGAGTAGGTGTC	150		

As you can see, this 150 nt sequence starts at 2471 and within the antigen segment. However, there is no homology identified when this is blasted against the Reference Sequence of SARS-CoV-2.

Sample 3

SRX7730881: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid
1 ILLUMINA (Illumina HiSeq 1000) run: 61.3M spots, 18.4G bases, 11.4Gb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the MGIEasy RNA Library Prep Set (96 RXN) (Cat. No.: 1000006384). Paired-end (150 bp) sequencing of the RNA library was performed on the MGISEQ-2000RS platform.

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads
[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)

[hide Abstract](#)
Discovery and characterization of a novel human coronavirus from five patients at the early stage of the Wuhan seafood market pneumonia virus outbreak.

Sample:
[SAMN14082197](#) • SRS6151292 • [All experiments](#) • [All runs](#)
Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:
Name: WIV04-2
Instrument: Illumina HiSeq 1000
Strategy: RNA-Seq
Source: METAGENOMIC
Selection: RANDOM
Layout: PAIRED

Runs: 1 run, 61.3M spots, 18.4G bases, [11.4Gb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092062	61,304,030	18.4G	11.4Gb	2020-02-16

ID: 10108894

Sample 4

SRX7730882: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid

1 ILLUMINA (Illumina HiSeq 3000) run: 34.3M spots, 10.3G bases, 6.4Gb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the MGIEasy RNA Library Prep Set (96 RXN) (Cat. No.: 1000006384). Paired-end (150 bp) sequencing of the RNA library was performed on the MGISEQ-2000RS platform .

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads

[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)

[hide Abstract](#)

Discovery and characterization of a novel human coronavirus from five patients at the early stage of the Wuhan seafood market pneumonia virus outbreak .

Sample:

[SAMN14082198](#) • SRS6151293 • [All experiments](#) • [All runs](#)

Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:

Name: WIV05

Instrument: Illumina HiSeq 3000

Strategy: RNA-Seq

Source: METAGENOMIC

Selection: RANDOM

Layout: PAIRED

Runs: 1 run, 34.3M spots, 10.3G bases, [6.4Gb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092061	34,255,843	10.3G	6.4Gb	2020-02-16

ID: 10108895

[https://www.ncbi.nlm.nih.gov/sra/SRX2913157\[accn\]](https://www.ncbi.nlm.nih.gov/sra/SRX2913157[accn])

Institute of Pathogen Biology, Chinese Academy of Medical Sciences & Peking Union Medical College

above has a few 125 nt hits between about 1950 to 3500 in adenovirus

Sequences used for the blast analyses

Adenovirus vaccine with CoV-1 Spike Protein

1 taactataac ggtcctaagg tagcgaaagc tcagatctgg atctcccgat cccctatggt
 61 cgactctcag tacaatctgc tctgatccg catagttaag ccagtatctg ctcctgctt
 121 gtgtgttga ggtcgtgag tagtgcgca gcaaaattta agctacaaca aggcaaggt
 181 tgaccgacaa ttgatgaag aatctgctta gggtaggcg tttgcgctg cttgcgatg
 241 tacgggccag atatacgctg tgacattgat tattgactag ttattaatag taatcaatta
 301 cggggtcatt agttcatagc ccatatatgg agttccgctg tacataactt acggtaaatg
 361 gcccgctgg ctgaccgccc aacgacccc gccattgac gtcaataatg acgtatgttc
 421 ccatagtaac gccaataggg acttccatt gacgtcaatg ggtggactat ttacggtaaa
 481 ctgccactt ggcagtacat caagtgtatc atatgccaag tacgcccct attgacgtca
 541 atgacggtaa atggcccgcc tggcattatg cccagtacat gaccttatgg gactttcta
 601 cttggcagta catctacgta ttagtcacg ctattacat ggtgatgcg tttggcagt
 661 acatcaatgg gcgtgtagat cggtttgact cacggggatt tccaagtct caccctatg
 721 acgtcaatgg gagttgttt tggcaccaaa atcaacggga cttccaaaa tgcgtaaca
 781 actccgcccc attgacgcaa atgggcggtta ggcgtgtacg gtgggaggtc tatataagca
 841 gagctctctg gctaactaga gaaccactg ctactggct tatcgaaatt aatacgactc
 901 actataggga gaccaagct ggctagcgtt taaacgggcc ctctagagtt gtggtttcaa
 961 gtgatattct tgtaataac taaacgaaca tgtttatttt cttattattt cttactctca
 1021 ctagtggtag tgacctgac cgggtgcacca ctttgatga tgtcaagct cetaattaca
 1081 ctcaacatac ttcactatg agggggggtt actatctga tgaaatttt agatcagaca
 1141 ctctttattt aactcaggat ttattcttc cattttatc taatgttaca gggttcata
 1201 ctattaatca tacgttgac aacctgtca taccttttaa ggatggtatt tatttgctg
 1261 ccacagagaa atcaaatgtt gtcctggtt gggttttgg ttctaccatg aacaacaagt
 1321 cacagtcggt gattattatt aacaattcta ctaatgtgt tatacagaca tgtaactttg
 1381 aattgtgtga caacctttc ttgtctgtt ctaaaccat gggtagacag acacatacta
 1441 tgatattcga taatgcattt aattgcactt tcgagtacat atctgatgcc tttcgcttg
 1501 atgtttcaga aaagtcaggt aattttaaac acttacgaga gttgtgttt aaaaataaag
 1561 atgggtttct ctatgtttat aagggtatc aacctataga tgtagttcgt gatctacctt
 1621 ctgggtttta cactttgaaa cctattttta agttgcctct tggattaaac attacaaatt
 1681 tttagacctt tttacagcc tttcacctg cgcaagacac ttggggcacg tcagctgcag
 1741 cctattttgt tggctattta aagccaacta catttatgct caagtatgat gaaaatggt
 1801 caatcacaga tgctgttgat tgttctcaa atccactgc tgaactcaa tgctctgta
 1861 agagctttga gattgacaaa ggaatttacc agacctctaa ttcagggtt gtccctcag
 1921 gagatgttgt gagattccct aatattacaa acttgtgtcc ttttgagag gttttaatg
 1981 ctactaaatt ccctctgtc tatgcatggg agggaaaaaa aatttctaat tgtgttgctg
 2041 attactctgt gctetacaac tcaacattt ttcaacctt taagtctat ggcgtttctg
 2101 ccactaagtt gaatgatctt tgettctca atgtctatgc agattcttt gtagcaagg
 2161 gagatgatgt aagacaaata gcgccaggac aaactggtgt tattgctgat tataattata
 2221 aattgccaga tgattcatg ggtgtgtcc ttgctggaa tactaggaac attgatgcta
 2281 ctccaactgg taattataat tataaatata ggtatcttag acatggcaag cttaggccct
 2341 ttgagagaga catatcta atgtgcctttt cccctgatgg caaaccttgc accccactg
 2401 ctcttaattg ttattggcca ttaaatgatt atggtttta caccactact ggcattggta
 2461 ccaagcttaa gtttaaccg ctgacagcc tcgactgtgc cttctagtgc ccagccatct

2521 gttgtttgcc cctccccctg gccttccttg accctggaag gtgccactcc cactgtcett
 2581 tcctaataaa atgaggaaat tgcacgcat tgtctgagta ggtgtcattc tattctgggg
 2641 ggtgggggtgg ggcaggacag caagggggag gattgggaag acaatagcag gcatgctggg
 2701 gatgcggtgg gctctatggc ttctgaggcg gaaagaacca gcagatctgc agatctgaat
 2761 tcactatgt cgggtgcgga gaaagaggtg atgaaatggc attatgggta ttatgggtct
 2821 gcattaatga atcgccaac gcgcggggag aggcgggttg cgtattgggc gctctccgc
 2881 ttctcgtc actgactcgc tgcgtcggc cgttcggctg cggcgagcgg taccagctca
 2941 ctcaaaggcg gtaatacggc tatccacaga atcaggggat aacgcaggaa agaacatgtg
 3001 agcaaaaggc cagcaaaagg ccaggaaccg taaaaaggcc gcgttgctgg cgttttcca
 3061 taggtccgc cccctgacg agcatcaca aaatcgacgc tcaagtcaga ggtggcgaaa
 3121 cccgacagga ctataaagat accaggcgtt tccccctgga agctccctcg tgcgtctcc
 3181 tgttcgacc ctgccgtta ccggatacct gtccgccttt cctccctcgg gaagcgtggc
 3241 gctttctcaa tgcacgct gtaggtatct cagttcgtg taggtcgtc gctccaagct
 3301 gggtgtgtg cacgaacccc ccgttcagcc cgaccgtgc gccttatccg gtaactatcg
 3361 tcttgagtc aacccggtaa gacacgactt atcgccactg gcagcagcca ctgtaacag
 3421 gattagcaga gcgaggtatg taggcgggtg tacagagttc ttgaagtggg ggcctaacta
 3481 cggctacact agaaggacag tatttggtat ctgcgtctg ctgaagccag ttacctcgg
 3541 aaaaagagtt gtagctctt gatccggcaa acaaacacc gctggtagcg gtggttttt
 3601 tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc cttgatctt
 3661 ttctacgggg tctgacgctc agtgaacga aaactcacgt taagggaatt tggcatgag
 3721 attatcaaaa aggatcttca cctagatcct ttgacctc cggcgttcag cctgtgccac
 3781 agccgacagg atggtgacca ccattgccc catatcaccg tcggtactga tccgctcgc
 3841 aataaaccca accgtacac cctgagcatc aaactcttt atcagttgga tcatgtcggc
 3901 ggtgtcggc ccaagacggt cgagcttctt caccagaatg acatcacctt cctccacct
 3961 catcctcagc aaatccagcc ctcccgatc tgtgaaactg ccggatgcct tgcggtaaa
 4021 gatgcggtta gctttacc ctcacatctt gagcgtgag gtctgcctcg tgaagaaggt
 4081 gttgtgact cataccaggc ctgaatgcc ccatcatcca gccagaaagt gagggagcca
 4141 cgggtgatga gagcttgtt gtaggtggac cagttggtga tttgaaactt ttgctttgcc
 4201 acggaacggt ctgcgttgc gggaagatgc gtgatctgat cctcaactc agcaaaagt
 4261 cgatttattc acaaaagccg ccgtcccgtc aagtcagcgt aatgctctgc cagtgttaca
 4321 accaattaac caattctgat tagaaaaact catcgagcat caaatgaaac tgcaatttat
 4381 tcatacagg attatcaata ccatatttt gaaaaagccg ttctgtaat gaaggagaaa
 4441 actcaccgag gcagttccat aggatggcaa gatcctggtg tcggtctgcg attccgactc
 4501 gtccaacatc aatacaacct attaatctc cctcgtcaaa aataagggtg tcaagtgaga
 4561 aatcaccatg agtgacgact gaatccggtg agaatggcaa aagcttatgc attctttcc
 4621 agactgttc aacaggccag ccattacgtc cgtcatcaaa atcactcga tcaaccaaac
 4681 cgttattcat tctgattgc gcctgagcga gacgaaatac gcgatcgtg taaaaggac
 4741 aattacaaac aggaatcgaa tgcaaccggc gcaggaacac tgccagcga tcaacaatat
 4801 ttacactga atcaggatat tcttcaata cctggaatgc tgtttccc gggatcgcag
 4861 tggtagtaa ccatgcatca tcaggagtag ggataaatg cttgatggtc ggaagaggca
 4921 taaattccgt cagccagttt agtctgacca tctcatctgt aacatcattg gcaacgctac
 4981 ctttgccatg ttcagaaac aactctggcg catcgggctt cccatacaat cगतगattg
 5041 tgcacctga ttgcccgaca ttatcgag cccatttata cccatataaa tcagcatcca
 5101 tgttgaatt taatcgcggc ctcgagcaag acgtttccc tgaatatgg ctcataacac
 5161 ccttgatt actgttatg taagcagaca gttttattgt tcatgatgat atattttat
 5221 cttgtgcaat gtaacatcag agattttgag acacaacgtg gctttgttga ataatcgaa

5281 cttttgctga gttgaaggat cagatcacgc atcttcccgga caacgcagac cgttccgtgg
 5341 caaagcaaaa gttcaaaatc accaactggt ccacctacaa caaagctctc atcaaccgtg
 5401 gctccctcac ttctggctg gatgatgggg cgattcaggc ctggtatgag tcagcaacac
 5461 cttcttcacg aggcagacct cagcgtaga ttattgaagc atttatcagg gttattgtct
 5521 catgagcgga tacatatattg aatgtattta gaaaaataaa caaatagggg ttccgcgcac
 5581 atttcccgga aaagtccac ctgacgt

SARS-CoV-2 Spike Protein gene

atgtttgt ttttctgtt ttattgccac tagtctctag
 21601 tcagtgtgtt aatcttaca ccagaactca attaceccct gcatacacta attcttca
 21661 acgtgggtgt tattaccctg acaaagttt cagatctca gttttacatt caactcagga
 21721 cttgttcta ctttctttt ccaatgttac ttgggtccat gctatacatg tctctgggac
 21781 caatgggtact aagaggttg ataaccctgt cctaccattt aatgatggtg tttatttgc
 21841 ttccactgag aagtctaaca taataagagg ctggattttt ggtactactt tagattcgaa
 21901 gacccagtc ctactattg ttaataacgc tactaatgtt gttattaaag tctgtgaatt
 21961 tcaattttgt aatgatccat tttgggtgt ttattaccac aaaaacaaca aaagtggat
 22021 ggaaagttag ttcagagttt attctagtgc gaataattgc acttttgaat atgtctctca
 22081 gccttttctt atggacctg aaggaaaaca gggtaatttc aaaaatctta gggaattgt
 22141 gtttaagaat attgatggtt atttataat atattctaag cacacgcta ttaattagt
 22201 gcgtgatctc cctcagggtt ttccggttt agaaccattg gtagatttgc caataggtat
 22261 taacatcact aggtttcaaa ctttacttgc ttacataga agttattga ctcctggtga
 22321 ttcttctca gtttgacag ctggtgctgc agcttattat gtgggttate tcaacctag
 22381 gacttttcta ttaaaatata atgaaatgg aaccattaca gatgctgtag actgtgact
 22441 tgacctctc tcagaaacaa agtgtacgtt gaaatcctc actgtagaaa aaggaatcta
 22501 tcaaaactct aactttagag tcaaccaac agaactctatt gtagatttc ctaatattac
 22561 aaactgtgc cttttggtg aagttttta cgcaccaga ttgcatctg tttatgctg
 22621 gaacaggaag agaatcagca actgtgttgc tgattattct gtctatata attcgcac
 22681 atttccact ttaagtgtt atggagtgc tctactaaa ttaaatgate tctgcttac
 22741 taatgtctat gcagattcat ttgtaattag aggtgatgaa gtcagacaaa tegtccagg
 22801 gcaaaactga aagattgctg attataatta taaattacca gatgatttca caggctgct
 22861 tatagcttgg aattctaaca atcttgatc taagggttgg gtaattata attacctga
 22921 tagattgtt aggaagtcta atctcaaacc tttgagaga gatatttcaa ctgaaatcta
 22981 tcaggccggt agcacacct gtaatggtgt tgaaggttt aattgttact ttctttaca
 23041 atcatatggt ttcaaccca ctaatggtgt tggttacca ccatcacag tagtagtact
 23101 ttcttttga cttctacatg caccagcaac tgtttgtga cctaaaaagt ctactaatt
 23161 ggttaaaaac aatgtgtca atttcaact caatggttca acaggcacag gtgttctac
 23221 tgagtctaac aaaaagtgc tgctttcca acaatttggc agagacattg ctgacactac
 23281 tgatgctgct cgtgatecac agacactga gattcttgc attacacat gttctttgg
 23341 tgggtgcagt gttataacac caggacaaa tacttctaac caggttgcgt ttctttatca
 23401 ggatgttaac tgcacagaag tccctgttgc tattcatgca gatcaactta ctcctactg
 23461 gcgtgtttat tctacagggt ctaatgttt tcaaacacgt gcaggctgtt taataggggc
 23521 tgaacatgct acaactcat atgagtgtga cataccatt ggtgcaggta tatgcgctag
 23581 ttatcagact cagactaatt ctctcggcg ggcacgtagt gtagctagtc aatccatcat
 23641 tgcttacct atgtcactg gtgcagaaaa ttcagttgct tactctaata actctattgc
 23701 cataccaca aattttacta ttagtgtac cacagaaatt ctaccagtgt ctatgaccaa

23761 gacatcagta gattgtacaa tgtacatttg tggtagtca actgaatgca gcaatcttt
23821 gttgcaatat ggcagtttt gtacacaatt aaaccgtgct ttaactggaa tagctgtga
23881 acaagacaaa aacaccaag aagttttgc acaagtcaaa caaatttaca aaacaccacc
23941 aattaaagat ttggtgggt ttaattttc acaaataa ccagatccat caaaaccaag
24001 caagaggtca ttattgaag atctacttt caacaaagt acactgcag atgctggctt
24061 catcaaaaa tatggtgatt gccttggtga tattgctgct agagacctca ttgtgcaca
24121 aaagttaac ggccttactg tttgccacc ttgtgcaca gatgaaatga ttgtcaata
24181 cactctgca ctgttagcgg gtacaatcac ttctggttg accttgggtg caggtgctgc
24241 attacaaata ccattgcta tgcaaatggc ttatagggtt aatggtattg gagttacaca
24301 gaatgtctc tatgagaacc aaaaattgat tgccaaccaa ttaaatagt ctattggcaa
24361 aattcaagac tcactttct ccacagcaag tgcacttga aaactcaag atgtggtaa
24421 ccaaatgca caagcttaa acacgcttg taaacaact agtccaatt ttgtgcaat
24481 ttcaagtgt ttaaatgata tctttcacg tctgacaaa gttgaggtg aagtgcaat
24541 tgataggtg atcacaggca gacttcaaag ttgcagaca tatgtgact aacaattaat
24601 tagagctgca gaaatcagag cttctgcta tctgtgct actaaatgt cagagtgtg
24661 acttgacaa tcaaaaagag ttgattttg tggaagggc tcatctta tgcctccc
24721 tcagtcagca cctcatggtg tagtctctt gcatgtgact tatgtcctg cacaagaaaa
24781 gaacttaca actgtcctg ccatttgta tgatggaaaa gcacacttc ctgtgaagg
24841 tgtctttgt tcaaatgca cacttggt tgtaacaaa aggaatttt atgaaccaca
24901 aatcattact acagacaaca cattgtgtc tggtactgt gatgtgtaa taggaattg
24961 caacaacaca gtttatgac ctttgaacc tgaattagac tcattcaagg aggagttaga
25021 taaatattt aagaatcata catcaccaga tttgattta ggtgacatc ctggcattaa
25081 tgettcagt gtaaacattc aaaaagaaat tgaccgctc aatgaggtg ccaagaattt
25141 aatgaatct ctcategac tccaagaact tggaagtat gagcagtata taaaatggcc
25201 atggtacatt tggctaggt ttatagctg ctgtattgcc atagtaattg tgacaattat
25261 gctttgctg atgaccagt gctgtagtg tctcaagggc tttgttct gtggatcctg
25321 ctgcaattt gatgaagac actctgagcc agtctcaaa ggagtcaat tacattacac
25381 ataa

***In silico* construct with Adenovirus vector shuttle containing CoV-2 Spike Protein gene**

```

1 taactataac ggtcctaagg tagcgaaagc tcagatctgg atctcccgat cccctatggt
61 cgactctcag tacaatctgc tctgatccg catagttaag ccagtatctg ctccctgctt
121 gtgtgttga ggtcgtgag tagtgcgca gcaaaattta agctacaaca aggcaaggct
181 tgaccgaaa ttgcatgaag aatctgctta gggtaggcg tttgcgctg ctgcgcatg
241 tacgggccag atatacgctg tgacattgat tattgactag ttattaatag taatcaatta
301 cggggtcatt agttcatagc ccatatatgg agttccgctg tacataactt acggtaaagt
361 gcccgctgg ctgaccgcc aacgacccc gccattgac gtcaataatg acgtatgttc
421 ccatagtaac gccaataggg actttccatt gacgtcaatg ggtggactat ttacggtaaa
481 ctgccactt ggcatgacat caagtgtatc atatgccaag tacgccccct attgacgtca
541 atgacggtaa atggcccgcc tggcattatg cccagtacat gacctatgg gactttcta
601 cttggcagta catctacgta ttatcatcgt ctattacat ggtgatgcg tttggcagt
661 acatcaatgg gctgtgatag cgtttgact cagggggatt tccaagtct caccacattg
721 acgtcaatgg gagtttgtt tggcacaaa atcaacggga cttccaaaa tgcgtaaca
781 actccgccc attgacgcaa atggcggtg ggcgtgtac gtgggaggtc tatataagca
841 gagctctctg gtaactaga gaaccactg cttactggt tategaaatt aatacgactc
901 actataggga gacccaagct ggctagcgt taaacgggc ctctagagt gtggttcaa
961 gtgatattct tgtaataac taaacgaaca tgtttgtt tctgttta tgccactag tctctag
21601 tcagtgtgtt aattctacaa ccagaactca attaccctt gcatacacta attcttca
21661 acgtgggtgt tattaccctg acaaagttt cagatctca gttttacatt caactcagga
21721 cttgttcta cttttttt ccaatgttac ttggtccat gctatacatg tctctgggac
21781 caatggfact aagaggttg ataaccctgt cctaccattt aatgatggtg tttatttgc
21841 ttccactgag aagtctaaca taataagagg ctggatttt ggtactact tagattcgaa
21901 gaccagtc ctactattg ttaataacgc tactaatgt gttattaaag tctgtgaatt
21961 tcaattttgt aatgatecat tttgggtgt ttattaccac aaaaacaaca aaagtggat
22021 ggaaagttag ttcagagttt attctagtgc gaataattgc acttttgaat atgtctctca
22081 gcctttctt atggacctg aaggaaaaca gggtaattc aaaaatctta gggaattgt
22141 gttaagaat attgatggtt attttaaaat atattctaag cacacgcta ttaattagt
22201 gcgtgatctc cctcagggtt ttccgcttt agaaccattg gtagatttgc caataggtat
22261 taacatcact aggttcaaaa cttacttgc ttacataga agttattga ctctggtga
22321 ttcttctca gtttgacag ctggtgctgc agcttattat gtgggttate ttaacctag
22381 gacttttcta taaaatata atgaaatgg aaccattaca gatgctgtag actgtgcact
22441 tgacctctc tcagaacaa agtgtagctt gaaatcttc actgtagaaa aaggaatcta
22501 tcaaaactct aacttagag tcaaccaac agaactatt gttagattc ctaattac
22561 aaacttgtgc cttttgtg aagttttta cgcaccaga ttgcatctg ttatgcttg
22621 gaacaggaag agaactagca actgtgtg tgattattct gtcctatata attccgcate
22681 attttcact ttaagtgt atggagtgc tctactaaa ttaatgate tctgcttac
22741 taatgtctat gcagattcat ttgtaattag aggtgatgaa gtcagacaaa tctctcagg
22801 gcaaaactga aagattgctg attataatta taaattacca gatgatttta caggctgcgt
22861 tatagcttgg aattctaaca atcttgatc taaggttggt gtaattata attacgtga
22921 tagattgttt aggaagtcta atctcaaacc tttgagaga gatatttcaa ctgaaatcta
22981 tcaggccggt agcacactt gtaatggtgt tgaaggttt aattgttact ttctttaca
23041 atcatatggt ttcaaccca ctaatggtgt tggftacca ccatacagag tagtagtact
23101 ttctttgaa cttctacatg caccagcaac tgtttgtga cctaaaaagt ctactaatt
23161 ggttaaaaac aaatgtgta atttcaact caatggttta acaggcacag gtgttctac
23221 tgagtetaac aaaaagttc tgcctttcca caaattggc agagacattg ctgacactac

```

23281 tgatgtgtc cgtgatccac agacacttga gattcttgac attacacat gttctttgg
 23341 tgggtgcagt gttataacac caggaacaaa tacttctaac caggttgctg ttctttatca
 23401 ggatgttaac tgcacagaag tcctgttgcc tattcatgca gatcaactta ctctacttg
 23461 gcgtgtttat tctacagggt ctaatgtttt tcaaacacgt gcaggctgtt taataggggc
 23521 tgaacatgtc aacaactcat atgagtgtga cataccattt ggtgcaggtg tatgcgctag
 23581 ttatcagact cagactaatt ctctcggcg ggcacgtagt gtagctagtc aatccatcat
 23641 tgcctacact atgtcacttg gtgcagaaaa ttcaattgct tactctaata actctattgc
 23701 catacccaca aattttacta ttagtgttac cacagaaatt ctaccagtgt ctatgacca
 23761 gacatcagta gattgtacaa tttacatttg tgggtattca actgaatgca gcaatcttt
 23821 gttgcaatat ggcagttttt gtacacaatt aaacctgtgt ttaactggaa tagctgttga
 23881 acaagacaaa aacaccaag aagtttttgc acaagtcaaa caaatttaca aaacaccacc
 23941 aattaaagat tttggtggtt ttaattttc acaaatatta ccagatccat caaaaccaag
 24001 caagagggtca ttattgaag atctactttt caacaaagt acactgcag atgtggtt
 24061 catcaacaa tatggtgatt gccttggtga tattgctgct agagacctca ttgtgcaca
 24121 aaagttaac ggccttactg tttgccacc ttgttcaca gatgaaatga ttgtcaata
 24181 cactctgca ctgttagcgg gtacaatcac ttctggttg acccttggtg caggtgctgc
 24241 attacaaata ccatttgcta tgcacatggc ttatagggtt aatggtattg gagttacaca
 24301 gaatgttctc tatgagaacc aaaaattgat tgccaaccaa ttaaatagtg ctattggcaa
 24361 aatcaagac tcaatttctt ccacagcaag tgcacttga aaactcaag atgtggtcaa
 24421 ccaaatgca caagctttaa acacgttgt taaacaactt agtccaatt ttggtgcaat
 24481 ttcaagtgtt ttaaatgata tcctttcacg tcttgacaaa gttgaggctg aagtgcacat
 24541 tgatagggtg atcacaggca gacttcaaag ttgcagaca tatgtgactc acaattaat
 24601 tagagctgca gaaatcagag ctctgctaa tcttgctgct actaaatgt cagagtgtg
 24661 acttgacaaa tcaaaaagag ttgattttt tggaaagggc tatcatctta tgccttccc
 24721 tcagtcagca cctcatggtg tagtctctt gcattgtgact tatgtccctg cacaagaaaa
 24781 gaacttaca actgctctg ccatttgta tgatggaaaa gcacacttc ctctgaagg
 24841 tgtctttgt tcaaatggca cacactggtt tgtaacaca aggaatttt atgaaccaca
 24901 aatcattact acagacaaca catttggtc ttgtaactgt gatgtgtta taggaattgt
 24961 caacaacaca gtttatgac ctttgcaacc tgaattagac tcattcaagg aggagttaga
 25021 taaatattt aagaatcata catcaccaga tttgattta ggtgacatct ctggcattaa
 25081 tgcctcagtt gtaaacattc aaaaagaaat tgaccgcctc aatgagggtg ccaagaattt
 25141 aatgaatct ctcatgac tccaagaact tggaaagtat gagcagtata taaatggcc
 25201 atggtacatt tggctaggtt ttatagctgg ctgattgcc atagtaattg tgacaattat
 25261 gctttgctgt atgaccagtt gctgtagttg tctcaagggc tttgttctt gtggatcctg
 25321 ctgcaaatgt gatgaagacg actctgagcc agtgctcaaa ggagtcacaa tacattacac
 25381 ataattg ccagccatct
 2521 gttgtttgcc cctccccctg gccttctttg accctggaag gtgccactcc cactgtcctt
 2581 tcctaataaa atgaggaaat tgcacgcgat tttctgagta ggtgtcattc tattctgggg
 2641 ggtgggggtgg ggcaggacag caaggggggag gattgggaag acaatagcag gcattgctggg
 2701 gatgcggtgg gctctatggc ttctgaggcg gaaagaacca gcagatctgc agatctgaat
 2761 tcatctatgt cgggtgctga gaaagaggta atgaaatggc attatgggta ttatgggtct
 2821 gcattaatga atcgcccaac gcgcggggag aggcgggttg cgtattgggc gctcttccgc
 2881 ttctcgtc actgactgc tgcgtcggg cgttcggctg cggcgagcgg tatcagctca
 2941 ctcaaaggcg gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacatgtg
 3001 agcaaaaggc cagcaaaagg ccaggaaccg taaaaaggcc gcgttgctgg cgttttcca
 3061 taggtctcgc cccctgacg agcatcaca aaatcgacgc tcaagtcaga ggtggcgaaa

3121 cccgacagga ctataaagat accaggcgtt tccccctgga agctccctcg tgcgtctcc
 3181 tgttccgacc ctgccgctta cggataacct gtccgccttt ctcccttcgg gaagcgtggc
 3241 gctttctcaa tgctcacgt gtaggtatct cagttcggtg taggtcgttc gctccaagct
 3301 gggctgtgtg cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg
 3361 tcttgagtc aacccggtaa gacacgactt atcgccactg gcagcagcca ctggtaacag
 3421 gattagcaga gcgaggtatg taggcggtgc tacagagttc ttgaagtggg ggcctaacta
 3481 cggctacact agaaggacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg
 3541 aaaaagagtt ggtagctctt gatccggcaa acaaacacc gctggtagcg gtggtttttt
 3601 tgttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt
 3661 ttctacgggg tctgacgtc agtggaaacga aaactcacgt taagggattt tggcatgag
 3721 attatcaaaa aggatcttca cctagatcct ttgatctc cggcgttcag cctgtgccac
 3781 agccgacagg atggtgacca ccatgtgcc catatcaccg tcggtactga tcccgctgc
 3841 aataaacga accgctacac cctgagcatc aaactcttt atcagttgga tcatgtcggc
 3901 ggtgtcggcg ccaagacggt cgagcttctt caccagaatg acatcacctt cctccacctt
 3961 catcctcagc aaatccagcc ctcccgatc tgttgaactg ccggatgcct tgcggtaaa
 4021 gatgcggtta gcttttacc ctgcatcttt gagcgtgag gtctgcctcg tgaagaagg
 4081 gttgctgact cataccaggc ctgaatgcc ccatcatcca gccagaaagt gagggagcca
 4141 cggttgatga gagctttgtt gtaggtggac cagttggtga ttttgaactt ttgctttgcc
 4201 acggaacggt ctgcgtgtc gggaagatgc gtgatctgat cttcaactc agcaaaagt
 4261 cgatttattc acaaaagccg cgtcccgtc aagtcagcgt aatgctctgc cagtgttaca
 4321 accaattaac caattctgat tagaaaaact catcgagcat caaatgaaac tgcaatttat
 4381 tcatacagg attatcaata ccatatttt gaaaaagccg ttctgtaat gaaggagaaa
 4441 actcaccgag gcagttccat aggatggcaa gatcctggta tcggtctgcg attccgactc
 4501 gtccaacatc aatacaacct attaatctc cctcgtaaa aataagggtt tcaagtgaga
 4561 aatcaccatg agtgacgact gaatccggtg agaatggcaa aagcttatgc atttcttcc
 4621 agactgttc aacaggccag ccattacgtc cgtcatcaaa atcactcgca tcaacaaac
 4681 cgttattcat tcgtgattgc gctgagcga gacgaaatac gcgacgctg taaaaggac
 4741 aattacaaac aggaatcgaa tgcaaccggc gcaggaacac tgccagcgca tcaacaatat
 4801 tttcacctga atcaggatat tcttctaata cctggaatgc tgtttcccg gggatcgag
 4861 tggtagtaaa ccatgcatca tcaggagtac ggataaaatg cttgatggc ggaagaggca
 4921 taaatccgt cagccagttt agtctgacca tctcatctgt aacatcattg gcaacgctac
 4981 ctttgccatg ttcagaaac aactctggcg catcgggctt cccatacaat cgatagattg
 5041 tcgcacctga ttgccgaca ttatcgcgag ccatttata cccatataaa tcagcatcca
 5101 tgttggaatt taatcgcggc ctgagcaag acgtttcccg ttgaatatgg ctcataaac
 5161 ccttgtatt actgtttatg taagcagaca gttttattgt tcatgatgat atattttat
 5221 cttgtgcaat gtaacatcag agattttgag acacaacgtg gctttgtga ataaatcgaa
 5281 cttttgctga gttgaaggat cagatcacgc atctcccgca caacgcagac cgtccgtgg
 5341 caaagcaaaa gttcaaaatc accaactggt ccactacaa caaagctctc atcaaccgtg
 5401 gtcctcac tttctggctg gatgatgggg cgattcaggc ctggtatgag tcagcaaac
 5461 cttcttcacg aggcagacct cagcgtaga ttattgaagc atttatcagg gttattgtct
 5521 catgagcgga tacatatttg aatgtattta gaaaaataaa caaatagggg ttccgcgcac
 5581 atttccccga aaagtgccac ctgacgt

Artificial Spike Protein in Chinese patent (not found in any patient specimens)

gaattgccg ccaccatgga cgccatgaag cggggcctct gctgtgttct gctgctctgc 60
 [0013] ggcgccgtgt tctgagtaa ctcgagccag tgcgtgaacc tgacaacaag gacacagctg 120
 [0014] cccctgcct acacaacag cttcactagg ggcgtgtact accccgacaa ggtgttcagg 180
 [0015] tccagcgtgc tgcacagcac acaggacctg ttctgccct tcttcagcaa cgtgacatgg 240
 [0016] ttccacgcca ttcacgtgag cgggaccaac gggaccaagc ggttcgataa cctgtcttg 300
 [0017] ccttcaacg atggcgtgta ctttgccagc accgagaagt ccaacatcat caggggctgg 360
 [0018] atcttggca caacctgga cagcaagacc cagagcctcc tgatcgtcaa caacgccaca 420
 [0019] aacgtcgtga tcaagggtg caggttcag ttctgcaac atccattcct gggcgtgtac 480
 [0020] taccataaga acaacaagtc ctggatggag agcgagtcc gggctactc cagcgccaac 540
 [0021] aactgcacct tcgagtacgt gagccagccc ttctgatgg acttgagggg gaagcagggc 600
 [0022] aactcaaga acctccggga gtctgtctt aagaacattg acggctactt caagatctac 660
 [0023] tccaagcaca ccccatcaa cctcgtcagg gatctgcccc aggggtttag cgccctggag 720
 [0024] cccctggtcg atctgcaat cggcatcaac atcacacggt ttcagaccct gctggccctg 780
 [0025] caccggtcct acctacccc tggcgatagc agctccggt ggacagccgg ggccgcccgc 840
 [0026] tactacgtcg gctacctca gcctcggact ttctgtga agtacaacga gaacgggaca 900
 [0027] ataccgatg ccgtggactg cgcctggat cccctcagcg agaccaagt cactgaag 960
 [0028] tccttactg tggagaagg gatctaccag acatccaact ttagggtgca gccaccgag 1020
 [0029] agcattgca ggtcccca catcacaac ctgtcccc ttggcgaggt gtcaacgcc 1080
 [0030] acaagattcg cttcgtgta cgcctggaac aggaagcggg ttagcaactg cgtggccgat 1140
 [0031] tactcgtcc tgtacaacag cgcctcttc tccacctca agtgctacgg cgtgtcccc 1200
 [0032] accaagctga acgatctgt cttactaac gtgtacgtg acagcttct gatcagaggc 1260
 [0033] gatgaggtgc ggcagatgc ccctgggcag acagggaaga tcgccgacta caactacaag 1320
 [0034] ctgcccgatg acttcacagg gtgcgtgat gcctggaact ccaacaacct ctagcaag 1380
 [0035] gtgggcggca actacaacta cctctacagg ctgttagga agtccaacct gaagccctt 1440
 [0036] gagcgggata ttagcacca gatctaccag gccgggagca cccctgtaa cggcgtcgag 1500
 [0037] gggtttaact gctacttcc tctgcagagc tacgggtcc agccacca cggggtcggg 1560
 [0038] taccagccat accgggtggt ggtgctgagc ttcagctgc tgcagcccc agccaccgtc 1620
 [0039] tgcggccca agaagtcac taactgggtg aagaacaagt gcgtgaactt caactcaac 1680
 [0040] ggctgacag ggacaggcgt gctgacagag tccaacaaga agttctccc ctccagcag 1740
 [0041] ttgggcggg acattgccga cacaaccgat gccgtgcggg acccagacac cctggagatc 1800
 ctggacatca caccctgcag cttcggcggg gtgagcgtga ttacaccgg cacaacacc 1860
 tccaaccagg tggcctgct gtaccaggat gtgaactgca cagaggtccc cgtggccatt 1920
 cagccgatc agctgacccc cactggcgg gtgtacagca ccggctcaa cgtgttcag 1980
 actagggccg gctgcctgat cggggccgag cagtgaca acagctacga gtgcgacatc 2040
 ccattgggg ccgggatctg cgcctctac cagacacaga caaacagccc taggcgggcc 2100
 aggtcgtgg ccagccagtc catcatgcc tacaccatga gcctgggcgc cgagaacagc 2160
 gtggcctaca gcaacaacag catcgtatc ccaacaaact ttaccatctc cgtgaccacc 2220
 gagatcctgc ccgtcagcat gactaagaca tccgtcagc gcaccatgta catctcggg 2280
 gagagcaccg agtgcctcaa cctgctgctg cagtacgggt ccttctgcac ccagctgaac 2340

agggccctga ctggcattgc cgtcgagcag gataagaaca cacaggaggt ctttgcacag 2400
 gtgaagcaga tctacaagac accccaatt aaggacttcg ggggttcaa ctttcccag 2460
 attctgctg acccagcaa gcccgcaag cggctcttca tcgaggacct gctgttcaac 2520
 aaggtgacac tggccgacgc cggctttatc aagcagtacg gcgactgcct cggcgacatc 2580
 gccgctaggg acctgatctg cgcccagaag ttcaacggcc tgacagtgc tccccctctg 2640
 ctgacagacg agatgatcgc ccagtacaca agcgccctgc tggccggcac catcacctcc 2700 ggggtggacat
 tcggggccgg gggcgccctg cagatcccct ttgcatgca gatggcctac 2760 aggttcaacg gcattggcgt gacacagaac
 gtgctgtacg agaaccagaa gctgatgcgc 2820 aaccagtta actccgccat cgggaagatc caggattccc tgagcagcac
 cgccagcgc 2880 ctgggcaagc tccaggatgt ggtgaaccag aacgcccagg cctcaacac cctggtgaag 2940
 cagctgtcct ccaacttcgg cgccattagc tccgtgctga acgacatcct gagccggctg 3000 gacaagggtg aggccgaggt
 gcagattgac cggctgatta ccggacggct gcagtcctg 3060 cagacctacg tgacacagca gctcatccgg gccgccgaga
 tccgcgctc cgccaacctg 3120 gccgccacta agatgtccga gtgcgtgctc ggccagagca agagggtgga ttctgcggg
 3180 aagggtacc acctgatgag ctccccag agcgcccc atgggggtgt gttcctgcac 3240 gtgacatacg
 tgctgcccga ggagaagaac taccaccg cccagccat ttgccacgac 3300 ggcaaggccc acttccctag ggagggcggt
 ttcgtgacg acgggacaca ctggttcgtg 3360 acccagcgga acttctacga gcccagatt atcaccagc ataaccctt
 tgttccggg 3420 aactgcgatg tctgattgg gatcgtcaac aacacagtct acgacccct gcagcccag 3480
 ctgcatagct ttaaggagga gctggataag tactttaaga accacacctc cctgatgtg 3540 gacctggggg atatcagcgg
 catcaacgcc agcgtggtga acatccagaa ggagatgat 3600 aggtgaacg aggtggccaa gaacctgaac
 gagtcctga tcgacctgca ggagctggg 3660 aagtacgagc agtacatcaa gtggccctgg tacatctggc tgggcttcat
 cgccgggctg 3720 atcgccatcg tgatggtgac cattatgctc tctgcatga ctactgctg ctctgctg 3780
 aagggtgct gcagctcgg gagctgctgc aagttgatg aggatgatag cgagccagt 3840 ctgaaggcgc tgaagctgca
 ctacacctg aagctt

Adenovirus 5 vector shuttle with Synthetic construct H7N9 HA gene 7640-9302

1 taactataac ggtcctaagg tagcgaaagc tcagatctgg atctcccgat ccctatggt
 61 cgactctcag tacaatctgc tctgatgccg catagttaag ccagtatctg ctccctgctt
 121 gtgtgttga ggtcgtgag tagtgcgca gcaaaatta agctacaaca aggcaaggct
 181 tgaccgaaa ttgatgaag aatctgctta gggtaggcg tttgcgctg ctccgcatg
 241 tacgggccag atatacgcgt tgacattgat tattgactag ttattaatag taatcaatta
 301 cggggtcatt agttcatagc ccatatatgg agttccgctg tacataact acggtaaatg
 361 gccgcctgg ctgaccgccc aacgaccccc gccattgac gtcaataatg acgtatgtt
 421 ccatagtaac gccaataggg acttccatt gacgtcaatg ggtggactat ttacggtaaa
 481 ctgcccactt ggcagtacat caagtgtatc atatgccaag tacgcccct attgacgtca
 541 atgacggtaa atggccccgc tggcattatg ccagttacat gaccttatgg gactttcta
 601 cttggcagta catctacgta ttagtcacg ctattacat ggtgatgcgg tttggcagt
 661 acatcaatgg gcgtggatag cggttgact cacggggatt tccaagtct caccctatg
 721 acgtcaatgg gagttgtt ttggacaaa atcaacggga cttccaaaa tgcgtaaca
 781 actccgccc attgacgcaa atggcggtga ggcgtgtac gtgggaggtc tatataagca
 841 gagctctctg gtaactaga gaaccactg ctactggct tatcgaatt aatagcact
 901 actataggga gaccaagct ggctagcgt taaacgggc ctctagagt gtggttcaa
 961 gtgatattct tgtaataac taaacgaac

7681 tetgcctcgg acatcatgcc gtgtcaaacg gaaccaaagt aaacacatta actgaaagag
7741 gagtggaaagt cgtcaatgca actgaaacag tggaacgaac aaacatcccc aggatctgct
7801 caaaagggaa aaggacagtt gacctcggtc aatgtggact cctggggaca atcactggac
7861 cacctcaatg tgaccaatc ctagaatctt cagccgattt aattattgag aggcgagaag
7921 gaagtgatgt ctgttatcct gggaaattcg tgaatgaaga agctctgagg caaattctca
7981 gagaatcagg cggaattgac aaggaagcaa tgggattcac atacagtgga ataagaacta
8041 atggagcaac cagtgcattg aggagatcag gatcttcatt ctatgcagaa atgaaatggc
8101 tcctgtcaaa cacagataat gctgcattcc cgcagatgac taagtcatat aaaaatacaa
8161 gaaaaagccc agctctaata gtatggggga tccatcattc cgtatcaact gcagagcaaa
8221 ccaagctata tgggagtgga acaaaactgg tgacagtggg gagttctaata tatcaacaat
8281 cttttgtacc gagtccagga gcgagaccac aagttaatgg tctatctgga agaattgact
8341 ttcattggct aatgctaaat cccaatgata cagtcacttt cagtttcaat ggggctttca
8401 tagctccaga cctgcaagc ttctgagag gaaaatctat gggaatccag agtggagtac
8461 aggttgatgc caatttgaa ggggactgct atcatagtgg agggacaata ataagtaact
8521 tgccatttca gaacatagat agcagggcag ttggaaaatg tccgagatat gtaagcaaa
8581 ggagtctgct gctagcaaca gggatgaaga atgttctga gattccaaaa ggaagaggcc
8641 tatttggtgc tatagcgggt ttattgaaa atggatggga aggcctaatt gatggttggt
8701 atggtttcag acaccagaat gcacaggag agggaaactgc tgcagattac aaaagcactc
8761 aatcggcaat tgatcaaata acaggaaaat taaaccggct tatagaaaaa accaaccaac
8821 aatttgagtt gatagacaat gaattcaatg aggtagagaa gcaaatcggg aatgtgataa
8881 attggaccag agattctata acagaagtgt ggtcatacaa tgctgaactc ttggtagcaa
8941 tggagaacca gcatacaatt gatctggctg attcagaaat ggacaaactg tacgaacgag
9001 tgaaaagaca gctgagagag aatgctgaag aagatggcac tggttgcttt gaaatatttc
9061 acaagtgtga tgatgactgt atggccagta ttagaaataa cacctatgat cacagcaaat
9121 acaggggaaga ggcaatgcaa aatagaatac agattgacc agtcaaacta agcagcggct
9181 acaaagatgt gatactttgg ttagcttcg gggcatcatg ttcatactt ctagecattg
9241 taatgggcct tgtcttcata tgtgtaaaga atggaacat gcggtgcact attgtatat
9301 aattg ccagcatct

2521 gttgtttgcc cctccccctg gccttccttg accctggaag gtgccactcc cactgtcctt
2581 tcctaataaa atgaggaaat tgcategc atgtctgagta ggtgtcattc tattctgggg
2641 ggtgggggtgg ggcaggacag caagggggag gattgggaag acaatagcag gcatgctggg
2701 gatgcgggtgg gctctatggc ttctgaggcg gaaagaacca gcagatctgc agatctgaat
2761 tcatttatgt cgggtgcgga gaaagaggta atgaaatggc attatgggta ttatgggtct
2821 gcattaatga atcgcccaac gcgcggggag aggcgggttg cgtattgggc gctctccgc
2881 ttctcgtc actgactgc tgcgtcggc cgttcggctg cggcgagcgg tatcagctca
2941 ctcaaaggcg gtaatacggc tatccacaga atcaggggat aacgcaggaa agaactatg
3001 agcaaaaggc cagcaaaagg ccaggaaccg taaaaggcc gcgttgctgg cgttttcca
3061 taggctccgc cccctgacg agcatcaca aaatcgacgc tcaagtcaga ggtggcgaaa
3121 cccgacagga ctataaagat accaggcgtt tcccctgga agctccctcg tgcgtctcc
3181 tgttccgacc ctgccgtta cggatacct gtccgcctt ctccctcgg gaagcgtggc
3241 gctttctcaa tgctacgct gtaggtatct cagttcggc taggtcgtc gctccaagc
3301 gggctgtgtg cacgaacccc cgttcagcc cgaccgtgc gccttatccg gtaactatc
3361 tcttgagtc aacccgtaa gacacgact atcgccactg gcagcagcca ctgtaacag
3421 gattagcaga gcgaggtat taggcggtgc tacagagttc ttgaagtggg ggcctaacta
3481 cggctacact agaaggacag tatttggtat ctgcgtctg ctgaagccag ttacctcgg
3541 aaaaagagtt ggtagctct gatccggcaa acaaacacc gctggtagcg gtggttttt
3601 tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc cttgatctt
3661 ttctacgggg tctgacgctc agtggaacga aaactcacgt taagggattt tggcatgag
3721 attatcaaaa aggatcttca cctagatcct ttgatcctc cggcgttcag cctgtgccac
3781 agccgacagg atggtgacca ccatttgc ccatacccg tcggtactga tcccgctgc
3841 aataaacga accgctacac cctgagcatc aaactcttt atcagttgga tcattgctgc
3901 ggtgtcgcgg ccaagacggc cgagcttct caccagaatg acatcacct cctccacct
3961 cactctcagc aaatccagc ctcccgatc tttgaactg ccggtgcct tgcggttaa
4021 gatgcgggta gcttttacc ctgcatctt gagcgtgag gtctgcctc tgaagaagg
4081 gttgctgact cataccagc ctgaatgcc ccatcatcca gccagaaag gagggagcca
4141 cgggtgatga gagctttgt gtaggtggc cagttggtga tttgaactt ttgcttgc

4201 acggaacggt ctgcgtgtgc gggaagatgc gtgatctgat ccttcaactc agcaaaagtt
4261 cgatttattc aacaagccg cgtcccgtc aagtcagcgt aatgctctgc cagtgttaca
4321 accaattaac caattctgat tagaaaaact catcgagcat caaatgaaac tgcaatttat
4381 tcatacagg attatcaata ccatattttt gaaaaagccg ttctgtaat gaaggagaaa
4441 actcaccgag gcagttccat aggatggcaa gatcctggta tcggtctgcg attccgactc
4501 gtccaacatc aatacaacct attaatttcc cctcgtcaaa aataagggtta tcaagtgaga
4561 aatcaccatg agtgacgact gaatccggtg agaatggcaa agcttatgc attttttcc
4621 agactgttc aacaggccag ccattacgct cgtcatcaaa atcactcgca tcaacaaaac
4681 cgttattcat tcgtgattgc gcctgagcga gacgaaatac gcgatcgtg taaaaggac
4741 aattacaaac aggaatcgaa tgcaaccggc gcaggaacac tgccagcgca tcaacaatat
4801 ttccacctga atcaggatat tcttctaata cctggaatgc tgtttcccg gggatcgag
4861 tggtagtaaa ccatgcatca tcaggagtac ggataaaatg cttgatggc ggaagaggca
4921 taaatccgt cagccagttt agtctgacca tctcatctgt aacatcattg gcaacgctac
4981 ctttgccatg ttccagaaac aactctggcg catcgggctt cccatacaat cgatagattg
5041 tcgcacctga ttgcccga ttatcgcgag cccattata cccatataaa tcagcatcca
5101 tgttggaatt taatcgggc ctcgagcaag acgtttcccg ttgaatatgg ctcataacac
5161 ccctgtatt actgtttatg taagcagaca gttttattgt tcatgatgat atattttat
5221 cttgtgcaat gtaacatcag agattttgag acacaacgtg gctttgtga ataatcgaa
5281 ctttgctga gttgaaggat cagatcacgc atctcccga caacgcagac cgtccgtgg
5341 caaagcaaaa gttcaaaatc accaactggt ccactacaa caaagctctc atcaaccgtg
5401 gtcctcac tttctggctg gatgatgggg cgattcaggc ctggtatgag tcagcaacac
5461 cttcttcag aggcagacct cagegctaga ttattgaagc attatcagg gttattgtct
5521 catgagcgga tacatatttg aatgtattta gaaaaataaa caaatagggg ttccgcgcac
5581 atttccccga aaagtccac ctgacgt


Expression vector pShuttle-SN, complete sequence

GenBank: AY862402.1


[FASTA](#) [Graphics](#)[Go to:](#) ☐

LOCUS AY862402 5607 bp DNA circular SYN 19-JUL-2005
 DEFINITION Expression vector pShuttle-SN, complete sequence.
 ACCESSION AY862402
 VERSION AY862402.1
 KEYWORDS .
 SOURCE Expression vector pShuttle-SN
 ORGANISM [Expression vector pShuttle-SN](#)
 other sequences; artificial sequences; vectors.
 REFERENCE 1 (bases 1 to 5607)
 AUTHORS Liu,R.Y., Wu,L.Z., Huang,B.J., Huang,J.L., Zhang,Y.L., Ke,M.L.,
 Wang,J.M., Tan,W.P., Zhang,R.H., Chen,H.K., Zeng,Y.X. and Huang,W.
 TITLE Adenoviral expression of a truncated S1 subunit of SARS-CoV spike
 protein results in specific humoral immune responses against
 SARS-CoV in rats
 JOURNAL Virus Res. 112 (1-2), 24-31 (2005)
 PUBMED [16022898](#)
 REFERENCE 2 (bases 1 to 5607)
 AUTHORS Liu,R.-Y., Huang,B.-J., Wu,L.-Z., Huang,J.-L., Zhang,R.-H.,
 Zeng,Y.-X. and Huang,W.
 TITLE Constructing recombinant adenovirus carrying the spike gene
 fragments as a vaccine against SARS-CoV by in vitro ligation
 JOURNAL Unpublished
 REFERENCE 3 (bases 1 to 5607)
 AUTHORS Liu,R.-Y., Huang,B.-J., Wu,L.-Z., Huang,J.-L., Zhang,R.-H.,
 Zeng,Y.-X. and Huang,W.
 TITLE Direct Submission
 JOURNAL Submitted (21-DEC-2004) Cancer Center, Sun Yat-Sen University, 651
 Dongfeng Road East, Guangzhou, Guangdong 510060, China
 FEATURES
 source 1..5607
 /organism="Expression vector pShuttle-SN"
 /mol_type="other DNA"
 /db_xref="taxon:[308969](#)"
 /country="China"
 CDS 990..2507
 /codon_start=1
 /transl_table=11
 /product="truncated SARS coronavirus spike glycoprotein S1
 subunit"
 /protein_id="[AAW56614.1](#)"
 /translation="MFIFLLFLTLTSGSDLDRCTTFDDVQAPNYQTHTSSMRGVYYPD
 EIFRSDTLYLTDQLFLPFYSNVTGFHTINHTFDNPVIFPKDGIYFAATEKSNVVRGWV
 FGSTMNKSQSVIIINNSTNVVIRACNFELCDNPFFAVSKPMGTQTHTMIFDNAFNCT
 FEYISDAFSLDVSEKSGNFKHLREFVFKNKDGLYVYKGYQPIDVVRDLPSGFNTLKP
 IFKLPLGINITNFRILTAFAQDTWGTSAAYFVGYLKPTTFMLKYDENGITIDAV
 DCSQNPLAELKCSVKSFEIDKGIYQTSNFRVVPDGVVRFNPITNLCPPGEVFNATKF
 PSVYAWEGKKISNCVADYSVLVYNSTFFSTFKCYGVSATKLNLCFSNVYADSFVVKGD
 DVRQIAPGQTGVIADYNYKLPPDFMGCVLAWNTRNIDATPTGNVYNYRYLRHGLRSP
 FERDISNVFSPDGKPTPPALNCYWPLNDYGFYTTTGIGTKLKFPLISLDCAF"
 misc_feature 990..2459
 /note="Region: SARS coronavirus spike glycoprotein"
 misc_feature 2460..2507
 /note="derived from pShuttle vector"

Source: <https://www.ncbi.nlm.nih.gov/nuccore/AY862402.1?report=GenBank>



Source: <https://www.nature.com/articles/s41467-020-18077-5>



Adenovirus vaccine sequences in patient specimen WIV02 from patient who is 32 y, male, hospitalized, ICU4G, outbreak 19 Dec 2019.

SRX7730880: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid
1 ILLUMINA (Illumina HiSeq 3000) run: 67.1M spots, 20.1G bases, 12.6Gb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the MGIEasy RNA Library Prep Set (96 RXN) (Cat. No.: 1000006384). Paired-end (150 bp) sequencing of the RNA library was performed on the MGISEQ-2000RS platform .

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads
[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)
[hide Abstract](#)
Discovery and characterization of a novel human coronavirus from five patients at the early stage of the Wuhan seafood market pneumonia virus outbreak .

Sample:
[SAMN14082196](#) • [SRS6151291](#) • [All experiments](#) • [All runs](#)
Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:
Name: WIV02-2
Instrument: Illumina HiSeq 3000
Strategy: RNA-Seq
Source: METAGENOMIC
Selection: RANDOM
Layout: PAIRED

Runs: 1 run, 67.1M spots, 20.1G bases, [12.6Gb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092063	67,083,195	20.1G	12.6Gb	2020-02-16

URL: <https://www.ncbi.nlm.nih.gov/sra/SRX7730880%5baccn%5d>

Adenovirus Expression vector pShuttle-SN, Synthetic construct H7N9 HA gene 7640-9302

Job Title: AY862402:Expression vector pShuttle-SN, Synthetic construct I
RID: [Z3E1GPKR01R](#) Search expires on 01-04 19:27 pm [Download All](#) [v](#)

Program: BLASTN [?](#) [Citation](#) [v](#)

Database: SRA [See details](#) [v](#)

SRA Blast search set information
[SRX7730880](#) [SRR11092063](#)

Query Length: 5711

Other reports: [Distance tree of results](#) [MSA viewer](#) [?](#)

Descriptions **Graphic Summary** Alignments


[hover to see the title](#) [click to show alignments](#)

Alignment Scores: ■ < 40 ■ 40 - 50 ■ 50 - 80 ■ 80 - 200 ■ >= 200 [?](#)


100 sequences selected [?](#)

Distribution of the top 100 Blast Hits on 100 subject sequences

534-3301 contiguous nt sequence (2768 nt) in H7N9 HA gene





Adenovirus with CoV-2 Spike Protein, full sequence



534-4573 contiguous (4040 nts)

Adenovirus with CoV-1 partial sequence

534-1905 (1372 nts) contiguous




Adenovirus vaccine with CoV-2 SP:


Bayesian Analysis of SARS-CoV-2 Origin

Steven C. Quay, MD, PhD


29 January 2021




Adenovirus H7N9




Adenovirus CoV-2




Adenovirus CoV-1



Adenovirus CoV-2



WIV-7 patient blast with Adenovirus to CoV-1



Blast analysis of early RNA seq raw reads from the Wuhan Institute of virology contain extensive reads matching “Expression vector pShuttle-SN” sequences, the same adenovirus vector used by the PLA Army for the creation of a vaccine.

Following the 2003 SARS epidemic, Liu et al. developed an adenoviral expression vector of a truncated S1 subunit of SARS-CoV spike protein that resulted in specific humoral immune responses against SARS-CoV in rats.¹⁴³ This same vector was used to create the CoV-2 adenovirus vector vaccine.¹⁴⁴

In order to test the hypothesis that CoV-2 began in the PLA Hospital as a vaccine challenge clinical trial that went awry, RNA-Seq raw reads from nasopharyngeal specimens of Wuhan COVID patients were blasted against the published genome sequence of the SARS-CoV-1 vaccine (GenBank [AY862402.1](#)). I used the SARS-CoV-1 vaccine because the PLA CoV-2 vaccine has not been published.

Nt Sequence	Function
1-990	Adeno virus genes
991-2506	Truncated N-terminus of SARS-CoV-1 Spike Protein
2507-5607	Adeno virus genes

The expected result would be the finding of RNA-Seq sequence raw reads that were homologous to the two Adenovirus regions but only partially homologous (about 80%) to the SARS-CoV-1 regions.

Eleven entries were found on GenBank of SRA data for RNA-Seq of early COVID-19 patients from Wuhan that were sequenced at either the WIV or the Hubei Provincial Center for Disease Control and Prevention (Hubei CDC). These entries are in the Text-Table below.

¹⁴³ <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7114075/>

¹⁴⁴ Chinese patent, attached herein.

- ☐ [RNA-Seq of Homo sapiens: hCov-19 infected patients Bronchoalveolar lavage uid](#)
- 1. 1 ILLUMINA (Illumina MiSeq) run: 5.8M spots, 1.7G bases, 634.3Mb downloads
Accession: SRX8032203
- ☐ [RNA-Seq of Homo sapiens: hCov-19 infected patients Bronchoalveolar lavage uid](#)
- 2. 1 ILLUMINA (Illumina MiSeq) run: 5.2M spots, 1.6G bases, 583.4Mb downloads
Accession: SRX8032202
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 3. 1 ILLUMINA (Illumina MiSeq) run: 5.2M spots, 1.5G bases, 772.9Mb downloads
Accession: SRX7730887
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 4. 1 ILLUMINA (Illumina MiSeq) run: 5.2M spots, 1.5G bases, 768.3Mb downloads
Accession: SRX7730886
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 5. 1 ILLUMINA (Illumina MiSeq) run: 8.3M spots, 2.2G bases, 1.2Gb downloads
Accession: SRX7730885
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 6. 1 ILLUMINA (Illumina HiSeq 3000) run: 38.5M spots, 11.5G bases, 7.1Gb downloads
Accession: SRX7730884
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 7. 1 ILLUMINA (Illumina HiSeq 3000) run: 29.7M spots, 8.9G bases, 5.6Gb downloads
Accession: SRX7730883
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 8. 1 ILLUMINA (Illumina HiSeq 3000) run: 34.3M spots, 10.3G bases, 6.4Gb downloads
Accession: SRX7730882
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 9. 1 ILLUMINA (Illumina HiSeq 1000) run: 61.3M spots, 18.4G bases, 11.4Gb downloads
Accession: SRX7730881
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 10. 1 ILLUMINA (Illumina HiSeq 3000) run: 67.1M spots, 20.1G bases, 12.6Gb downloads
Accession: SRX7730880
- ☐ [RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid](#)
- 11. 1 ILLUMINA (Illumina MiSeq) run: 3.6M spots, 1G bases, 548.1Mb downloads
Accession: SRX7730879

The WIV entry with the greatest read depth, Number 10 above, is described below:

SRX7730880: RNA-Seq of Homo sapiens: bronchoalveolar lavage fluid
 1 ILLUMINA (Illumina HiSeq 3000) run: 67.1M spots, 20.1G bases, 12.6Gb downloads

Design: Total RNA was extracted from bronchoalveolar lavage fluid using the QIAamp Viral RNA Mini Kit (50) following the manufacturers instructions. An RNA library was then constructed using the MGIEasy RNA Library Prep Set (96 RXN) (Cat. No.: 100006384). Paired-end (150 bp) sequencing of the RNA library was performed on the MGISEQ-2000RS platform .

Submitted by: Wuhan Institute of Virology, Chinese Academy of Sciences

Study: Severe acute respiratory syndrome coronavirus 2 Raw sequence reads
[PRJNA605983](#) • [SRP249613](#) • [All experiments](#) • [All runs](#)
[show Abstract](#)

Sample:
[SAMN14082196](#) • [SRS6151291](#) • [All experiments](#) • [All runs](#)
Organism: [Severe acute respiratory syndrome coronavirus 2](#)

Library:
Name: WIV02-2
Instrument: Illumina HiSeq 3000
Strategy: RNA-Seq
Source: METAGENOMIC
Selection: RANDOM
Layout: PAIRED

Runs: 1 run, 67.1M spots, 20.1G bases, [12.6Gb](#)

Run	# of Spots	# of Bases	Size	Published
SRR11092063	67,083,195	20.1G	12.6Gb	2020-02-16

Unexpectedly, over 100 sequences producing significant alignment were identified:

BLAST » blastn suite-SRA » results for RID-S76CAHY001R

Home Recent Results Saved Strategies Help

[Edit Search](#) [Save Search](#) [Search Summary](#)

How to read this report? [BLAST Help Videos](#) [Back to Traditional Results Page](#)

Filter Results

Percent Identity: to E value: to Query Coverage: to

[Filter](#) [Reset](#)

Job Title: [gb|AY862402.1|](#)

RID: [S76CAHY001R](#) Search expires on 10-13 07:47 am [Download All](#)

Program: BLASTN [Citation](#)

Database: SRA [See details](#)

Query ID: [AY862402.1](#)

Description: Expression vector pShuttle-SN, complete sequence

Molecule type: nucleic acid

Query Length: 5607

Other reports: [Distance tree of results](#) [MSA viewer](#)


[Descriptions](#) [Graphic Summary](#) [Alignments](#)

Sequences producing significant alignments [Download](#) [Manage Columns](#) [Show](#) 100 [?](#)

☒ select all 100 sequences selected [Graphics](#) [Distance tree of results](#)

Description	Max Score	Total Score	Query Cover	E value	Per. Ident	Accession
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.66604450.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.66456076.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.63120099.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.63120099.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.62730385.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.61105636.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.60748776.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.60011402.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.59155252.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.59155252.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.58125883.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.57571550.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.57484454.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.56079039.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.56036194.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.55663455.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.55111993.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.53777284.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.53579813.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.52965281.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.51414706.1
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.51016881.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.50609371.2
<input checked="" type="checkbox"/> SRX7730880	278	278	2%	2e-70	100.00%	SRA-SRR11092063.50609371.1

A graphical display of the alignments shows they are not in the Spike Protein region (961 to 2507) of the adenovirus vector but outside of those regions.



An examination of individual reads shows 100% homology over the entire 150 nt segments and outside of the Spike Protein region. The first set of reads are immediately downstream of the Spike Protein segment. The other read is from the 5' boundary of the Adenovirus vector with the Spike Protein region.

To test if this was the actual SARS-CoV-1 vaccine vector and had been given to the patients as an desperate attempt to create immunity during an infection, the Spike Protein region of the vaccine was blasted against the above sample, looking for a near 100% homology. The only reads were a 38 nt segment of 1482-1518, with one gap, as expected. The absence of long reads for the SARS-CoV-1 Spike Protein establishes that this vaccine was not a CoV-1 vaccine.

To test if the homology seen between lavage specimens of patients in Wuhan with the CoV-1 Adenovirus vaccine was due to homology with human sequences, the Expression vector itself was blasted against *Homo sapien* sequences, but no matches were found, as shown below.

BLAST[®] » blastn suite » results for RID-S793VKCV01R

[< Edit Search](#) [Save Search](#) [Search Summary](#) ▼

i Your results are filtered to match records that include: *Homo sapiens* (taxid:9606)

Job Title	AY862402:Expression vector pShuttle-SN, complete...
RID	S793VKCV01R Search expires on 10-13 08:34 am Download All ▼
Program	? Citation ▼
Database	nt See details ▼
Query ID	AY862402.1
Description	Expression vector pShuttle-SN, complete sequence
Molecule type	nucleic acid
Query Length	5607
Other reports	?

! No significant similarity found. For reasons why [click here](#)